

Specimen Label



Garlon[®] 4

Specialty Herbicide

®Trademark of Dow AgroSciences LLC

For the control of woody plants and annual and perennial broadleaf weeds in non-crop industrial manufacturing and storage sites, rights-of-way such as electrical power lines, communication lines, pipelines, roadsides, railroads, forests and in the establishment and maintenance of wildlife openings. Use on these sites may include application to grazed areas.

Active Ingredient:

triclopyr: 3,5,6-trichloro-2- pyridinyloxyacetic acid, butoxyethyl ester.....	61.6%
Other Ingredients	38.4%
Total.....	100.0%

Contains petroleum distillates
Acid equivalent: triclopyr - 44.3% - 4 lb/gal

EPA Reg. No. 62719-40

Keep Out of Reach of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation • Harmful If Swallowed • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

Applicators and other handlers who handle this pesticide must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables are given, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the WPS (40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

The requirements in this box apply to forestry uses.

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves
- Shoes plus socks
- Protective eyewear

Non-Agricultural Use Requirements

The requirements in this box apply to all use sites on this label except for forestry uses.

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: For applications to non-cropland areas, do not allow entry into areas until sprays have dried.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal. Open dumping is prohibited.

Pesticide Storage: Store above 28°F or agitate before use.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Storage and Disposal (Cont.)

Nonrefillable containers 5 gallons or less:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers 5 gallons or larger:

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable containers 5 gallons or larger:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

General Information

Use Garlon® 4 specialty herbicide for the control of woody plants and annual and perennial broadleaf weeds in non-crop industrial manufacturing and storage sites, rights-of-way such as electrical power lines, communication lines, pipelines, roadsides, railroads, forests and in the establishment and maintenance of wildlife openings. Use on these sites may include application to grazed areas.

Garlon 4 is an oil soluble, emulsifiable liquid product containing the herbicide triclopyr. Garlon 4 may be applied to woody or herbaceous broadleaf plants as a foliar spray or as a basal bark or cut stump application to woody plants. As a foliar spray, Garlon 4 controls only herbaceous plants that have emerged from the soil or woody plants that are in full leaf at the time of application. Small amounts of Garlon 4 can kill or injure many broadleaf plants. To prevent damage to crops and other desirable plants, follow all directions and precautions.

General Use Precautions and Restrictions

In Arizona: The state of Arizona has not approved Garlon 4 for use on plants grown for commercial production; specifically forests grown for commercial timber production, or on designated grazing areas.

Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications:

1. The distance of the outer most operating nozzles on the boom must not exceed 3/4 the length of the rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the following Aerial Drift Reduction Advisory. [This information is advisory in nature and does not supersede mandatory label requirements.]

Aerial Drift Reduction Advisory

Information on Droplet Size: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

Controlling Droplet Size:

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produced larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length: For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height: Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind

Mixing Directions

Garlon 4 may be foliarly applied by diluting with water or by preparing an oil-water emulsion. For woody plant control, an oil-water emulsion performs more dependably under a broader range of conditions than a straight water dilution and is recommended for aerial applications.

Oil-Water Mixture Sprays

Prepare a premix of oil, surfactant and Garlon 4 in a separate container using diesel fuel, fuel oil, or kerosene plus an emulsifier such as Sponto 712 or Triton X-100. Use a jar test to check spray mix compatibility before preparing oil-water emulsion sprays in the mixing tank. Do not allow any water or mixtures containing water to get into the premix or Garlon 4 since a thick "invert" (water in oil) emulsion may form that will be difficult to break. Such an emulsion may also be formed if the premix or Garlon 4 is put into the mixing tank before the addition of water. Fill the spray tank about one-half full with water, then slowly add the premix with continuous agitation and complete filling the tank with water. Continue moderate agitation.

Ground Application: Add oil to the spray mix at a rate of 5 to 10% of the total mix, up to a maximum of 1 gallon of oil per acre, using agricultural spray emulsifiers according to mixing instructions below.

Aerial Application: Use oil and water in the spray mixture in a 1:5 ratio (1 part oil to 5 parts water), up to a maximum of 1 gallon of oil per acre according to mixing instructions below.

Oil Mixture Sprays for Basal Treatment

Prepare oil-based spray mixtures using either diesel fuel, No. 1 or No. 2 fuel oil, kerosene or a commercially available basal oil. Substitute other oils or diluents only as recommended by the oil or diluent's manufacturer. When preparing an oil mixture, read and follow the use directions and precautions on the manufacturer's product label. Add Garlon 4 to the required amount of oil in the spray tank or mixing tank and mix thoroughly. If the mixture stands over 4 hours, reagitiation is required.

Oil Mixtures of Garlon 4 and Tordon K: Tordon K and Garlon 4 may be used in tank mix combination for basal bark treatment of woody plants. These herbicides are incompatible and will not form a stable mixture when mixed together directly in oil. Make a stable tank mixture for basal bark application by first combining each product with a compatibility agent prior to final mixing in the desired ratio. (See product bulletin for mixing instructions.) Tordon K is not registered for use in the states of California and Florida.

Water Dilutions

For water dilutions, an agricultural surfactant at the manufacturer's recommended rate may be added to the spray mixture to provide improved wetting of foliage. To help minimize spray drift, a drift control and deposition aid cleared for application to growing crops is recommended.

Tank Mixing

Garlon 4 may be applied in tank mix combination with labeled rates of other herbicides provided (1) the tank mix product is labeled for the timing and method of application for the use site to be treated; and (2) tank mixing is not prohibited by the label of the tank mix product. When tank mixing Garlon 4 with other materials, a compatibility test (jar test) using relative proportions of the tank mix ingredients should be conducted prior to mixing ingredients in the spray tank. Use a clear glass quart jar with lid and mix the tank mix ingredients in the required order and their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, jels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

Mixing Order for Tank Mixes: Add one-half of the needed water to the mixing tank and start agitation. Add different materials in the order indicated below, allowing time for complete dispersion and mixing after addition of each product.

1. Water soluble herbicide (if used)
2. Premix of oil, emulsifier, Garlon 4 and other oil-soluble herbicide (if used); see below

Add the remaining water. During the final filling of the tank, add a drift control and deposition aid cleared for application to growing crops (if used), plus an agricultural surfactant (if a water dilution rather than an oil-water emulsion spray is used). Maintain continuous agitation of the spray mixture during mixing, final filling and throughout application to ensure spray uniformity.

Premixing: Prepare a premix of oil, emulsifier (if oil-water emulsion), and Garlon 4 plus other oil-soluble herbicide (if used), e.g., 2,4-D ester. **Note:** Do not allow water or mixtures containing water to get into the premix or Garlon 4 since a thick "invert" (water in oil) emulsion may form that will be difficult to break. Such an emulsion may also be formed if the premix or Garlon 4 is put into the mixing tank before the addition of water.

Tank Mixing Precautions:

- Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels.
- Do not exceed recommended application rates. If products containing the same active ingredient are tank mixed, do not exceed the maximum allowable active ingredient use rates.
- For direct injection or other spray equipment where the product formulations will be mixed in undiluted form, special care should be taken to ensure tank mix compatibility.
- Always perform a (jar) test to ensure the compatibility of products to be used in tank mixture.

Mixing with Liquid Fertilizer for Broadleaf Weed Control

Garlon 4 may be tank mixed with liquid nitrogen fertilizer and foliarly applied for weed control and fertilization of grass pastures. Use Garlon 4 in accordance with recommendations for grass pastures as given on this label. Apply at rates recommended by supplier or Extension Service Specialist. **Note:** Garlon 4 is not recommended for use with liquid fertilizer on woody plants (brush). Foliage burn caused by liquid fertilizer may reduce herbicide effectiveness on woody plants. Test for mixing compatibility using desired procedure and spray mix proportions in clear glass jar before mixing in spray tank. A compatibility aid such as Unite or Compex may be needed in some situations. **Compatibility is best with straight liquid nitrogen fertilizer solutions. Mixing with N-P-K solutions or suspensions may not be satisfactory even with the addition of compatibility aid.** Premixing Garlon 4 with 1 to 4 parts water may help in difficult situations.

Fill in the spray tank about half full with the liquid fertilizer, then add the herbicide with agitation and complete filling the tank with fertilizer. Apply immediately and continue agitation in the spray tank during application.

Do not store liquid fertilizer spray mixtures. Application during very cold weather (near freezing) is not advisable. The likelihood of mixing or compatibility problems with liquid fertilizer increases under cold conditions.

Note: Do not use spray equipment for other applications to land planted, or to be planted, to susceptible crops or desirable plants **unless** it has been determined that all phytotoxic herbicide residue has been removed by thoroughly cleaning the equipment.

Plants Controlled by Garlon 4

Woody Plant Species

alder	cottonwood	maple (except bigleaf, vine ³)	sweetbay magnolia
arrowwood	crataegus (hawthorn)	milkweed vine ³	sweetgum
ash	dogwood	mulberry	sycamore
aspen	Douglas-fir	oaks	tanoak
bear clover (bearmat)	elderberry	osage orange	thimbleberry
beech	elm (except winged elm)	pepper vine ³	tree-of-heaven
birch	gallberry	persimmon, eastern	(<i>Ailanthus</i>) ¹
blackberry	gorse	pine	trumpet creeper ³
blackbrush	granjeno	poison ivy	tulip poplar
blackgum	guajillo	poison oak	twisted acacia
boxelder ¹	guava ³	poplar	Virginia creeper ³
Brazilian pepper	hazel	salmonberry	wax myrtle (top growth)
buckthorn	hickory	saltbush	wild rose
cascara	hornbeam	(<i>Braccharis</i> spp.) ³	willow
ceanothus	huisache (suppression)	salt cedar ¹	willow primrose
cherry ³	kudzu ²	sassafras	winged elm
chinquapin	locust	scotch broom	
choke cherry	madrone	sumac	

¹For best control, use either a basal bark or cut stump treatment.

²For complete control, re-treatment may be necessary.

³Basal or dormant stem applications only.

Annual, Biennial and Perennial Broadleaf Weeds

Note: Numbers in parentheses refer to footnotes below table.

black medic	dandelion (top growth)	Oxalis	vetch
bull thistle	dogfennel	plantain	wild carrot
burdock	field bindweed	purple loosestrife	(Queen Anne's lace)
Canada thistle	goldenrod	ragweed	wild lettuce
chicory	ground ivy	sericea lespedeza (1)	wild violet
cinquefoil	lambsquarters	smartweed	yarrow
clover	lespedeza	sulfur cinquefoil (2)	
creeping beggarweed	matchweed	sweet clover	
curly dock	mustard	tropical soda apple (3)	

1. **Sericea lespedeza:** Apply 1 to 2 pints of Garlon 4 per acre. For best results, apply after maximum foliage development in the late spring to early summer, but prior to bloom.
2. **Sulfur cinquefoil:** Apply 1 to 2 pints of Garlon 4 per acre. For best results, apply to plants in the rosette stage.
3. **Tropical soda apple:** Apply 2 pints of Garlon 4 per acre when tropical soda apple plants reach the first flower stage. For best results, apply in a total spray volume of 40 gallons per acre using ground equipment. An agricultural surfactant may be added at the manufacturer's recommended rate to provide more complete wetting and coverage of the foliage. Spot treatments may be used to control sparse plant stands. For spot treatment use a 1 to 1.5% solution of Garlon 4 in water (1 to 1 1/2 gallons of Garlon 4 in 100 gallons total spray mixture) and spray the entire plant to completely wet the foliage. **In Florida,** control of tropical soda apple may be improved by using the following management practices:

- Mow plants to a height of 3 inches every 50 to 60 days or whenever they reach flowering. Continue the mowing operation through April.
- In late May to June (50 to 60 days after the April mowing), apply Garlon 4 as a broadcast treatment.
- Use spot treatment to control any remaining plants or thin stands of plants that germinate following a broadcast treatment.

Application Methods

Use Garlon 4 at rates of 1 to 8 quarts per acre to control broadleaf weeds and woody plants. It is suggested that rates higher in this rate range be used to control woody plants. In all cases, use the amount specified in enough water to give uniform and complete coverage of the plants to be controlled. The order of addition to the spray tank is water, spray thickening agent (if used), surfactant (if used), additional herbicide (if used), and Garlon 4. If a standard agricultural surfactant is used, use at a rate of 1 to 2 quarts per acre. Use continuous adequate agitation.

Before using any recommended tank mixtures, read the directions and all precautions on both labels.

For best results apply when woody plants and weeds are actively growing. When hard to control species such as ash, blackgum, choke cherry, elm, maples (other than vine or big leaf), oaks, pines, or winged elm are prevalent, during applications made during late summer when the plants are mature, or during drought conditions, use the higher rates of Garlon 4 alone or in combination with Tordon® 101 Mixture specialty herbicide or Tordon K herbicide. Tordon 101 Mixture and Tordon K are restricted use pesticides. Tordon 101 Mixture and Tordon K are not registered for use in the states of California and Florida.

When using Garlon 4 in combination with 2,4-D low volatile ester herbicide, generally the higher rates of Garlon 4 should be used for satisfactory brush control.

Use the higher dosage rates when brush approaches an average of 15 feet in height or when the brush covers more than 60% of the area to be treated. If lower rates are used on hard to control species, resprouting may occur the year following treatment.

On sites where easy to control brush species dominate, rates less than those listed may be effective. Consult state or local extension personnel for such information.

Foliage Treatment With Ground Equipment

Use sufficient spray volume to completely and uniformly cover foliage. For ground application, apply 10 gallons or more of total spray volume per acre. Use higher spray volumes for ground applications to ensure adequate coverage with increased depth and density of foliage, particularly for treatment of woody plants.

High Volume Foliage Treatment

For control of woody plants, use Garlon 4 at the rate of 2 to 6 quarts per 100 gallons of spray mixture, or Garlon 4 at 2 to 4 quarts may be tank mixed with labeled rates of 2,4-D low volatile ester herbicide, Tordon 101 Mixture, or Tordon K and diluted to make 100 gallons of spray. Do not apply more than 2 gallons of Garlon 4 per acre. On rangeland and permanent pasture sites, make 1 application per year and apply no more than 2 quarts of Garlon 4 (2 lb ae of triclopyr) per acre. Apply at a volume of 100 to 400 gallons of total spray per acre depending upon size and density of woody plants. Tordon 101 Mixture and Tordon K are not registered for use in the states of California and Florida. When tank mixing, follow applicable use directions and precautions on each manufacturer's label.

Depending upon the size and density of the woody plants, apply sufficient spray volume to thoroughly wet all leaves, stems, and root collars. To minimize spray drift, select the minimum spray pressure that provides adequate plant coverage without forming a mist and direct sprays no higher than the top of the target plants. Use a drift control additive cleared for application to growing crops to reduce spray drift. Before using any tank mixture, read the directions and use precautions on both labels. For best results, apply when woody plants and weeds are actively growing.

Table 1: The following table is provided as a guide to the user to achieve the proper rate of Garlon 4.

Total Spray Volume (gallons/acre)	Rate of Garlon 4	
	Forestry Sites (qt/100 gallons of spray) ¹	Non-Cropland Sites (qt/100 gallons of spray) ²
400	1.5	2
300	2	2.7
200	3	4
100	6	8
50	12	16
40	15	20
30	20	26.7
20	30	40
10	60	80

¹Do not exceed the maximum use rate of 6 qt of Garlon 4 (6 lb ae of triclopyr) per acre per year.

²Do not exceed the maximum use rate of 8 qt of Garlon 4 (8 lb ae of triclopyr) per acre per year for non-grazable areas, or 2 qt (2 lb ae of triclopyr) per acre per year for grazed areas, except on portions of grazed areas that meet the following requirement. Portions of grazed areas that intersect treated non-cropland, rights-of-way and forestry sites may be treated at up to 8 lb ae per acre if the area to be treated on the day of application comprises no more than 10% of the total grazable area.

Table 2

Application Rates per 100 Gallons of Spray		
Garlon 4	Plus Tank Mix Product	Rate (qt)
1 - 4 qt	--	--
1 - 2 qt	Grazon® P+D specialty herbicide	4
1 - 2 pt	2,4-D low volatile ester herbicide	1 - 2
1 - 2 qt	Tordon 22K	1 - 2
2 qt	Reclaim® specialty herbicide ^{1,2}	2

¹Reclaim is registered for use only in Arizona, Texas, Oklahoma and New Mexico.

²See directions for Mesquite Control Using High Volume Foliage Treatment below.

Mesquite Control Using High Volume Foliage Treatment: For control of mesquite infestations of low to moderate density, apply Garlon 4 and Reclaim in a tank mixture to individual plants with backpack or hand-held sprayers or a vehicle-mounted sprayer with hand-held spray wand or spray gun. For individual plant treatment, use 2 quarts of Garlon 4 in combination with 2 quarts of Reclaim per 100 gallons of total spray solution (1/2% v/v of each product). Apply in water or as an oil-water emulsion as described in Mixing Directions. If using an oil-water emulsion, add the oil at a rate of 5% of the total spray volume. Apply as a complete spray-to-wet foliar application, including all leaves. Thorough coverage is necessary for good results, but do not spray to the point of runoff. Do not apply when mesquite foliage is wet. The total amount of Garlon 4 applied should not exceed 1 1/3 pints per acre. For best results, follow information given elsewhere in this label concerning effect of environmental conditions and application timing on control. This application method works best for brush less than 8 feet tall since efficient treatment and thorough coverage of taller brush is difficult to achieve with this method. To minimize drift, select a spray nozzle and pressure that provides good coverage while forming a coarse spray. Additionally, drift may be reduced by using the minimum pressure necessary to obtain plant coverage without forming a mist and by directing sprays no higher than the top of target plants. If desired, a spray dye may be added to the spray mixture to mark the treated plants.

Low Volume Foliage Treatment

To control susceptible woody plants, mix up to 20 quarts of Garlon 4 in 10 to 100 gallons of finished spray. The spray concentration of Garlon 4 and total spray volume per acre should be adjusted according to the size and density of target woody plants and kind of spray equipment used. With low volume sprays, use sufficient spray volume to obtain uniform coverage of target plants including the surfaces of all foliage, stems, and root collars (see General Use Precautions and Restrictions). For best results, a surfactant should be added to all spray mixtures. Match equipment and delivery rate of spray nozzles to height and density of woody plants. When treating tall, dense brush, a truck mounted spray gun with spray tips that deliver up to 2 gallons per minute at 40 to 60 psi may be required. Backpack or other types of specialized spray equipment with spray tips that deliver less than 1 gallon of spray per minute may be appropriate for short, low to moderate density brush.

Tank Mixing: As a low volume foliage spray, up to 12 quarts of Garlon 4 may be applied in tank mix combination with labeled rates of Tordon K or Tordon 101 Mixture in 10 to 100 gallons of finished spray. Tordon 101 Mixture and Tordon K are not registered for use in the states of California and Florida.

Broadcast Applications With Aerial or Ground Equipment

Environmental conditions and application timing influence brush and weed control results. For best results, apply when woody plants and weeds are actively growing. For woody species, apply after the rapid growth period of early spring when leaf tissue is fully expanded and terminal growth has slowed. Brush regrowth should be at least 4 ft high prior to treatment to insure adequate foliage for herbicide absorption. Adequate soil moisture before and after treatment as well as the presence of healthy foliage at the time of application are important factors contributing to optimal herbicidal activity.

Use sufficient spray volume to completely and uniformly cover foliage. For ground application, apply 10 gallons or more of total spray volume per acre. For aerial application, apply at least 2 gallons of total spray volume per acre. Use higher spray volumes for ground or aerial applications to ensure adequate coverage with increased depth and density of foliage, particularly for treatment of woody plants.

Mesquite: The herbicidal response of mesquite is strongly influenced by foliage condition, growth stage and environmental conditions. For best results, apply when new growth foliage has turned from light to dark green, when the soil temperature is above 75°F at a depth of 12 to 18 inches, and soil moisture is adequate for plant growth. Apply within 60 days after the 75°F minimum soil temperature at the 12- to 18-inch depth has been reached. Product performance may be adversely affected if application is made before mesquite foliage has turned from light to dark green or if foliage has been injured or removed by late frost, insects, hail or plant diseases. Do not treat if mesquite exhibits new (light green) terminal growth in response to recent heavy rainfall during the growing season. Rate of soil warm-up at the 12- to 18-inch depth may vary with soil texture and drainage. Coarse-textured (sandy) soils warm up sooner than fine-textured (clay) soils and dry soils warm up more quickly than wet soils. Mesquite regrowth should be at least 4 ft high prior to treatment to insure adequate foliage for herbicide absorption.

Mesquite Only

Apply 1/2 to 1 pint of Garlon 4 per acre in combination with 2/3 to 1 1/3 pint per acre of Reclaim. See label for Reclaim for additional treatment recommendations and information on mesquite control. Apply aerially as an oil:water emulsion in 4 gallons or more total volume per acre or with ground equipment in 10 gallons or more total volume per acre. Use a maximum of 1 gallon of oil per acre for aerial or ground application.

Mesquite and Pricklypear Cactus

If pricklypear cactus is a target species in association with mesquite, apply a tank mix of 1/2 to 1 pint of Garlon 4 with 1 to 2 pints of Tordon 22K per acre. (The 2 pint per acre rate of Tordon 22K provides a higher and more uniform plant kill of pricklypear.) Tordon 22K may also be applied in combination with Reclaim to control pricklypear while providing improved control of mesquite. See labels for Tordon 22K and Reclaim for additional information and treatment recommendations. Apply aerially as an oil:water emulsion in 4 gallons or more total volume per acre or with ground equipment in 10 or gallons or more total volume per acre. If mesquite canopy is dense, use higher spray volumes. Use a maximum of 1 gallon of oil per acre for aerial or ground application.

South Texas Mixed Brush (Mesquite, Pricklypear Cactus, Blackbrush, Twisted Acacia and Granjeno)

Use 1 to 2 pints of Garlon 4 in a tank mix with 2 pints of Tordon 22K per acre if pricklypear is a problem, or with 2/3 to 1 1/3 pints of Reclaim per acre if mesquite is the prevalent species. Garlon 4 contributes to the control of non-legume species such as granjeno and oaks. However, if woody legume species are predominate, apply 2 pints of Tordon 22K per acre in combination with 2/3 to 1 1/3 pints of Reclaim per acre for improved control. See labels for Tordon 22K and Reclaim for additional information and treatment recommendations. Apply aerially in an oil:water emulsion in 4 gallons or more total volume per acre or with ground equipment in 15 gallons or more total volume per acre. Use a maximum of 1 gallon of oil per acre for aerial or ground application. The use of an oil:water emulsion is critical and good spray coverage is essential for acceptable brush control.

Sand Shinnery Oak Suppression

In Texas, New Mexico and Oklahoma, apply Garlon 4 alone at a rate of 1/2 to 2 pints per acre for suppression of shinnery oak growing on sandy soils. Grass response following suppression may be impressive where rainfall is adequate. Grazing deferment following application together with proper grazing management is recommended to allow for the reestablishment of grass stands.

Post Oak and Blackjack Oak - Regrowth Stands

Apply in the late spring (May) to early summer (June-July) when oak leaves are fully developed (expanded). Use 2 quarts of Garlon 4 alone or in tank mix combination with 0.5 to 1 pints of 2,4-D low-volatile ester herbicide per acre. Apply in an oil:water emulsion or water surfactant dilution in sufficient total volume per acre to assure thorough coverage, usually 5 gallons or more per acre by fixed-wing aircraft or helicopter or 15 to 25 gallons per acre by ground equipment. Use a maximum of 1 gallon of oil per acre for aerial or ground application. Lower rates may be used for suppression only. Control will require at least 3 consecutive treatments. **Note:** Regrowth plants have a large root mass relative to top growth when compared to undisturbed plants. In order for top growth to intercept and translocate enough herbicide to control the roots, delay broadcast treatment until top growth is at least 4 ft tall.

High Volume Foliage Treatment: For regrowth less than 4 ft tall, apply 2 quarts of Garlon 4 per 100 gallons of water and 2 quarts of ag surfactant alone or in tank mix combination with 1 gallon of Grazon P+D or 1 quart of Tordon 22K. Apply as a high volume leaf-stem treatment to individual plants using ground equipment.

Post Oak and Blackjack Oak - Mature Stands

For control of mature stands (greater than 5 ft tall), apply 2 quarts of Garlon 4 per acre in late spring (May) to early summer (June-July) when oak leaves are fully developed (expanded). Understory species such as winged elm, buckbrush, tree huckleberry and ash occurring in some areas will not be controlled (only suppressed or defoliated) by using Garlon 4 alone. Where these understory species occur, control may be improved by tank mixing 2 quarts of Garlon 4 with 1 quart of Tordon 22K or 4 quarts of Grazon P+D per acre. For best results, apply as an oil:water emulsion in a total volume of 5 gallons per acre or more by fixed-wing aircraft or helicopter.

Other Susceptible Woody Plants

Apply 2 to 4 pints of Garlon 4 alone or in combination with 2 to 3 quarts of 3.8 lb/gal 2,4-D low volatile ester or amine formulation per acre. If difficult to control species such as ash, choke cherry, elm, maple or oaks are prevalent, and during applications made when plants are mature late in the summer or during drought conditions, use the higher rates of Garlon 4, alone or with 2,4-D. Garlon 4 may also be applied in a tank mixture with Grazon P+D or Tordon 22K for increased control of certain species. See labels for Grazon P+D and Tordon 22K for additional information and treatment recommendations. Apply aerially in 4 gallons or more total volume per acre or with ground equipment in 10 gallons or more total volume per acre. For best results on blackberry, apply during or after bloom. For management of kudzu, apply 1 quart of Garlon 4 per acre. Repeat application may be necessary to achieve desired level of control.

Susceptible Broadleaf Weeds

Use 2 pints of Garlon 4 per acre in a water spray. Apply as a broadcast spray in a total volume of 10 gallons or more per acre by ground equipment or aerially in a total volume of 2 gallons or more per acre. Apply anytime the weeds are actively growing. Garlon 4 at 1/2 to 3 pints may be tank mixed with 1 to 2 quarts of 3.8 lb/gal 2,4-D amine or low volatile ester.

Woody Plant Control

Foliage Treatment: Use 4 to 8 quarts of Garlon 4 in enough water to make 5 gallons or more per acre of total spray, or 1 1/2 to 3 quarts of Garlon 4 may be combined with labeled rates of 2,4-D low volatile ester, Tordon 101 Mixture, or Tordon K in sufficient water to make 5 gallons or more per acre of total spray. Tordon 101 Mixture and Tordon K are not registered for use in the states of California and Florida.

Broadleaf Weed Control

Use Garlon 4 at rates of 1 to 4 quarts in a total volume of 5 gallons or more per acre as a water spray mixture. Apply anytime weeds are actively growing. Garlon 4 at 0.25 to 3 quarts may be tank mixed with labeled rates of 2,4-D amine or low volatile ester, Tordon K, or Tordon 101 Mixture to improve the spectrum of activity. For thickened (high viscosity) spray mixtures, Garlon 4 can be mixed with diesel oil or other inverting agent. When using an inverting agent, read and follow the use directions and precautions on the product label. Tordon 101 Mixture and Tordon K are not registered for use in the states of California and Florida.

Foliage Treatment (Utility and Pipeline Rights-of-Way)

Use 4 to 8 quarts of Garlon 4 alone, or 3 to 4 quarts of Garlon 4 in a tank mix combination with labeled rates of 2,4-D low volatile ester, Tordon 101 Mixture or Tordon K and apply in a total spray volume of 10 to 30 gallons per acre. Use the higher rates and volumes when plants are dense or under drought conditions. Tordon 101 Mixture and Tordon K are not registered for use in the states of California and Florida.

Portions of grazed areas that intersect treated non-cropland, rights-of-way and forestry sites may be treated at up to 8 lb ae per acre if the area to be treated on the day of application comprises no more than 10% of the total grazable area.

Basal Bark, Dormant Stem and Cut Surface Treatments

Individual plant treatments such as basal bark and cut surface applications may be used on any use site listed on this label at a maximum use rate of 8 lb ae of triclopyr per acre. These types of applications are made directly to ungrazed parts of plants and, therefore, are not restricted by the grazing maximum rate of 2 lb ae of triclopyr per acre.

Basal Bark Treatment

To control susceptible woody plants with stems less than 6 inches in basal diameter, mix 1 to 5 gallons of Garlon 4 in enough oil to make 100 gallons of spray mixture. Apply with knapsack sprayer or power spraying equipment using low pressure (20 to 40 psi). Spray the basal parts of brush and tree trunks to a height of 12 to 15 inches from the ground, thoroughly wetting the indicated area. Spray until runoff at the ground line is noticeable. Old or rough bark requires more spray than smooth young bark. Apply anytime, including the winter months, except when snow or water prevent spraying to the ground line. **Mixing with oil requires vigorous agitation to form an oil solution.** Once a solution is formed it will stay stable.

Low Volume Basal Bark Treatment

To control susceptible woody plants with stems less than 6 inches in basal diameter, mix 20 to 30 gallons of Garlon 4 in enough oil to make 100 gallons of spray mixture. Apply with a backpack or knapsack sprayer using low pressure and a solid cone or flat fan nozzle. Spray the basal parts of brush and tree trunks to a height of 12 to 15 inches from the ground in a manner that thoroughly wets the lower stems, including the root collar area, but not to the point of runoff. Herbicide concentration should vary with size and susceptibility of species treated. Apply anytime, including the winter months, except when snow or water prevent spraying to the ground line or when stem surfaces are saturated with water. See Table 1 for relationship between mixing rate, spray volume and maximum application rate. **Note:** The addition of a soil active herbicide to a basal bark mixture with Garlon 4 may result in damage to surrounding non-target vegetation. Care should be taken to assess the areas in which these soil active herbicides are used in combination with Garlon 4 in basal bark applications. **Mixing with oil requires vigorous agitation to form an oil solution.** Once a solution is formed it will stay stable.

Garlon 4 Plus Tordon K in Oil Tank Mix: Garlon 4 and Tordon K may be used in tank mix combination as a low volume basal bark treatment to improve control of certain woody species such as ash, elm, maple, poplar, aspen, hackberry, oak, oceanspray, birch, hickory, pine, tanoak, cherry, locust, sassafras, and multiflora rose. (See product bulletin for mixing instructions.) Tordon K is not registered for use in the states of California and Florida.

Streamline Basal Bark Treatment (Southern States)

To control or suppress susceptible woody plants for conifer release, mix 20 to 30 gallons of Garlon 4 in enough oil to make 100 gallons of spray mixture. Streamline basal bark treatments are most effective on stems less than 4 inches in basal diameter. Apply with a backpack or knapsack sprayer using equipment that provides a directed straight stream spray. Apply the spray in a 2- to 3-inch wide band to one side of stems less than 3 inches in basal diameter. When the optimum amount of spray mixture is applied, the treated zone should widen to encircle the stem within approximately 30 minutes. Treat both sides of stems which are 3 to 4 inches in basal diameter. Direct the spray at bark that is approximately 12 to 24 inches above ground. Pines (loblolly, slash, shortleaf, and Virginia) up to 2 inches in diameter breast height (dbh) can be controlled by directing the spray at a point approximately 4 feet above ground. Vary spray mixture concentration with size and susceptibility of the species being treated. Better control is achieved when spray is applied to thin juvenile bark and above rough thickened mature bark. This technique is not recommended for scrub and live oak species, including blackjack, turkey, post, live, bluejack and laurel oaks, or bigleaf maple. Apply anytime, including winter months, except when snow or water prevents spraying at the desired height above ground level. **Note:** Best results with some hardwood species occur when applications are made from approximately 6 weeks prior to leaf expansion in the spring until approximately 2 months after leaf expansion is completed. **Mixing with oil requires vigorous agitation to form an oil solution.** Once a solution is formed it will stay stable.

Low Volume Stem Bark Band Treatment (North Central and Lake States)

To control susceptible woody plants with stems less than 6 inches in basal diameter, mix 20 to 30 gallons of Garlon 4 in enough oil to make 100 gallons of spray mixture. Apply with a backpack or knapsack sprayer using low pressure and a solid cone or flat fan nozzle. Apply the spray in a 6- to 10-inch wide band that completely encircles the stem. Spray in a manner that completely wets the bark, but not to the point of runoff. The treatment band may be positioned at any height up to the first major branch. For best results, apply the band as low as possible. Spray mixture concentration should vary with size and susceptibility of species to be treated. Applications may be made anytime, including winter months. **Mixing with oil requires vigorous agitation to form an oil solution.** Once a solution is formed it will stay stable.

Thinline Basal Bark Treatment

To control susceptible woody plants with stems less than 6 inches in diameter, apply Garlon 4, either undiluted or mixed at 50 to 75% v/v with oil, in a thin stream to all sides of the lower stems. The stream should be directed horizontally to apply a narrow band of Garlon 4 around each stem or clump. Use a minimum of 2 to 15 milliliters of Garlon 4 or oil mixture with Garlon 4 to treat single stems and from 25 to 100 milliliters to treat clumps of stems. Use an applicator metered or calibrated to deliver the small amounts required. **Mixing with oil requires vigorous agitation to form an oil solution.** Once a solution is formed it will stay stable.

Dormant Stem Treatment

Dormant stem treatments control susceptible woody plants and vines with stems less than 2 inches in diameter. Plants with stems greater than 2 inches in diameter may not be controlled and resprouting may occur. This treatment method is best suited for sites with dense, small diameter brush. Dormant stem treatments of Garlon 4 can also be used as a chemical side-trim for controlling lateral branches of larger trees that encroach onto roadside, utility, or other rights-of-way.

Mix 4 to 8 quarts of Garlon 4 in 2 to 3 gallons of crop oil concentrate or other recommended oil and add this mixture in enough water to make 100 gallons of spray solution. Use continuous adequate agitation. Apply with knapsack or power spraying equipment, using low pressure (20 to 40 psi). In western states, apply anytime after woody plants are dormant and most of the foliage has dropped. In other areas apply anytime within 10 weeks of budbreak, generally February through April. Thoroughly wet the upper parts of the stems and use the remainder to wet the lower 12 to 15 inches above the ground to the point of runoff. For root suckering species such as sumac, sassafras and locust, also spray the ground under the plant to cover small root suckers which may not be visible above the soil surface. For oil-water mixture application, mix 6 quarts of Garlon 4, 25 gallons of oil and 1.5 gallons of an approved agricultural spray emulsifier such as Sponto 712 or Triton X-100 as indicated in the mixing directions. Treat as above. Garlon 4 may be mixed with 4 quarts of Weedone 170 herbicide to improve the control of black cherry and broaden the spectrum of herbicidal activity. Do not apply to wet or saturated bark as poor control may result.

Cut Stump Treatment

To control resprouting, mix 20 to 30 gallons of Garlon 4 in enough oil to make 100 gallons of spray mixture. Apply with a backpack or knapsack sprayer using low pressures and a solid cone or flat fan nozzle. Spray the root collar area, sides of the stump, and the outer portion of the cut surface, including the cambium, until thoroughly wet, but not to the point of runoff. Spray mixture concentration should vary with the size and susceptibility of species treated. Apply anytime, including in winter months, except when snow or water prevent spraying to the ground line.

Mixing with oil requires vigorous agitation to form an oil solution. Once a solution is formed it will stay stable.

Cut Stump Treatment in Western States

To control resprouting of salt cedar and other *Tamarix* species, bigleaf maple, tanoak, Oregon myrtle, and other susceptible species, apply undiluted Garlon 4 to wet the cambium and adjacent wood around the entire circumference of the cut stump. Treatments may be applied throughout the year; however, control may be reduced with treatment during periods of moisture stress as in late summer. Cut stumps so that they are approximately level to facilitate uniform coverage of Garlon 4. Use an applicator which can be calibrated to deliver the small amounts of material required.

Growing Point and Leaf Base (Crown) Treatment of Yucca

Prepare a 2% v/v solution of Garlon 4 in diesel or fuel oil (13 fl oz of Garlon 4 in 5 gallons of spray mixture). Thoroughly wet the center of the plant including growing point and leaf bases to the soil surface. Complete coverage of leaves is not necessary.

Forest Management Applications

For broadcast applications, apply 1 to 6 quarts of Garlon 4 per acre in a total spray volume of 5 to 25 gallons per acre by air or 10 to 100 gallons per acre by ground. Use spray volumes sufficient to provide thorough coverage of treated foliage. Nozzles or additives that produce larger droplets of spray may require higher spray volumes to provide adequate coverage.

Plant Back Interval for Conifers: Conifers planted sooner than 1 month after treatment with Garlon 4 at less than 4 quarts per acre or sooner than 2 months after treatment at 4 to 6 quarts per acre may be injured. When tank mixtures of herbicides are used for forest site preparation, labels for all products in the mixture should be consulted and the longest recommended waiting period before planting observed.

Forest Site Preparation (Not for Conifer Release)

Southern States including Alabama, Arkansas, Delaware, Florida, Georgia, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, and Virginia: To control susceptible woody plants and broadleaf weeds, apply Garlon 4 at a rate of 4 to 6 quarts per acre. To broaden the spectrum of woody plants and broadleaf weeds controlled, apply 2 to 4 quarts of Garlon 4 per acre in tank mix combination with labeled rates of Tordon 101 Mixture or Tordon K. Tordon 101 Mixture and Tordon K are not registered for use in the state of Florida. Where grass control is also desired, Garlon 4, alone or in tank mix combination with Tordon K or Tordon 101 Mixture, may be applied with labeled rates of other herbicides registered for grass control in forests. Use of tank mix products must be in accordance with the most restrictive of label limitations and precautions. Do not exceed labeled application rates. Garlon 4 cannot be tank mixed with any product containing a label prohibition against such mixing.

Western, Northeastern, North Central, and Lake States (States not Listed Above as Southern States): To control susceptible woody plants and broadleaf weeds, apply Garlon 4 at a rate of 3 to 6 quarts per acre. To broaden the spectrum of woody plants and broadleaf weeds controlled, apply 1.5 to 3 quarts of Garlon 4 per acre in tank mix combination with labeled rates of Tordon 101 Mixture, Tordon K, or 2,4-D low volatile ester. Tordon 101 Mixture and Tordon K are not registered for use in the state of California. Where grass control is also desired, Garlon 4, alone or in tank mix combination with Tordon 101 Mixture or Tordon K, may be applied with labeled rates of other herbicides registered for grass control in forests. When applying tank mixes, follow applicable use directions and precautions on each product label.

Southern Coastal Flatwoods: To control susceptible broadleaf weeds and woody species such as gallberry and wax-myrtle, and for partial control of saw-palmetto, apply 2 to 4 quarts of Garlon 4 per acre. To broaden the spectrum of species controlled to include fetterbush, staggerbush, titi, and grasses, apply 2 to 3 quarts of Garlon 4 per acre in tank mix combination with labeled rates of Arsenal Applicator's Concentrate herbicide. Where control of gallberry, wax-myrtle, broadleaf weeds, and grasses is desired, apply 2 to 3 quarts of Garlon 4 per acre in tank mix combination with labeled rates of Accord Concentrate or Accord SP herbicide.

These treatments may be broadcast during site preparation of flat planted or bedded sites or, on bedded sites, applied in bands over the top of beds. For best results, apply in late summer or fall. Efficacy may not be satisfactory when applications are made in early season prior to August.

Note: Do not apply after planting pines.

Directed Spray Applications for Conifer Release

To release conifers from competing hardwoods and brush such as red maple, sugar maple, striped maple, sweetgum, red and white oaks, ash, hickory, alder, birch, aspen, pin cherry, *Ceanothus* spp., blackberry, chinquapin, and poison oak, mix 4 to 20 quarts of Garlon 4 in enough water to make 100 gallons of spray mixture. This spray mixture should be directed onto foliage of competitive hardwoods using knapsack or backpack sprayers with flat fan nozzles or equivalent anytime after the hardwoods and brush have reached full leaf size, but before autumn coloration. The majority of treated hardwoods and brush should be less than 6 feet in height to ensure adequate spray coverage. Care should be taken to direct spray away from contact with conifer foliage, particularly foliage of desirable pines. See Table 1 for relationship between mixing rate, spray volume and maximum application rate.

Note: Spray may cause temporary damage and growth suppression where contact with conifers occurs; however, injured conifers should recover and grow normally. Over-the-top spray applications can kill pines.

Broadcast Applications for Mid-Rotation Understory Brush Control in Southern Coastal Flatwoods Pine Stands (Ground Equipment Only)

For control of susceptible species such as gallberry and wax-myrtle and broadleaf weeds, apply 2 to 4 quarts of Garlon 4 per acre. To broaden the spectrum of woody plants controlled to include fetterbush, staggerbush, and titi, apply 2 to 3 quarts of Garlon 4 per acre in tank mix combination with labeled rates of Arsenal Applicator's Concentrate. Saw-palmetto will be partially controlled by use of Garlon 4 at 4 quarts per acre or by mixtures of Garlon 4 at 2 to 3 quarts per acre in tank mix combination with either Arsenal Applicator's Concentrate or Escort herbicide. These mixtures should be broadcast applied over target understory brush species, **but to prevent injury to pines, make applications underneath the foliage of pines.** Apply sprays in 30 gallons or more per acre of total volume. For best results, apply in late summer or fall. Efficacy may not be satisfactory when applications are made in early season prior to August.

Broadcast Applications for Conifer Release in the Pacific Northwest and California

Dormant Conifers Before Bud Swell (Excluding Pines): To control or suppress deciduous hardwoods such as vine maple, bigleaf maple, alder, scotch broom, or willow **before leaf-out**, or evergreen hardwoods such as madrone, chinquapin, and *Ceanothus* spp., use Garlon 4 at 1 to 2 quarts per acre. Use diesel or fuel oil as a diluent, or use water plus 1 to 2 gallons per acre of diesel oil or a suitable surfactant or oil substitute at manufacturer's recommended rates. **Mixing with oil as the only diluent requires vigorous agitation to form an oil solution.** Once a solution is formed it will stay stable.

Conifer Plantations (Excluding Pines) After Hardwoods Begin Growth and Before Conifer Bud Break ("Early Foliar" Hardwood Stage): Use Garlon 4 at 1 to 1.5 quarts alone or with 2,4-D low volatile ester herbicide in water carrier to provide no more than 3 lb ae per acre from both products. After conifer bud break, these sprays may cause more serious injury to the crop trees. Use of a surfactant may cause unacceptable injury to conifers especially after bud break.

Conifer Plantations (Excluding Pines) After Conifers Harden Off in Late Summer and While Hardwoods are Still Actively Growing: Use Garlon 4 at rates of 1 to 1.5 quarts per acre alone or with 2,4-D low volatile ester to provide no more than 3 lb ae per acre from both products. Treat as soon after conifer bud hardening as possible so that hardwoods and brush are actively growing. Use of oil, oil substitute, or surfactant may cause unacceptable injury to the conifers.

Broadcast Applications for Conifer Release in the Eastern United States

To release spruce, fir, red pine, and white pine from competing hardwoods such as red maple, sugar maple, striped maple, alder, birch (white, yellow, and grey), aspen, ash, pin cherry, and *Rubus* spp. and perennial and annual broadleaf weeds, use Garlon 4 at rates of 1.5 to 3 quarts per acre alone or with 2,4-D amine or low volatile ester to provide no more than 4 lb ae per acre from both products. Apply in late summer or early fall after conifers have formed their overwintering buds and hardwoods are in full leaf and prior to autumn coloration.

Broadcast Applications for Conifer Release in the Lake States Region

To release spruce, fir, and red pine from competing hardwoods such as aspen, birch, maple, cherry, willow, oak, hazel, and *Rubus* spp. and perennial and annual broadleaf weeds, use Garlon 4 at rates of 1.5 to 3 quarts per acre. Apply in late summer or early fall after conifers have formed their overwintering buds and hardwoods are in full leaf and prior to autumn coloration.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC

Produced for Dow AgroSciences LLC • Indianapolis, IN 46268 USA

Label Code: D02-102-026

Replaces Label: D02-102-025

LOES Number: 010-00085

EPA accepted 04/18/07

Revisions:

1. Product may be applied by fixed wing aircraft or helicopter.
2. Added mixing directions section.
3. Added blackbrush, granjeno, guajillo, guava, milkweed vine, osage orange, pepper vine, trumpet creeper, twisted acacia, Virginia creeper and willow primrose to list of woody plants controlled.
4. Added biennial broadleaf weeds to list of weeds controlled.
5. Added dormant stem and cut surface treatments.