


2015 Aerial Insect and Disease Survey

USGS 100K Quad: SKYKOMISH RIVER - E147121; 4C



Mortality Agents		Defoliators	
Code	Damaging Agent	Code	Damaging Agent
1	Douglas-fir beetle	BS	W spruce budworm
4	Pine engraver		
6L	Mtn. pine beetle		
BEAR	Bear		

Other Damaging Agents	
Code	Damaging Agent
AB	Balsam woolly adelgid
DH	Dying hemlock
FIRE	Fire
HDM	Hardwood decline
WATR	Water damage

USGS 100K Quad: SKYKOMISH RIVER - E147121; 4C
2015 Aerial Insect and Disease Survey
Map Scale: 1:100,000
Date: 24 November 2015

Legend

- Defoliating Agents
- Mortality Agents
- Other Damage
- WDNR Managed Lands
- Areas Not Flown During Overview Survey
- 2015 Large Fires

Source: USGS GeoMAC - <http://www.geomac.gov/>

Source: Washington Dept. of Natural Resources

How the Aerial Surveys Are Conducted

Data represented on this map are based on trees visibly affected by forest insects and diseases detected and recorded during aerial survey flights conducted by the USDA Forest Service, the Washington Department of Natural Resources and the Oregon Department of Forestry. Observers have just a few seconds to recognize the color difference between healthy and damaged trees of different species; diagnose causal agents correctly; estimate intensity; delineate the extent of damage; and precisely record this information on a georeferenced, digital map. Air turbulence, cloud shadows, distance from aircraft, haze, smoke and observer experience can all affect the quality of the survey. These data summaries provide an estimate of conditions on the ground and may differ from estimates derived by other methods.

The aerial survey provides information on the current status for many causal agents and is important when examining insect activity trends by comparing historical and current survey data over large areas.

Overview surveys are a 'snap shot' in time and therefore may not be timed to accurately capture the true extent or severity of a particular disturbance activity. Specially designed surveys with modified flight patterns and timing may be conducted to more accurately delineate the extent and severity of a particular disturbance agent. Special surveys, such as Swiss needle cast surveys, are conducted when resources are available to address situations of sufficient economic, political or environmental importance.

DIRECT ALL INQUIRIES TO:

Washington State Department of Natural Resources
 Wildfire Division
 Forest Health
 1111 Washington St. SE
 Olympia, WA 98504

-- OR --

USDA Forest Service, Region 6
 State and Private Forestry
 Forest Health Protection
 PO Box 3623
 Portland, Oregon 97208

DISCLAIMER
 Forest Health Protection (FHP), Washington Department of Natural Resources (WDNR) and Oregon Department of Forestry (ODF) serve to maintain an accurate Aerial Detection Survey (ADS) Dataset, but due to the conditions under which the data are collected FHP, WDNR and ODF shall not be held responsible for missing or inaccurate data. ADS are not intended to replace more specific information. An accuracy assessment has not been done for this dataset; however, ground checks are completed in accordance with local and national guidelines. <http://www.fs.fed.us/foresthealth/visibility/quality/insurance.shtml>. Maps and data may be updated without notice. Please cite: "USDA Forest Service, Forest Health Protection, Washington Department of Natural Resources, Wildfire Division, Forest Health, and Oregon Department of Forestry, Forest Health Management" as the source of this data.

The basemap is an ESRI map service of TOPOI data (Copyright 2013 National Geographic) which is comprised of seamless, scanned images of USGS paper topographic maps. For more info on this map see http://globe.arcgisonline.com/maps/USA_Topo_Maps.

A data dictionary, digital copies of this map and geospatial insect and disease data are available at: <http://www.fs.usda.gov/detail/r6/forest-grasslandhealth/insects-diseases/>

