CHAPTER 2. EXISTING VEGETATION – FOREST SCALE

North Fork Valley

Current Vegetation

Vegetation mapping for the Grand Mesa, Uncompahgre and Gunnison (GMUG) National Forests is generated primarily from aerial photo interpretation, with periodic updates resulting from field verification, management activities and natural disturbances (i.e., wildfires). Vegetation is classified by cover type. Cover type is determined by the dominant cover or species present at the time of classification. As a result, cover type classifications are more simplified than conditions that actually occur on the ground. (For example: Spruce/fir cover type includes areas where Englemann spruce and/or subalpine fir tree species are the majority of the vegetation. Aspen tree species may or may not be present as subdominant species.)

Current vegetation cover type composition on NFS lands within the North Fork Valley Geographic Area is shown in Figure 1. A map of current vegetation is shown in Figure 2.

Figure 1. Composition of Current Vegetation Cover Types on NFS lands within the North Fork Valley Geographic Area



Figure 2. Map of Current Vegetation, North Fork Valley Geographic Area



Habitat Structural Stages

Vegetation is also characterized by structure. Structure is described by habitat structural stages. Habitat structural stages are defined by size class, tree diameter, and canopy closure (or canopy density measured as crown cover percent). Table 1 displays habitat structural stage definitions (Hoover and Wills, 1987).

Table 1. Habitat Structural Stage Definitions

Habitat Structural Stage	Size Class	Diameter	Crown Cover Percent
1T[1]/1M[2]	Grass-Forb	Not applicable	0 – 10%
2T[1]/2S[3]	Shrub-Seedling	< 1 inch	0 - 10%
3A	Sapling-Pole	1 – 9 inches	11 – 40%
3B	Sapling-Pole	1 – 9 inches	41 - 70%
3C	Sapling-Pole	1 – 9 inches	71 - 100%
4A	Mature	9+ inches	11 – 40%
4B	Mature	9+ inches	41 - 70%
4C	Mature	9+ inches	71 - 100%

1 Opening in forest cover type created by some type of disturbance
2 Natural meadow
3 Shrub cover type

Habitat structural stages can be used as indicators for:

- *Wildlife habitat* - Different wildlife species have different habitat requirements. For example, species that require large diameter trees in dense stand conditions find habitat in areas with 4B and 4C habitat structural stages.

- *Potential risk for future fire, insect and/or pathogen activity* - Trees growing in dense stands can be stressed by competition for water, nutrients and sunlight, making them more susceptible to insect attack.

- *Time since disturbances (fire, harvest, etc.) affected an area*. Age of vegetation is not directly linked to habitat structural stage; however, relative stages of succession can be implied. Early stages of succession are shown as habitat structural stages 1 or 2; later successional stages are usually in habitat structural stage 4A/ 4B/4C. Sometimes structural stages 3A/3B/3C may be the same age as 4A/4B/4C.

Current habitat structural stages for NFS lands within the North Fork Valley Geographic Areas are shown in Figure 3. Please note, on this map natural meadows in grass-forb cover types are shown as structural stage 1M, and shrub cover types are shown as structural stage 2S.

Figure 3. Current Habitat Structural Stages, North Fork Valley Geographic Area



Figure 4 displays the distribution of habitat structural stages for the forest and woodland cover types (spruce-fir, aspen, Douglas-fir, ponderosa pine, cottonwood, pinyon-juniper) on NFS portion of the North Fork Valley Geographic Area. Note: Natural meadows and shrublands are not included in the chart below.

Figure 4. Habitat Structural Stage Distribution of Forest and Woodland Cover Types, North Fork Valley Geographic Area



This information can also be summarized by looking at the distribution of size classes (Figure 5) and densities or canopy closure (Figure 6) of forest and woodland cover types on NFS lands within the Geographic Area. Again, natural meadows and shrublands are not included in the data shown in Figures 5 and 6.

Figure 5. Size Class Distributions of Forest and Woodland Cover Types



Figure 6. Canopy Closure Distributions in Forest and Woodland Cover Types



Characterization of Current Vegetation Conditions

Additional information on current conditions of different cover types is summarized in Table 2. (Also see timber, range and fire management sections.)

Table 2. Current Vegetation Characterization for NFS lands on North Fork Valley Geographic Area

Cover Type[1]	Composition of GA[1]	Age Distribution[2]	Habitat Structural Stages[1]	Canopy Conditions[1]	Past Activities (1955 – 2003)[3]	Effects of Roads/Trails[4]
Spruce-fir (Englemann spruce – subalpine fir)	121,800 acres 23%.	Age data not available	5%-3A 14%-3B 2%-3C 13%-4A **38%-4B** 27%-4C	28% - single-storied 72% - multi-storied 99% - continuous canopies	6% of type has been affected by timber harvest	74% of type is within ½ mile of a road or trail. Includes: 53 miles roads 132 miles trails
Aspen	212,600 acres 40%.	Age data not available	<1%-2T 7%-3A **41%-3B** 6%-3C 3%-4A 16%-4B 26%-4C	40% - single-storied 60% - multi-storied 99% - continuous canopies	2% of type has been affected by timber harvest	83% of type is within ½ mile of a road or trail. Includes: 138 mile roads 328 miles trails
Mixed Conifer (Douglas-fir, ponderosa pine)	1,500 acres <1%.	Age data not available	3%-3A 27%-3B 5%-3C **29%-4A** 26%-4B 10%-4C	28% - single-storied 72% - multi-storied 100% - continuous canopies	1% of these types have been affected by timber harvest.	80% of type is within ½ mile of an open road or trail. Includes: 1 mile road 3 miles trails
Pinyon-Juniper	3,200 acres 1%	Age data not available	24%-3A 2%-3B 29%-4A **40%-4B** 4%-4C	89% - single-storied 11% - multi-storied 100% - continuous canopies	3% of type has been mechanically treated to improve habitat for big game.	75% of type is within ½ mile of a road or trail. Includes: 2 miles roads 3 miles trails
Cottonwood	1,100 acres <1%	No age data available.	6%-3A 2%-3B **61%-4A** 32%-4B	62% - single-storied 38% - multi-storied 100% - continuous canopies	None recorded.	64% of these types are within ½ mile of a road or trail. Includes: 3 miles roads 2 miles trails

Cover Type[1]	Composition of GA[1]	Age Distribution[2]	Habitat Structural Stages[1]	Canopy Conditions[1]	Past Activities (1955 – 2003)[3]	Effects of Roads/Trails[4]
Gamble Oak – Mixed Mountain Shrub	97,500 acres 18%	No age data available.	100% 2S 44% large (>6.5') 48% med. (2.5-6.4') 8% small (<2.5')	31% - single-storied 69% - multi-storied 86% - >40% cover 14% - <= 40% cover	6% of these types have been treated with prescribed fire, <1% of these types have been mechanically treated to improve habitat for big game.	78% of these types are within ½ mile of a road or trail. Includes: 101 miles roads 153 miles trails
Sagebrush	8,600 acres 2%.	No age data available.	100% 2S 74% med. (2.5-6.4') 26% small (<2.5')	33% - single-storied 67% - multi-storied 91% - >40% cover 9% - <= 40% cover	None recorded in this period.	60% of type is within ½ mile of a road or trail. Includes: 21 miles roads 6 miles trails
Willow	10,300 acres 2%	No age data available	100% 2S 33% large (>6.5') 61% med. (2.5-6.4') 7% small (<2.5')	31% - single-storied 69% - multi-storied 61% - >40% cover 39% - <= 40% cover	None recorded.	73% of these types are within ½ mile of a road or trail. Includes: 11 miles roads 24 miles trails
Grass/Forb	40,200 acres 8%	No age data available.	100% 1M	Not applicable	None recorded for this period.	79% of these types are within ½ mile of a road or trail. Includes: 70 miles roads 79 miles trails.

1 Common Vegetation Unit/R2VEG database
2 Stand exam data for Grand Mesa Geographic Area.
3 RMACT database

INFRA transportation data related to R2VEG data. Includes all inventoried roads and trail, not just routes open to public use.

Comparison of Current Conditions to Historic Conditions

Environmental factors such as soils, slope, aspect, climate, and elevation determine the plant communities that potentially can grow on a given area. The stable plant community that establishes in the absence of any disturbances (i.e., fire, insect/pathogen mortality, windthrow, drought, harvest) is called the climax plant community. The area where a given climax plant community can grow is classified as a Potential Natural Vegetation (PNV) type, and is named for the climax plant community. For example, spruce-fir forests are the climax plant community at elevations from 10,000 to 12,000 feet, in the subalpine climatic zone (30-40 inches of precipitation annually, 50-70 frost free days, 30-40ºF mean annual air temperature) (Johnston et al. 2001). Areas where spruce-fir is the climax plant community are classified as the spruce-fir PNV type.

Draft PNV type classification has been completed for this Geographic Area. The PNV type composition on NFS lands within the North Fork Valley Geographic Area is shown in Figure 7. (PNV types with <0.1 percent composition are not included in Figure 7.) A map of PNV for the North Fork Valley GA is shown in Figure 8.

Figure 7. Composition of Potential Natural Vegetation Types on NFS lands within the North Fork Valley Geographic Area



* Unclassified, grass/forb or shrub types where the PNV type has not yet been assigned.

Figure 8. Potential Natural Vegetation on NFS portion of North Fork Valley Geographic Area



Historically, for any given PNV type, natural disturbances (i.e., fires, insect outbreaks) occurred at characteristic intervals and intensities, called regimes. When a disturbance was intense enough to change the existing plant community (e.g., some or all of the overstory was killed), the remaining vegetation followed a natural progression, or succession, of plant communities that changed over time. If no further disturbances

occurred, an area eventually returned to the climax plant community. However, if multiple disturbances occurred on the same area earlier seral plant communities would be perpetuated on a site for extended periods. The natural disturbance regimes and succession cycles resulted in shifting mixtures (or ranges) of different seral plant communities (or stages) within any given PNV type at any point in time.

The Vegetation Dynamics Development Tool (VDDT) (Beukema et al. 2003) was used to model the expected range in seral conditions that would have existed under historic disturbance regimes for forest, woodland and shrub PNV types. The results displayed in the following tables are conditions that would have been distributed across an entire PNV type on the Geographic Area. Results are scale dependant. For example, if a large fire burned an entire watershed, all of a given PNV type in that watershed could be set back to an early seral condition. However, when the watershed is considered as part of the Geographic Area, the mixture of seral conditions for that PNV type would be within the expected ranges. Both situations would have occurred naturally.

The next sections include comparisons of current conditions to historic conditions for the most common PNV types on the North Fork Valley Geographic Area, to see where differences occur. This information can be used to identify future management opportunities.

Spruce-Fir PNV Type

The spruce-fir PNV type is mapped on 12 percent of the North Fork Valley Geographic Area. The climax community in this PNV type includes Englemann spruce and supalpine fir as the overstory species. In some areas Englemann spruce may be the only overstory species (Romme et al. 1999). The spruce-fir PNV type includes 44 percent of what is currently classified as spruce-fir cover type.

Table 3 describes the seral stages and the timeline for succession in the spruce-fir PNV type. This table also displays a comparison of the modeled historic range of seral conditions to the current seral conditions (as percentages) within the spruce-fir PNV type.

Table 3. Succession (Seral Stages) in Spruce-Fir PNV Type

	Early	**Early-Mid**	**Late-Mid**	**Late**
Seral Stage Descriptions*	Initially grass/forb, low shrub with spruce seedlings eventually becoming established. May last 50 to-200 years, depending on the time it takes trees to become established.	Dense pole-sized spruce, grass and forb understory, lasting up to 150 years	Mature spruce overstory with fir and spruce trees growing in the understory. May persist 150 to 300 years.	Variable density of mature conifer, multiple age and size classes. New trees can become established in gaps in canopy. Lasts until next stand replacing disturbance
VDDT Modeled Range of Seral Conditions	27-32%	20-24%	12-13%	31-40%
Current Seral Conditions	14%	32%	53% (Limited age data makes it difficult to differentiate between late-mid and late seral conditions.)	

*Romme et al. 2003

Currently much of the spruce-fir PNV type is in mid- to late seral conditions (including both early-mid and late-mid), less in early seral conditions. This PNV type occurs at high elevations and over 70 percent of this type is on steeper slopes (> 40 percent slopes). Much of this PNV type is within or adjacent to areas affected by extensive fires on the Grand Mesa in the mid to late 1800s (Sudworth 1900). The current conditions expressed in the spruce-fir PNV type demonstrate the effect of large scale disturbances setting large areas to the same seral condition. Not enough time has passed since the last major disturbance (the late 1800s) to allow a full range of seral conditions to develop in this PNV type.

The historic fire regime for the spruce-fir PNV type was long return interval (> 200 years), stand replacing fires, which could cover areas from 1,000 to 10,000 acres (Peet 1981 as cited in Neely et al. 2001) mixed with infrequent low-intensity surface fires that affected much smaller areas. Fire return intervals tended to be longer at higher elevations and in moist depressions and valley bottoms, up to 500 years (Romme et al. 1999). Recent fires have been very infrequent and very small in this PNV type (see Fire Management Section).

A variety of insect and pathogen agents affect spruce-fir forests (see Forest Health section). Spruce beetle outbreaks have occurred in several areas of this PNV type in the late 1970s early 1980s.

Because much of the spruce-fir PNV type occurs in inaccessible areas, past harvest activities in this PNV type have been localized in certain areas (Black Mesa, Hubbard and Horse Ranch Park areas). The majority of harvest in this PNV type has occurred on Black Mesa in the southern portion of the GA. Clearcuts in the 1960s, shelterwood harvests in the 1970s, commercial thinning and selective sanitation/salvage harvests in the 1980s were the silvicultural methods used on these areas. The clearcuts are currently in the early seral stage. The sanitation/salvage areas are included in the early-mid

conditions. Shelterwood harvest practices reduced the density of stands but did not alter the seral condition.

Spruce-Fir-Aspen PNV Type

The spruce-fir-aspen PNV type is mapped on 32 percent of the North Fork Valley Geographic Area. The climax community in this PNV type includes Englemann spruce and subalpine fir as the dominant overstory species with aspen occurring in all seral stages. The spruce-fir-aspen PNV type is comprised of 50 percent of the current spruce-fir cover type and approximately 46 percent of the current aspen cover type.

Table 4 describes the seral stages and timeline for succession in the spruce-fir-aspen PNV type. This table also displays a comparison of the modeled historic range of seral conditions to the current seral conditions (as percentages) within the spruce-fir-aspen PNV type.

Table 4. Succession (Seral Stages) in Spruce-Fir-Aspen PNV Type

	Early	**Early-Mid**	**Late-Mid**	**Late**
Seral Stage Descriptions*	New stand of aspen seedlings/suckers with grass and forb understory lasting 30-50 years	Dense pole-sized aspen, grass and forb understory, lasting up to 100 years	Mature aspen overstory with conifer trees growing in the understory. May take 100 to 200 years for conifers to dominate stand.	Mature conifer, scattered mature aspen in overstory. New trees can become established in gaps in canopy. Lasts until next stand replacing disturbance
VDDT Modeled Range of Seral Conditions	13-19%	22-29%	13-16%	35-49%
Current Seral Conditions	6%	69%	25% (Limited age data makes it difficult to differentiate between late-mid and late seral conditions.)	

*Romme et al. 2003

This PNV type is generally below the spruce-fir PNV type in elevation. Approximately 60 percent of the spruce-fir-aspen PNV type currently is dominated by aspen in the overstory, and 36 percent currently has more conifer in the overstory. As evidenced by the percentages in the table above, basically all this PNV type is in mid seral conditions (includes both early-mid and late-mid). Current conditions are the result of large scale fires that burned through this area in 1878 to 1879, with less extensive burning occurring in 1883 to 1885 and again in 1890 to 1892 (Sudworth 1900). The large amount of area in the early-mid seral condition is due to a pulse of aspen regeneration that occurred following these fires. The majority of these stands are in the 80 to 100 year old range. The current conditions expressed in the spruce-fir-aspen PNV type demonstrate the effect of large scale disturbances setting large areas to the same seral condition. The time since the last major disturbance has not been sufficient to allow a full range of seral conditions to develop in this PNV type.

The fire regime in the spruce-fir-aspen PNV type is similar to that described above for the spruce-fir PNV type (long return interval stand replacing fires affecting large areas, mixed with infrequent low-intensity fires affecting small areas). Recent fires have been infrequent and very small in this PNV type. The Black Mesa portion of the Geographic Area was not affected by the fires mentioned above.

Little of this PNV type across the North Fork Valley Geographic Area has been affected by harvest activity. What has occurred is localized in the Black Mesa and Mendicant Ridge areas. Clearcuts to treat aspen and more recently commercial thinning have occurred in these areas.

Aspen PNV Type

In the aspen PNV type, pure aspen is the climax plant community, with no conifer species present. The aspen PNV type is mapped on 21 percent of the North Fork Valley Geographic Area. Approximately 49 percent of what is currently shown as aspen cover type is included in the aspen PNV type.

Table 5 depicts the seral stages and timeline for this PNV type as well as a comparison of the modeled historic range of seral conditions to the current seral conditions (as percentages).

Table 5. Succession (Seral Stages) in Aspen PNV Type

	Early	**Early-Mid**	**Late-Mid**	**Late**
Seral Stage Descriptions*	New stand of aspen seedlings/suckers with grass and forb understory lasting 10 to 20 years	Dense pole-sized aspen, grass and forb understory, lasting 50 to 80 years	Mature aspen overstory, aspen regeneration in the understory where overstory gaps results from individual tree mortality, lasting up to 80 years	Stable multi-storied, multi-aged aspen stand, predominantly forb understory.Lasts until next stand replacing disturbance
VDDT Modeled Range of Seral Conditions	8-14%	23-26%	17-24%	23-43%
Current Seral Conditions	5%	58%	36% (Limited age data makes it difficult to differentiate between late-mid and late seral conditions.)	

*Romme et al. 2003

The Aspen PNV type is also predominantly in mid to late seral conditions. Few areas are in early seral conditions. As with the previous PNV type, the distribution of seral conditions in the aspen PNV type is the result of past large scale fire activity in the late 1800s (Sudworth 1900).

The same fires that influenced the distribution of seral conditions in the spruce-fir-aspen PNV type also affected this aspen PNV type. Because aspen matures faster than spruce

or fir, this PNV type shifts into the later seral stages more rapidly, as refelected in the higher amount of late-mid seral condition as compared to the spruce-fir-aspen PNV type.

The historic fire regime for aspen is more frequent than for the two previous PNV types. Aspen forest appear to be dominated by a regime of relatively infrequent, large fires at higher elevations, and smaller more frequent fires at lower elevations (Kulakowski and Veblen, Draft 2004). Recent fires have been very infrequent and small in this type.

Two percent of this PNV type has been affected by timber harvest. Clearcuts were done mostly in the 1980s and 1990s in the Leroux Creek, Cunningham Creek, Terror Creek, Alder Creek and Black Mesa areas. These harvested areas are included in the early-mid seral conditions listed in the table above.

Oak-Serviceberry PNV Type

The Oak-Serviceberry PNV type has been mapped on 17 percent of NFS lands on the North Fork Valley Geographic Area. Historically, fires occurred in this PNV type at frequent intervals (every 15-30 years). The dominant shrub species Gambel oak and serviceberry resprout following fire so areas quickly returned to shrubs. This type transitioned from one seral stage to another at approximately 30 year timesteps. Table 6 compares current conditions in this PNV type to the expected historic mix of seral conditions.

Table 6. Succession (Seral Stages) in Oak-Serviceberry PNV Type

	Early	**Mid**	**Late**
Seral Stage Descriptions*	Stands are dominated by resprouting shrubs with grasses and forbs in the understory. Persists to 20 or 30 years	Dense clumps of shrubs, usually < 6 feet high. Grass and forbs between clumps. Persists from 30 to 40 years.	Shrubs > six feet high, more open canopy. Grass/forbs between and beneath shrub canopies. Persists until next stand replacing disturbance.
VDDT Modeled Range of Seral Conditions	30-70%	28-34%	0-36%
Current Seral Conditions	6%	50%	44%

*Romme et al. 2003

Most of the oak-serviceberry PNV type in the North Fork Valley Geographic Area is currently in mid and late seral conditions, and very little is in early seral conditions. The disparity between the historic conditions and current conditions is because there have been few recent disturbance in this PNV type, primarily due to fire suppression efforts over the past 100 years.

Frequent fires are the historical regime in this PNV type (0 - 35 years). Fires would have started below the forest on what is mostly private land now, and burned into the oak-serviceberry PNV type on the Geographic Area. Fire suppression efforts, cover type conversion and past livestock grazing have not allowed fires to burn in this PNV type as

frequently as in the past. As a result, the majority of this vegetation type has aged into mid and late seral conditions.

Approximately seven percent of this PNV type has been treated with prescribed fire (6,100 acres) and mechanical treatments (700 acres). These projects occurred in the mid 1980s, and mid 1990s, and account for the early seral conditions represented in the tables above. Some of the old treatments are approaching the mid seral condition. The purposes for these treatments was to improve habitat for big game species by restoring earlier seral conditions that provide more forage than later seral conditions.

Key Findings

- Aspen is currently the dominant tree species occurring on the North Fork Valley Geographic Area, with stands dominated by aspen occurring on 40 percent of the Geographic Area and stands of aspen mixed with spruce-fir cover types currently occupying 23 percent of the Geographic Area.
- The large extend of aspen is the result of large scale fires in 1878 to 1879, with less extensive burning occurring in 1883 to 1885 and again in 1890 to 1892 (Sudworth 1900), that affected the northern two-thirds of the North Fork Valley Geographic Areas. As a result, the majority of this geographic area is currently in mid seral conditions.
- Approximately 31 percent of forest and woodland cover types are in the sapling/pole size class (mostly in the aspen cover type), and 58 percent are in mature size class (mostly in the spruce-fir cover type).
- The majority of the current forest and woodland vegetation conditions – 87 percent - have dense canopy closures (> 40 percent canopy closure).
- There is very little early seral condition in any cover type on the North Fork Valley Geographic Area, as shown in Tables 3 through 6.
- When comparing the compositions of current vegetation cover types to PNV types (Figure 1 through Figure 7), the forest types and bare areas are approximately equivalent. The biggest differences occur in the grass/forb types; however, much of the current grass/forb cover type is also the early seral stage of forest PNV types. Grass/forb PNV types are classified on only a very small amount (< one percent) of the North Fork Valley Geographic Area.

Vegetation Trends

- The trend across all vegetation cover types on the North Fork Valley Geographic Area is to continue successional progress predominantly with the absence of either natural or human-caused disturbances. Structural and compositional conditions in each cover type will continue to progress along successional timelines. A shift from aspen dominated forests to conifer dominated forests is also occurring as a result of successional changes.

Vegetation Management Implications

Management implications of vegetation conditions relate to differences between current and historic vegetation conditions. Where landscape conditions are similar to historic conditions landscapes are more resilient following disturbances, suitable habitat is provided for all species that evolved there, and ecosystems are productive and sustainable over time. Where current conditions deviate from historic ranges, there are increased risks for natural disturbances to be of higher intensities and affect larger areas than in the past; some habitats may become very limited or lost, threatening species viability; and loss of productivity can result in both ecological and economic losses. Management activities that can include both human and natural disturbances can be used to alter vegetation conditions. Management implications for specific cover types are discussed below.

Spruce-Fir

- The spruce-fir cover type occurs in both the spruce-fir and the spruce-fir-aspen PNV types. Disturbances (human or natural) that occur in the spruce-fir-aspen PNV type will result in aspen regeneration within a very short period of time. Disturbances in the spruce-fir PNV type do not have aspen as an early seral species, and may take 50 to 200 years to regenerate to spruce.

- Both the spruce-fir and spruce-fir-aspen PNV types have large percentages in mature, dense structural conditions. These conditions are ripe for fire and epidemic insect and pathogen outbreaks (see Forest Health and Fire Management sections). Because so much of the landscape is in the same condition, there is a high probability that large areas can be affected at the same time. Large scale disturbances (both fire and insect outbreaks) are the natural regimes for these PNV types. Management activities that increase diversity in age, size and seral condition, could be used to reduce the potential effects of natural disturbances.

- The majority of this cover type in the North Fork Valley Geographic Area (90,100 acres, or 74 percent) is in Wilderness and currently unroaded areas. Only natural processes will alter vegetation conditions in Wilderness areas, and potentially in unroaded areas where future management decisions (i.e., theme designations) limit or restrict management activities.

Aspen

- The aspen cover type occurs in both the spruce-fir-aspen PNV type (as the early through late-mid seral conditions) and the aspen PNV type (all seral stages). Based on the percentages in each seral stage displayed in the above tables for these PNV types, aspen currently dominates the North Fork Valley Geographic Area landscape, as a result of fires in the late 1800s.

- Aspen stands become very susceptible to cankers and root rots as they mature. A large percentage of the aspen cover type on the North Fork Valley Geographic Area is estimated to be between the age of 80 and 120 years, based on time since

last major fire disturbance. An increasing amount of mortality caused by these fungal agents is expected in the future.

- Approximately half of the aspen stands on the North Fork Valley Geographic Area include conifer species in the understory or as codominant species. In the absence of some disturbance, these stands will eventually succeed to conifer dominated cover types. A shift from an aspen cover type to various conifer cover types will change the types of habitat available on the North Fork Valley Geographic Area to favor those species that require mature conifer forests.

Gambel Oak and Mountain Shrubs

- Current conditions in the Gambel oak and mixed mountain shrub cover types have less patchiness and structural stage diversity than would have occurred historically. The result is current conditions are more susceptible to higher intensity fires that may affect larger areas of land than would have occurred in the past. There is also less diversity in the types of habitat provided by these cover types than would have been present.

References

Beukema, S.J., W.A. Kurz, C.B. Pinkham, K. Milosheva, L. Frid. 2003. Vegetation Dynamics Development Tool users' guide, Version 4.4., ESSA Technologies Ltd. Vancouver, BC. 234 pp.

Hoover, R.L., and D.L. Wills, ed. 1984. Managing forested lands for wildlife. Colorado Division of Wildlife in cooperation with USDA Forest Service, Rocky Mountain Region. Denver, CO, 459 pp.

Johnston, B.C., L. Huckaby, T.J. Hughes and J. Pecor. 2001. Ecological types of the Upper Gunnison Basin: Vegetation-soil-landform-geology-climate-water land classes for natural resource management. Technical Report R2-RR-2001-01. USDA Forest Service, Rocky Mountain Region. Lackwood, CO. 858 pp.

Kulakowski D. and T.T. Veblen. 2004. Draft Historic Range of Variability Assessment for Forest Vegetation of the Grand Mesa National Forest, Colorado. Department of Geography, University fo Colorado, Boulder, CO. 131 pp.

Neely, B., P.Commer, C.Mortiz, M. Lammert, R. Rondeau, C. Pague, G. Bell, H. Copeland, J. Humke, S. Spackman, T. Schulz, D. Theobald, and L. Valutis. 2001. Southern Rocky Mountains: An Ecoregional Assessment and Conservation Blueprint. Prepared by the Nature Conservancey with the support from the USDA Forest Service, Rocky Mountain Region, Colorado Division of wildlife, and Bureau of Land Management.

Romme, W.H., M.L. Floyd, D. Hanna, J.S. Redders, K. McGarigal, J.P. Lindsey, M. Crist. 1999. Landscape Condition Analysis for the South Central Highlands Section, Southwestern Colorado and Northwestern New Mexico, Chapter 5: Spruce-Fir Forests. Forest Sciences Department, Colorado State University, Fort Collins, CO. 62 pp.

Romme, W.H., K. McGarigal and D. Goodwin. 2003. Vegetation Dynamics of the North Fork Valley Landscape, Southwestern Colorado (Draft). Department of Forest, Rangeland and Watershed Stewardship. Colorado State University. Fort Collins, CO. 62 pp.

Sudworth, G.B. 1900. Battlement Mesa Forest Reserve, in The White River Plateau and Battlement Mesa Forest Reserves. USDI U.S. Geological Survey. Washington Government Printing Office. Pages 181-243.