

Field Guide for Managing African Rue in the Southwest



Cover Photos

Top left: G.A. Cooper, USDA-PLANTS Database, Bugwood.org

Bottom left: G.A. Cooper, USDA-PLANTS Database, Bugwood.org

Right: Joseph M. DiTomaso, University of California-Davis, Bugwood.org

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotape, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at http://www.ascr.usda.gov/complaint_filing_cust.html and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: program.intake@usda.gov.

USDA is an equal opportunity provider, employer and lender.

African rue (*Peganum harmala*)

Caltrop family (Zygophyllaceae)

African rue is listed as a noxious weed in both Arizona and New Mexico. This field guide serves as the U.S. Forest Service's recommendations for management of African rue in forests, woodlands, and rangelands associated with the Service's Southwestern Region. The Southwestern Region covers Arizona and New Mexico, which together have 11 national forests. The Region also administers 4 national grasslands that are located in northeastern New Mexico, western Oklahoma and the Texas panhandle.

Description

African rue (synonyms: wild rue, rue weed, Syrian rue, harmal, esfand, ozallaik, ruin weed) is a succulent perennial herb with a robust, deep-growing root system that is a major obstacle to plant control.

Growth Characteristics

- Perennial bushy herb that grows 1 to 1.5 feet tall and 3-4 feet in diameter.
- Has a hardy woody taproot that reaches 25 feet or more downwards into the soil profile. Creeping, lateral roots are produced about 12-15 inches below the surface that can extend up to 20 feet away from the parent plant.
- Bright green leaves are alternate, smooth and divided deeply into narrow lobes. Leaves have a very disagreeable odor when crushed.
- Single flowers are borne along the stem and in the forks of leaves. Flowers have five white petals and produce a green, orange, or brown-colored cylindrical seed pod (2-4 celled) with 45-60 seeds.
- Reproduces by seed, but new shoots and plants are also produced from adventitious buds along lateral roots. Seed is secured in a leathery fruit capsule; each plant may produce as many as 1,000 capsules.

Ecology

Impacts/Threats

This weed is extremely toxic to cattle, sheep, horses, and humans as it contains at least four poisonous alkaloids. The seeds and fruit are the most toxic, followed by young leaves and mature leaves. However, animals typically avoid eating African rue because of its bad taste and smell. In addition, most parts of the plant have allelopathic chemicals that reduce the growth of surrounding native plants.

Site/Distribution

Common in disturbed and barren areas such as roadsides, parking lots, oil pads, corrals, stockyards, open fields, abandoned crop fields, river banks, ditches and arid/semi-arid desert areas, especially in moist soils. It can grow in many soil types, including alkaline or highly saline soils.

African rue occurs throughout most western States, including New Mexico, Arizona, Texas, California, Montana, Nevada, Oregon, and Washington. In New Mexico, it is primarily found in southern counties with isolated populations in central and northwestern counties. It is common in eastern Arizona and western Texas.

Spread

Seeds and root fragments are readily transported by running water. Seeds can be carried in mud that sticks to vehicles, railcars, and earth-moving equipment; thus, new plants easily spread along transportation corridors. Seed may also adhere to the feet, fur, or feathers of animals. Seed longevity is unknown but is probably persistent.

Invasive Features

African rue undergoes robust and rapid vegetative growth when soil moisture is available. It favors disturbed areas such as those impacted by earth-moving equipment, excessive grazing, or vehicle use. Disturbed areas with moist soil such as roadsides, river or stream banks, riparian corridors, waterways, and drainage or irrigation ditches are especially susceptible.

This invasive weed is extremely drought tolerant and has “drought evader” growth characteristics. In southern New Mexico, plants die back to their roots during winter and initiate new annual growth in mid-to-late March. After spring growth and flowering, plants may become senescent (decrepit-looking) and die back to the woody base as soils dry during summer. With cooler temperatures and additional moisture coming later in the season, African rue usually undergoes a second vegetative growth phase that lasts until freeze occurs in early November.

Management

Once established, African rue is difficult to eradicate without impacting associated desirable plants. However, methods to control the plant other than by herbicidal application are quite limited. Grazing is not an option due to toxicity, and there are no known classic biocontrol agents at this time. The plant grows back quickly after mowing or burning, and deep cultivation only fragments and spreads the roots. Since African rue is able to regenerate from root fragments, grubbing or cultivation may actually increase the population size.

The following actions should be considered when planning a management approach for African rue:

- Maintain healthy plant communities to prevent or limit infestations of African rue.
- Detect and eradicate new populations of African rue as early as possible, especially along roadways, waterways, and ditches.
- Use foliar or soil-active herbicides at recommended rates for effective control of African rue populations.

Table 1 summarizes some management options for common situations involving African rue. Further details on these management options are explained below. Choice of control method(s) for African rue depends on the land use and current site conditions (accessibility, terrain, soil and air temperature, weather, extent and density of infestations, other flora and fauna present, etc.). Other considerations include treatment effectiveness, cost, and the time needed to achieve control.

Physical Control

Because African rue has a complex root system consisting of lateral roots and a deep tap root, it is extremely difficult to control the plant solely through the use of mechanical methods. The more practical and economical way to obtain effective long-term control is to use herbicide treatments alone or in combination with a physical method.

Manual Methods

Hand-pulling, digging, or hoeing for control of mature plants is virtually impossible due to the depth of the root system. Hand removal of seedlings and young plants that is continued year after year may eventually control individual plants, but this approach is tedious and usually impractical.

Mechanical Methods

Controlling African rue by plowing or grubbing individual plants is ineffective since the roots are too deep to dig out and any root parts left in the ground will likely re-sprout. In many instances, cultivation may actually lead to an increase in population size. In certain agronomic situations, repeated deep cultivation combined with reseeding perennial plants has been found to be somewhat successful; but this is an expensive option. Mowing or blading also causes the plant to re-sprout and spread.

Prescribed Fire

Fire is not an effective control method as plants grow back rapidly after burning.

Cultural Control

Early detection and plant removal, especially for seedlings and young plants, are critical for preventing African rue establishment and spread. Land managers, road crews, and the local public should be educated on identification so they can report suspected populations. Vehicles, humans, and livestock should be discouraged from traveling through infested areas; a program to check and remove seeds from vehicles and livestock after going through infested areas should be implemented to help stop the spread. Hay, straw mulch, planting seeds, fill, and other related materials should be certified to be weed free before use in areas undergoing treatment.

Table 1. Management options*

Site	Physical Control	Cultural Control	Biological Control	Chemical Control
Roadsides	Few options available. Combine suppression methods (e.g., repeat deep cultivation or mowing) with reseeding desired native perennial plants.	Use seed, mulch, and fill materials certified to be weed-free. Limit disturbances along roadsides. Implement requirements for vehicle operations and for reporting infestations along roads.	None known at this time.	Use truck or tractor spraying equipment. Wash underneath to prevent spread.
Ditches and waterways	Few options available. Hand extraction or hoeing can be used to target seedlings.	Limit disturbances along waterways and ditches. Increase public awareness and reporting of plant presence along ditches and roadways.	Same as above.	Use herbicide with approved aquatic labeling when near or in water.
Rangeland	Few options available.	Use seed and forage hay certified to be weed-free; use pellets for horses in backcountry areas. When moving livestock or vehicles through infested areas, inspect mud on animals, clothing, and vehicles and remove any seeds before entering uninfested areas.	Same as above.	Use ground broadcast spraying with ATV or tractor; however, backpack spraying may be more practical in areas with difficult access. Consider individual plant treatment in areas with less dense infestation or near desirable vegetation.
Wilderness and other natural areas	Few options available. Hand removal repeated year after year may be required for effective control.	Use seed and forage hay certified to be weed-free; use pellets for horses in backcountry areas. Post signs warning visitors to remove seeds.	Same as above.	Use backpack sprayers for individual plant treatment or ground broadcast spraying with an ATV or tractor, if allowed.

* Choice of a particular management option must be in compliance with existing regulations for the land resource.

Biological Control

Grazing

Grazing is not a viable option for removing African rue. Due to its bad taste and smell, livestock usually avoid the plant unless other forages are unavailable. Occasionally, African rue is found as a contaminant in hay bales. Effects of poisoning on livestock include loss of appetite, trembling, and loss of coordination. Severe poisoning can result in hemorrhaging in the heart and liver.

Classical Biological Control

There are no known classical biocontrol agents (insects, pathogens, etc.) available for African rue at this time.

Chemical Control

Numerous herbicides have been investigated in field trials for African rue control. These tests show that only chemicals translocated deep into the plant's root system can effectively kill the weed (see table 2). Certain selective, systemic herbicides (e.g., metsulfuron) will suppress seasonal top

growth; however, most plants recover and regrow within a year of treatment. Unfortunately, the few herbicides that provide effective long-term control (e.g., imazapyr, tebuthiuron, and hexazinone) are nonselective and may cause unacceptable injury to desirable species, especially grasses.

It is very important to read the herbicide label carefully and follow all instructions when mixing and applying any herbicide. Equipment used to spray herbicide should be calibrated. Aquatically approved formulations should be used in or near water. When using aquatic herbicide formulations, an aquatically approved surfactant should also be added to the spray mix.

Considerations for Herbicide Use

Based on research and practical experience, the following guidelines should be used for herbicidal control of African rue:

- **Plant condition and growth stage** at the time of application are critical for control success. When using a foliar spray, treating African rue that is healthy and robust in late summer (September-October) is optimal. Plants in the early vegetative and bloom stage in spring (April) can also be successfully controlled provided soil moisture is adequate. Do not spray African rue that has been stressed from drought, disease, insects, or other causes that might have resulted in dieback, yellowing, or other plant damage.
- **Weather conditions** at the time of spraying are important for successful herbicide activity. Coverage is best with low wind speeds (3 to 8 mph) and cool temperatures (between 50 °F and 85 °F). Air temperatures above 85 °F increase spray solution volatility and thus reduce the amount of herbicide available to be absorbed by the plant. Do not spray a foliar-active herbicide if a rainstorm is expected within 6 hours of application.
- **The plant community** associated with the African rue population should be closely evaluated before treating an area. Use of a nonselective herbicide should not be made in areas where grass

preservation is desired. Individual plant spraying will minimize grass damage relative to broadcast spraying, but some damage should still be expected.

- **Time needed to gain control** is slow, requiring a year or more to kill a plant when either sprayed foliarly or treated with a soil-active herbicide.

Herbicide Application Options

Broadcast foliar spray treatment – The ideal time to spray African rue foliarly varies from year to year because of specific weather conditions. Expect average to above-average winter moisture to produce a prolific burst of new vegetation growth in spring, which is needed before spraying to maximize chemical uptake and movement (translocation) through all portions of the plant. In droughty years, African rue will have poor foliage and flower growth; therefore, spraying is not recommended. Similarly, healthy foliage that results from above-average summer rainfall can be sprayed late in the season. At the time of spraying, soil temperature at a 6-inch depth should be between 55 °F and 70 °F. In years with average to above-average winter precipitation, the desired growth stage and soil temperature range occurs roughly from April 1 to May 15, plus or minus 15 days. In late season, the desired growth stage is usually from September 1 to October 1.

Using imazapyr alone or in combination with other herbicides has consistently provided the best African rue control. Imazapyr (e.g., Arsenal®) is the most common commercial product used for African rue control. Imazapyr is labeled for pasture/rangeland, noncropland sites, rights-of-way, industrial areas, fence rows, non-irrigation ditch banks, establishment and maintenance of wildlife openings, and for bare ground weed control. When broadcast spraying larger infestations, use a 1.75 lbs. a.i. per acre rate, which is equivalent to 3 pints of product per acre. When spot treating smaller populations or isolated plants, use an IPT approach as described below. A surfactant should be added to the herbicide mixture at rates specified on the herbicide label. Imazapyr is a nonselective herbicide so damage to grasses and other associated plants should be anticipated.

A new herbicide product called Viewpoint® (active ingredients: aminocyclopyrachlor + imazapyr +

Table 2. Herbicide recommendations

Common Chemical Name (active ingredient)	Product Example(s) ¹	Broadcast Treatment (rate per acre)	Spot Treatment (spray solution and pellets) ²	Time of Application	Remarks
Imazapyr	Arsenal Habitat [others available]	3 pints per acre	3%	Spray healthy regrowth foliage in good condition in late summer or early fall.	Imazapyr is a nonselective herbicide; anticipate damage to non-target plants from overspray, runoff, movement in soil, or root exudates. When performing IPT with a backpack sprayer, adjust nozzle to wet foliage thoroughly.
Aminocyclopyrachlor + metsulfuron + imazapyr	Viewpoint	3–18 ounces per acre	NA	Same as above.	This herbicide combination provides nonselective control and should only be used in non-crop areas.
Metsulfuron	Escort Ally	3.2–6.4 ounces per acre	NA	Same as above.	Metsulfuron is selective and provides seasonal foliage suppression but low mortality in African rue.
Tebuthiuron	Spike 20P	10–15 pounds of pellets	NA	Anytime but optimal time is just before a rain event.	Apply only on sandy or coarse soils; will likely damage associated vegetation.
Hexazinone	Velpar	NA	Mix 50:50 with water in a backpack sprayer. Apply 4 ml per plant (equivalent to 2 ml undiluted Velpar).	Anytime.	To apply as an IPT, squirt 4 ml of mixture on the soil surface next to each plant. Not recommended as a broadcast because it is nonselective and will damage or kill associated vegetation.

¹ Trade names for products are provided for example purposes only, and other products with the same active ingredient(s) may be available. Individual product labels should be examined for specific information and appropriate use with African rue.

² Spray solution is the herbicide/water ratio in a spray mix that may be used for spot treatment with backpack or hand-held sprayers. The amount of product applied during an annual growing season must not exceed the maximum application rate per acre as specified by the product label – refer to the product label for the site type and application.

metsulfuron) provides good to excellent control of African rue when applied at a rate of 13 to 18 ounces of product per acre. Viewpoint is formulated as a dispersible granule that is mixed in water and applied as a liquid spray. This herbicide is labeled specifically for non-crop situations only, such as spraying rights-of-way, oil pads, parking lots, etc. It provides broad-spectrum control of brush and weeds species. Care should be exercised when using this product in

the vicinity of crops or other desirable plants because this herbicide combination may injure or kill them. Read the herbicide label carefully for use directions.

Broadcast soil-active treatment – African rue growing on rangeland with sandy to loamy soils may be controlled with a broadcast rate of tebuthiuron (e.g., Spike 20P®) applied at 2 to 3 lbs. a.i. per acre. Unlike the liquid foliar-active herbicides, tebuthiuron is a dry pellet formulation that is

applied to the soil surface. These pellets remain intact until they are dissolved by rainfall and the chemical is moved into the soil where it is taken up by roots. Tebuthiuron is applied by aerial or ground methods and provides fair to good African rue control. However, damage to associated vegetation should always be anticipated at the relatively high rate of application required.

Individual plant treatment (IPT) – IPT is an effective way to control the plant; however, the method is often time-consuming and costly, especially if the African rue is dense. For practical reasons, IPT is ordinarily used only for small areas where African rue is growing sparsely and the infestation is easy to treat. Two active ingredients, hexazinone and imazapyr, are preferred IPT choices, but they are applied in different ways as described below:

- Hexazinone (e.g., Velpar®) can be applied as a liquid directly to the soil surface next to the canopy of individually treated plants. For IPT, hexazinone should be diluted as a 50:50 mix with water in a backpack sprayer. A quick squirt of the handle with the nozzle turned to a straight stream generally administers about 4 ml of solution (or 2 ml of active product), which is the desired rate of application. Before spraying, check the handle pressure and practice with water in the sprayer to calibrate this delivery rate. Adding a blue dye to the spray mixture helps to identify treated plants. Hexazinone moves into the soil profile with rain and is taken up by roots. Thus, application prior to anticipated rainfall is preferred, but the product usually works at any time of year.
- Imazapyr diluted as a 3 percent mixture with water is effective for IPT. Most herbicide activity is through the foliage so use a backpack sprayer with its nozzle adjusted to wet the top growth of individual plants completely. Add blue dye to the spray mixture to help view spray coverage and to identify treated plants.

Management Strategies

Early detection and control of new African rue populations – especially near waterways and ditches – are important activities for slowing the spread of this noxious weed. In

most cases, effective control of African rue will require herbicide spraying. Since each treatment situation is unique, site-specific management should be used. Depending on the intended land use and available budget, consider reseeding desired plant communities, as healthy plant populations can help deter spread of African rue.

Even with herbicide use, it may take years to reduce the presence of this plant, which can regenerate from deep-growing root parts as well as seed. Since it is ordinarily useless to treat an area only one time without retreatment, sufficient resources must be allocated for the area where control is attempted. After initial treatment, it is especially important that resources are also available to respray or retreat the treated area as necessary. Always anticipate the need for monitoring and additional control measures such as spot spraying with a backpack or hand-held sprayer.

References and Further Information

- Abbot, L.B., D. Lepak, and D.L. Daniel. 2007. Vegetative and Reproductive Phenology of African rue in the Northern Chihuahuan Desert. *The Southwestern Naturalist* 52(2):209–218.
- Duncan, K., K. McDaniel, and M. Renz. 2005. Chemical Weed and Brush Control for New Mexico Rangelands. Las Cruces: New Mexico State Univ. College of Agriculture and Home Economics and Cooperative Extension Service. Pp. 6. Available at http://aces.nmsu.edu/pubs/_circulars/CR_597.pdf.
- Fact Sheet: Stop the Spread of African Rue. November 2009. Published by the Government of South Australia/South Australian Arid Lands Natural Resources. Available at <http://www.saalnrn.sa.gov.au/> (accessed Sept. 2014)
- Hart, C.R. 2002. African rue - Biology and Management. University of Texas A&M Publication L-5429 10-02. Available at <http://pecosbasin.tamu.edu/media/2652/L-5429.pdf> (accessed Sept, 2014)
- La Plata County Weed Office, 2500 Main Avenue, Durango, CO. Available at www.lpcweeds.org/Africanrue.htm (accessed Sept. 2014)

McDaniel, K.C. and K. Duncan. 2006. African Rue Control: Ground Applications. New Mexico Cooperative Extension Service Publication BC-6. New Mexico State Univ., Las Cruces, NM.

Michelmore, M. 1997. African rue Management—Distribution, Biology, Impact and Control Strategies for *Peganum harmala* L. (Zygophyllaceae) in South Australia. Primary Industries, SA, Port Augusta, South Australia.

Northam, E. and W. Meyer et al. 2009. Non-Native Invasive Plants of Arizona. Produced by Conservation Districts and RC&D Areas of Arizona and the University of Arizona Cooperative Extension. Peer reviewed publication AZ 1482. Available at <http://cals.arizona.edu/pubs/natresources/az1482.pdf> (accessed Sept. 2014)

Renz, M. and F. Sholedice. 2006. Troublesome Weeds of New Mexico. Las Cruces: New Mexico State University College of Agriculture and Home Economics and Cooperative Extension Service. Pp. 54-55. Available at <http://www.emnrd.state.nm.us/fd/Publications/documents/NMTroublesomeWeeds.pdf> (accessed Sept. 2014)

The Plant Database. 2010. U.S. Department of Agriculture NRCS National Plant Database. Baton Rouge, LA. Available at <http://plants.usda.gov/java/profile?symbol=PEHA> (accessed Sept. 2014))

Suggested Web Sites

Dirty Dozen Invaders of the Southwest. 2002. NMSU, UA, USU, and U.S. Dept. of Agriculture cooperating. Available at http://www.invasiveweeds.com/dirty/african_rue/

For information on invasive species:
Plant Health and Pest Prevention Services:
<http://www.cdffa.ca.gov/phpps/ipc/weedinform/peganum.htm>

National Invasive Species Information Center:
<http://www.invasivespeciesinfo.gov/>

Invasive and Exotic Species of North America
<https://www.invasive.org/>

For information about calibrating spray equipment:
NMSU Cooperative Extension Service Guide A-613, *Sprayer Calibration*. Available at http://aces.nmsu.edu/pubs/_a/A-613.pdf

Oregon Department of Agriculture, Plant Division Noxious Weed Control. Available at http://www.oregon.gov/ODA/PLANT/WEEDS/profile_africanrue.shtml

**For more information or
other field guides, contact:**

USDA Forest Service
Southwestern Region
Forest Health
333 Broadway Blvd., SE
Albuquerque, NM 87102

**Or visit the Southwestern Region's
website for invasive species:**

<http://www.fs.usda.gov/goto/r3/invasivespecies>



The use of trade or firm names in this publication is for reader information only and does not imply endorsement of any product or service by the U.S. Department of Agriculture. Recommendations made here for pesticide use are not obligatory, nor do they imply that discussed uses have been registered. All uses of pesticides must be registered by appropriate State and/or Federal agencies before they can be applied.



CAUTION: Pesticides can be injurious to humans, domestic animals, desirable plants, and fish or other wildlife—if they are not handled or applied properly. Use all pesticides carefully and lawfully. Follow recommended practices for the disposal of surplus pesticides and pesticide containers.