


***Conservation Assessment  
for  
Grand Caverns Blind Cave Milliped (Trichopetalum weyeriense)***



*(From Barr, 1973)*

***USDA Forest Service, Eastern Region***  
December 2001

Julian J. Lewis, Ph.D.  
J. Lewis & Associates, Biological Consulting  
217 W. Carter Avenue  
Clarksville, IN 47129  
[Lewisbioconsult@aol.com](mailto:Lewisbioconsult@aol.com)



*This Conservation Assessment was prepared to compile the published and unpublished information on the Trichopetalum weyeriensis (Grand Caverns Blind Cave Milliped). It does not represent a management decision by the U.S. Forest Service. Though the best scientific information available was used and subject experts were consulted in preparation of this document, it is expected that new information will arise. In the spirit of continuous learning and adaptive management, if you have information that will assist in conserving the subject community and associated taxa, please contact the Eastern Region of the Forest Service Threatened and Endangered Species Program at 310 Wisconsin Avenue, Milwaukee, Wisconsin 53203.*

**Table of Contents**

**EXECUTIVE SUMMARY.....4**  
**NOMENCLATURE AND TAXONOMY .....4**  
**DESCRIPTION OF SPECIES.....4**  
**LIFE HISTORY .....4**  
**HABITAT.....5**  
**DISTRIBUTION AND ABUNDANCE .....5**  
**RANGEWIDE STATUS.....5**  
**POPULATION BIOLOGY AND VIABILITY .....5**  
**POTENTIAL THREATS .....5**  
**SUMMARY OF LAND OWNERSHIP AND EXISTING HABITAT**  
**PROTECTION ..... 7**  
**SUMMARY OF MANAGEMENT AND CONSERVATION ACTIVITIES 7**  
**RESEARCH AND MONITORING .....7**  
**RECOMMENDATIONS.....7**  
**REFERENCES ..... 7**

## EXECUTIVE SUMMARY

The Grand Caverns Blind Cave Milliped (*Trichopetalum weyerienseis*) is designated as a Regional Forester Sensitive Species on the Monongahela National Forest in the Eastern Region of the Forest Service. The purpose of this document is to provide the background information necessary to prepare a Conservation Strategy, which will include management actions to conserve the species.

*Trichopetalum weyerienseis* is a rare troglobitic (obligate cavernicolous) milliped that occurs in West Virginia from 15 caves distributed through Greenbrier, Monroe, Pendleton, and Pocahontas counties as well as eight caves in the adjacent part of Virginia.

## NOMENCLATURE AND TAXONOMY

**Classification:** Class Diplopoda  
Order Chordeumatida  
Family Trichopetalidae

**Scientific name:** *Trichopetalum weyerienseis* (Causey)

**Common name:** Grand Caverns Blind Cave Milliped

**Synonyms:** none

Causey (1963, in litt., reported in Holsinger, et. al. 1976) has raised the possibility that *Trichopetalum weyerienseis* may be a subspecies of *Trichopetalum whitei* rather than a distinct species. The genus *Trichopetalum* is being revised by Shear (in progress) and at present *T. weyerienseis* is being recognized as a full species.

## DESCRIPTION OF SPECIES

*Trichopetalum weyerienseis* is an eyeless, unpigmented (white) milliped. Typical of all trichopetalids this species has rows of elongate segmental setae extending in rows along the dorsal side. Identification of this species requires microscopic examination and dissection of the gonopods (copulatory apparatus) by a specialist familiar with the identification of millipeds. This species can be recognized in the field only at sites from which it has been previously identified by a taxonomist.

## LIFE HISTORY

Little is known of the life history of this species. Shear (1971) summarized the findings of Schubart (1934), who reported some observations on the mating of other millipeds of the Order Chordeumatida. In those animals the male secreted sperm from the seminal pores on the coxae of the second legs into coxal sacs on the postgonopodal legs. The secretions from the coxal sacs then form the seminal fluid into a spermatophore which is then transferred to

the cyphopods of the female during mating. Oviposition has not been observed, although some North American members of the order produce silk chambers for the egg laying.

Feeding is presumed to consist of picking up or scraping material from the substrate with the mouthparts then grinding it with the mandibles. It is presumably omnivorous although, again, nothing is known of the specifics about its feeding preferences.

## **HABITAT**

Trichopetalum weyerienses is a troglobite and occurs only in caves. Although nothing was stated about the specific habitat of this milliped, it presumably occurs under rocks, around organic debris and on damp riparian mudbanks as per related Trichopetalum species reported in eastern West Virginia by Holsinger, et. al. (1976).

## **DISTRIBUTION AND ABUNDANCE**

As presently understood the range of Trichopetalum weyerienses straddles the West Virginia/Virginia state line. The species is reported in West Virginia from 15 caves distributed through Greenbrier, Monroe, Pendleton, and Pocahontas counties (Holsinger, et. al. 1976). In Virginia the milliped has been reported from 8 caves distributed through Augusta, Page, Rockingham and Shenandoah counties (Holsinger and Culver, 1988).

## **RANGEWIDE STATUS**

**Global Rank:** G2/3 = imperiled/vulnerable; G2 typically includes species known from between 6-20 sites, G3 includes those known from 21-99 sites. The G2/G3 ranking signifies uncertainty about the rank. Although reported from 23 caves, at least two of the known sites are probably parts of the same cave system (Grand Caverns and Madison Saltpetre Cave) and would count as one occurrence of the species rather than two. Thus, Trichopetalum weyerienses straddles high G2/low G3 in rarity.

**West Virginia State Rank:** S2 = imperiled; S2 typically encompasses a species known from between 6-20 sites within the state. Trichopetalum weyerienses is known from 15 sites in West Virginia.

## **POPULATION BIOLOGY AND VIABILITY**

Nothing is known of the population biology of Trichopetalum weyerienses.

## **POTENTIAL THREATS**

Due to the presence of Trichopetalum weyerienses solely in the restricted cave environment, it is susceptible to a wide variety of disturbances (Elliott, 1998). Caves are underground drainage conduits for surface runoff, bringing in significant quantities of nutrients for cave communities. Unfortunately, contaminants may be introduced with equal ease, with devastating effects on cave animals. Potential contaminants include (1) sewage or fecal

contamination, including sewage plant effluent, septic field waste, campground outhouses, feedlots, grazing pastures or any other source of human or animal waste (Harvey and Skeleton, 1968; Quinlan and Rowe, 1977, 1978; Lewis, 1993; Panno, et al 1996, 1997, 1998); (2) pesticides or herbicides used for crops, livestock, trails, roads or other applications; fertilizers used for crops or lawns (Keith and Poulson, 1981; Panno, et al. 1998); (3) hazardous material introductions via accidental spills or deliberate dumping, including road salting (Quinlan and Rowe, 1977, 1978; Lewis, 1993, 1996).

Habitat alteration due to sedimentation is a pervasive threat potentially caused by logging, road or other construction, trail building, farming, or any other kind of development that disturbs groundcover. Sedimentation potentially changes cave habitat, blocks recharge sites, or alters flow volume and velocity. Keith (1988) reported that pesticides and other harmful compounds like PCB's can adhere to clay and silt particles and be transported via sedimentation.

Impoundments may detrimentally affect cave species. Flooding makes terrestrial habitats unusable and creates changes in stream flow that in turn causes siltation and drastic modification of gravel riffle and pool habitats. Stream back-flooding is also another potential source of introduction of contaminants to cave ecosystems (Duchon and Lisowski, 1980; Keith, 1988).

Smoke is another potential source of airborne particulate contamination and hazardous material introduction to the cave environment. Many caves have active air currents that serve to inhale surface air from one entrance and exhale it from another. Potential smoke sources include campfires built in cave entrances, prescribed burns or trash disposal. Concerning the latter, not only may hazardous chemicals be carried into the cave environment, but the residue serves as another source of groundwater contamination.

Numerous caves have been affected by quarry activities prior to acquisition. Roadcut construction for highways passing through national forest land is a similar blasting activity and has the potential to destroy or seriously modify cave ecosystems. Indirect effects of blasting include potential destabilization of passages, collapse and destruction of stream passages, changes in water table levels and sediment transport (Keith, 1988).

Oil, gas or water exploration and development might encounter cave passages and introduce drilling mud and fluids into cave passages and streams. Brine produced by wells is extremely toxic, containing high concentrations of dissolved heavy metals, halides or hydrogen sulfide. These substances can enter cave ecosystems through breach of drilling pits, corrosion of inactive well casings, or during injection to increase production of adjacent wells (Quinlan and Rowe, 1978).

Cave ecosystems are unfortunately not immune to the introduction of exotic species. Out-competition of native cavernicoles by exotic facultative cavernicoles is becoming more common, with species such as the exotic milliped Oxidus gracilis affecting both terrestrial and aquatic habitats.

With the presence of humans in caves comes an increased risk of vandalism or littering of the habitat, disruption of habitat and trampling of fauna, introduction of microbial flora non-native to the cave or introduction of hazardous materials (e.g., spent carbide, batteries). The construction of roads or trails near cave entrances encourages entry.

## **SUMMARY OF LAND OWNERSHIP AND EXISTING HABITAT PROTECTION**

Grand Caverns is a gated, commercially operated cave, of which Madison Saltpetre Cave is a disjunct section. General Davis Cave is owned by The Nature Conservancy.

## **SUMMARY OF MANAGEMENT AND CONSERVATION ACTIVITIES**

There are no species specific management activities for Trichopetalum weyerienseis.

The existing (1985) Monongahela Land and Resource Management Plan does not provide management direction for caves although they are being considered in the Forest Plan revision currently underway. A Forest Plan Amendment in progress for Threatened and Endangered Species will include management for the caves on the forest.

## **RESEARCH AND MONITORING**

Holsinger, et. al. (1976) reported on a bioinventory of West Virginia cave fauna that encompassed collections from 190 caves in 14 counties. Most of what is known about Trichopetalum weyerienseis in the area of the Monongahela National Forest was gathered during that project. A similar bioinventory was conducted in Virginia caves by Holsinger and Culver (1988).

## **RECOMMENDATIONS**

Retain on list of Regional Forester Sensitive Species.

## **REFERENCES**

- Barr, Thomas C., Jr. 1973. Refugees of the Ice Age. *Natural History*, 82 (5): 26-35, 72-73.
- Duchon, K. and E.A. Lisowski. 1980. Environmental assessment of Lock and Dam Six, Green River navigation project, on Mammoth Cave National Park. Cave Research Foundation, Dallas, Texas, 58 pages.
- Elliott, William R. 1998. Conservation of the North American cave and karst biota. *Subterranean Biota (Ecosystems of the World)*. Elsevier Science. Electronic preprint at [www.utexas.edu/depts/tnhc/.www/biospeleology/preprint.htm](http://www.utexas.edu/depts/tnhc/.www/biospeleology/preprint.htm). 29 pages.

- Harvey, S.J. and J. Skeleton. 1968. Hydrogeologic study of a waste-disposal problem in a karst area at Springfield, Missouri. U.S. Geological Survey Professional Paper 600-C: C217-C220.
- Holsinger, John R., Baroody, Roger A. and David C. Culver. 1976. The invertebrate cave fauna of West Virginia. West Virginia Speleological Survey Bulletin 7, 82 pages.
- Holsinger, John R. and David C. Culver. 1988. The invertebrate cave fauna of Virginia and a part of eastern Tennessee: Zoogeography and Ecology. *Brimleyana*, 14: 1- 162.
- Keith, J.H. 1988. Distribution of Northern cavefish, Amblyopsis spelaea DeKay, in Indiana and Kentucky and recommendations for its protection. *Natural Areas Journal*, 8 (2): 69-79.
- Keith, J.H. and T.L. Poulson. 1981. Broken-back syndrome in Amblyopsis spelaea, Donaldson-Twin Caves, Indiana. Cave Research Foundation 1979 Annual Report, 45-48.
- Lewis, Julian J. 1993. Life returns to Hidden River Cave: The rebirth of a destroyed cave system. *National Speleological Society News*, (June) 208-213.
- Lewis, Julian J. 1996. The devastation and recovery of caves affected by industrialization. Proceedings of the 1995 National Cave Management Symposium, October 25-28, 1995, Spring Mill State Park, Indiana: 214-227.
- Loomis, Harold F. 1939. The millipeds collected in Appalachian caves by Mr. Kenneth Dearolf. *Bulletin of the Museum of Comparative Zoology*, 86: 165-193.
- Panno, S. V., I.G. Krapac, C.P. Weibel and J.D. Bade. 1996. Groundwater contamination in karst terrain of southwestern Illinois. *Illinois Environmental Geology Series EG 151*, Illinois State Geological Survey, 43 pages.
- Panno, S.V., C.P. Weibel, I.G. Krapac and E.C. Stormont. 1997. Bacterial contamination of groundwater from private septic systems in Illinois' sinkhole plain: regulatory considerations. Pages 443-447 In B.F. Beck and J.B. Stephenson (eds.). *The engineering geology and hydrology of karst terranes. Proceedings of the sixth multidisciplinary conference on sinkholes and the engineering and environmental impacts on karst.* Spring, Missouri.
- Panno, S.V., W.R. Kelly, C.P. Weibel, I.G. Krapac, and S.L. Sargent. 1998. The effects of land use on water quality and agricultural loading in the Fogelpole Cave groundwater basin, southwestern Illinois. Proceedings of the Illinois Groundwater Consortium Eighth Annual Conference, Research on agriculture chemicals in Illinois groundwater, 215-233.

Quinlan, J.F. and D.R. Rowe. 1977. Hydrology and water quality in the central Kentucky karst. University of Kentucky Water Resources Research Institute, Research Report 101, 93 pages.

Quinlan, J.F. and D.R. Rowe. 1978. Hydrology and water quality in the central Kentucky karst: Phase II, Part A. Preliminary summary of the hydrogeology of the Mill Hole sub-basin of the Turnhole Spring groundwater basin. University of Kentucky Water Resources Research Institute, Research Report 109, 42 pages.

Schubart, O. 1934. Tausendfüssler oder Myriapoda. 1: Diplopoda: In Die Tierwelt Deutschlands, 28 Teil. Jena: Gustav Fischer, 318 pages.

Shear, William A. 1971. The milliped Family Conotylidae in North America, with a description of the new Family Adritylidae (Diplopoda: Chordeumida). Bulletin of the Museum of Comparative Zoology, 141(2): 55-97.

Shear, William 1972. Studies in the milliped Order Chordeumida (Diplopoda): A revision of the Family Cleidogonidae and reclassification of the Order Chordeumida in the New World. Bulletin of the Museum of Comparative Zoology, 144: 151-352.