

Table G-WS-8a. Mapping units of the Terrestrial Ecological Unit Inventory* on the Chippewa National Forest.

| ELT group# | ELT Group Description | ELT's from '86 Forest Plan |
|--|--|---|
| 1 | Droughty, low fertility sites | B28, B29, F72, F 73, K22, K23, K24, L42, L43, M58, M59, M60, M61, N77, N78, N79 |
| 2 | Wet, medium fertility sites | B26, B27, F78, F79, K32, K33, K34, L48, L50, M68, N83, N84 |
| 3 | Moist, fertile sites | K27, K28, K29, K31, L46, M65, M67, N82 |
| 4 | Wet, fertile sites | K30, K35, L47, L49, M66, M69 |
| 5 | Moist, medium fertility sites | B30, B31, F76, F77, K25, L45, M63, M64, N81 |
| 6 | Dry, medium fertility sites, glacial tills | B32, B33, F74, F75, K26, L44, M62, N80, |
| 7 | Very wet, medium fertility sites | B34, B37, F80, F81, F83, K36, K37, L51, L53, L54, M70, M72, M73, N85, N86, N87 |
| 8 | Very wet, low fertility sites, organic soil sites | B35, B36, F82, F84, K38, K39, L52, L55, M71, M74, N88, N89 |
| Phase Group# | Phase Group Description | Phases from LTP Mapping ** |
| 9 | Excessively well-drained or somewhat excessively well-drained, glacial outwash, dune sand or lacustrine sand, low fertility sites. | b, c, g |
| 10 | Well-drained, stratified sands, silts and glacial till, medium fertility sites. | d, e |
| 11 | Moderately well-drained or well-drained glacial till, medium fertility sites | h |
| 12 | Well-drained, lacustrine, high fertility sites | k, t |
| 13 | Moderately well-drained, lacustrine, high fertility sites | s |
| 14 | Moderately well-drained or well-drained, sand over glacial till, medium fertility sites | hs, qs, p, r |
| 15 | Moderately well-drained, glacial till, medium fertility sites | i, q |
| 16 | Moderately well-drained wind-blown sands, medium fertility sites | f |
| 17 | Poorly drained or somewhat-poorly drained medium fertility sites | j, ba |
| 18 | Poorly drained and very poorly drained forested wetlands, low fertility sites | bb, bs, ta, wc |
| <p>** Each LT phase can be found on different LTA's. For example, phase h can be found on the Blackduck till plain (Btp) and the Itasca moraine (Im). For the purpose of this table they are regarded as the same, but there will be some variation in soil properties between LTA's.</p> <p>*Changes in the developing system include a new Landtype Phase (LTP) layer. In some Landtype Associations (LTA's), LTP mapping is completed and will replace the Ecological Landtype (ELT) outlined in the Ecological Classification System Handbook, 1985. In those LTA's where phase unit mapping has not been completed, use of the ELT's will continue. Until the entire Forest is mapped to the scale of LTP's, both mapping conventions will be used to apply the management direction found in Table G-WS-8. As the phase level inventories are completed, the interpretations will become available to resource managers. The table above provides the key to using map units from either system.</p> | | |

