



April

SUN

MON

TUE

WED

THU

FRI

SAT

1st Week

2nd Week

3rd Week

Garlic mustard emerging

4th Week

Garlic mustard - herbicide, flame torch, or pull

NOTES ON TREATMENT

- **Garlic mustard** - Treatment with glyphosate (roundup) from April to just prior to seed set (about end April)
- **Equipment**— Check all equipment to make sure it is functional
- **Supplies**—Check herbicide supply and order more

CHEQUAMEGON-NICOLET

Chequamegon-Nicolet



May

SUN

MON

TUE

WED

THU

FRI

SAT

1st Week

Garlic mustard - herbicide, or pull

2nd Week

Garlic mustard - herbicide, or pull

3rd Week

Garlic mustard - herbicide, or pull (bag if pods are starting to form)
Canada thistle - rosettes appear

4th Week

Garlic mustard - herbicide, or pull (bag if pods are starting to form)

NOTES ON TREATMENT

- **Garlic mustard** - Treatment with glyphosate (roundup) from April to just prior to seed set (about end April)
- **Garlic mustard** - manual control until seeds form. Pull or cut low to ground with a weed-whacker
- **3rd week** - Siberian Pea shrub in bloom - easy to see in woods

CHEQUAMEGON-NICOLET

Chequamegon-Nicolet



June

SUN

MON

TUE

WED

THU

FRI

SAT

1st Week

Canada thistle - herbicide or manual
Purple loosestrife - Galerucella beetle release

2nd Week

Canada thistle - herbicide or manual
Leafy spurge - weed whacker (if tall enough)

3rd Week

Canada thistle - herbicide or dig or first mowing of 3
Leafy spurge - weed whacker
Wild parsnip - herbicide or manual
Spotted knapweed - herbicide

4th Week

Wild parsnip - herbicide or manual
Spotted knapweed - herbicide or mow
Canada thistle - herbicide or manual

NOTES ON TREATMENT

- **1st week** -Canada thistle is about 8-10 inches tall, very susceptible to herbicide .clopyralid or glyphosate when 6-10 " tall
- **Canada thistle** - Manual pull or mow until early bud stage (mid June)
- **Wild Parsnip** - Manual control by pulling (wear gloves) or by pushing blade of shovel to cut root just below ground
- **Spotted knapweed** - Transline (clopyralid) from mid-June to early July. Roundup Pro (glyphosate) from early growth to bolting stage.
- **Wild parsnip** - Basal rosettes and plants until seed set: pull or herbicide . June - July
- **Leafy spurge** - When plants are a foot or so tall, whack them down so they will re-sprout; they get herbicide in September

CHEQUAMEGON-NICOLET

Chequamegon-Nicolet



July

SUN

MON

TUE

WED

THU

FRI

SAT

1st Week

Leafy spurge - beetle release

Spotted knapweed - herbicide or mow

2nd Week

Spotted knapweed - herbicide or mow

Canada thistle - continue mowing

3rd Week

Spotted knapweed - herbicide or mow

Phragmites - herbicide or cut (need weed trimmer w/ blade)

4th Week

Phragmites - herbicide or cut (need weed trimmer w/ blade)

Purple loosestrife - herbicide

NOTES ON TREATMENT

- **Spotted knapweed** - herbicide or mow until prior to seed set
- **Purple loosestrife** - herbicide when in bloom (easier to find)
- **Leafy spurge** - Release beetles in early July
- **Oriental bittersweet** - herbicide (Garlon , Vine-X, or glyphosate) July thru October. Identification easiest in summer so treat then, or flag for fall treatment.
- **Canada thistle** - If mowing or pulling, should be done at least 2 times (same patch) 3 is better

CHEQUAMEGON-NICOLET

Chequamegon-Nicolet



August

SUN

MON

TUE

WED

THU

FRI

SAT

1st Week

Purple loosestrife - herbicide

2nd Week

Purple loosestrife - herbicide

3rd Week

Buckthorn, Honeysuckle, Barberry, & Pea shrub - begin herbicide
Canada thistle - third mowing

4th Week

Begin Leafy spurge - herbicide with Plateau (imazapic)

NOTES ON TREATMENT

- **Purple loosestrife** - treat manual or herbicide until prior to seed set.
- **Oriental bittersweet** - Herbicide July to October
- **Leafy spurge** - Plateau (imazapic) is applied quite late in the season;
- **Canada thistle** - third mowing in mid to late August

mid September is best

CHEQUAMEGON-NICOLET



September

SUN

MON

TUE

WED

THU

FRI

SAT

1st Week

Leafy spurge - herbicide with Plateau (imazapic)

Buckthorn, Honeysuckle, Barberry, Autumn Olive, Bittersweet, & Pea shrub - herbicide

2nd Week

Leafy spurge - herbicide - Plateau (imazapic)

Buckthorn, Honeysuckle, Barberry, Autumn Olive, Bittersweet, & Pea shrub - herbicide

3rd Week

Buckthorn, Honeysuckle, Barberry, Autumn Olive, Bittersweet, & Pea shrub - herbicide

Leafy spurge - herbicide - Plateau (imazapic)

Swamp thistle - herbicide new rosettes or dig

4th Week

Buckthorn, Honeysuckle, Barberry, Autumn Olive, Bittersweet, & Pea shrub - herbicide

Leafy spurge - herbicide - Plateau (imazapic)

NOTES ON TREATMENT

- **Leafy spurge** - herbicide with Plateau (imazapic) clopyralid to new rosettes which over -winter
- **Buckthorn, Honeysuckle, Barberry, Siberian peashrub-** Herbicide (Roundup, Garlon)
- **Swamp thistle** - glyphosate or

CHEQUAMEGON-NICOLET



October

SUN

MON

TUE

WED

THU

FRI

SAT

1st Week

Buckthorn, Honeysuckle, Barberry, Autumn Olive, Bittersweet, & Pea shrub - herbicide

2nd Week

Buckthorn, Honeysuckle, Barberry, Autumn Olive, Bittersweet, & Pea shrub - herbicide

3rd Week

4th Week

NOTES ON TREATMENT

- **Shrubs** - herbicide until temps reach below 30 degrees F. Use Garlon

CHEQUAMEGON-NICOLET



November

SUN

MON

TUE

WED

THU

FRI

SAT

1st Week

Buckthorn, Honeysuckle, Barberry, Autumn Olive, Bittersweet, & Pea shrub - herbicide

2nd Week

3rd Week

4th Week

NOTES ON TREATMENT

- **Shrubs** - herbicide until temps drop below 30 degrees F

CHEQUAMEGON-NICOLET

Chequamegon-Nicolet