

# Montana Forest Insect and Disease Conditions and Program Highlights

# 2009

United States



Department of  
Agriculture

Forest Service

Northern Region

Forest Health  
Protection


Report 10-1



Montana  
Department of  
Natural Resources  
and Conservation  
Forestry Division



**Mountain Pine Beetle in  
Whitebark Pine**



**Rhabdocline Needle Cast in  
Douglas-fir**

“The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all of part of an individual’s income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large prints, audiotape, etc.) should contact USDA’s TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, DC 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.”

# **MONTANA**

## **Forest Insect and Disease Conditions and Program Highlights – 2009**

**Report 10-01**

**2010**

**Compiled By:**

**Amy Gannon, Montana Department of Natural Resources and Conservation,  
Forestry Division**

**Scott Sontag, USDA Forest Service, Northern Region, State and Private  
Forestry, Forest Health Protection**

**Contributors:**

**Gregg DeNitto, Ken Gibson, Marcus Jackson, Blakey Lockman, Scott Sontag,  
Brytten Steed, and Nancy Sturdevant, of the USDA Forest Service, Northern  
Region, State and Private Forestry, Forest Health Protection;**

**Amy Gannon of the Montana Department of Natural Resources and  
Conservation, Forestry Division; Brennan Ferguson of Ferguson Forest  
Pathology Consulting Inc.**

**Data Summary:**

**William O'Donnell, USDA Forest Service, Region One, Engineering**

**Map Production:**

**International Forestry Consultants Inc.**

**Cover Photos:**

**Mountain Pine Beetle in Whitebark Pine, by Ken Gibson, USDA Forest Service**

**Rhabdocline Needle Cast in Douglas-fir, by Scott Sontag, USDA Forest Service**

## TABLE OF CONTENTS

|  |           |
|--|-----------|
| <b>ABBREVIATIONS</b> .....                           | <b>1</b>  |
| <b>INTRODUCTION</b> .....                            | <b>1</b>  |
| <b>SUMMARY OF CONDITIONS</b> .....                   | <b>2</b>  |
| <b>Bark Beetles</b> .....                            | <b>2</b>  |
| <b>Defoliators</b> .....                             | <b>3</b>  |
| <b>Root Diseases</b> .....                           | <b>3</b>  |
| <b>Foliage Diseases</b> .....                        | <b>4</b>  |
| <b>Dwarf Mistletoes</b> .....                        | <b>4</b>  |
| <b>White Pine Blister Rust</b> .....                 | <b>4</b>  |
| <b>Abiotic Damage</b> .....                          | <b>5</b>  |
| <b>ANNUAL AERIAL SURVEY</b> .....                    | <b>5</b>  |
| <b>INSECT AND DISEASE CONDITIONS BY COUNTY</b> ..... | <b>7</b>  |
| <b>COMMON AND SCIENTIFIC NAMES</b> .....             | <b>39</b> |
| <b>PUBLICATIONS 2009</b> .....                       | <b>40</b> |
| <b>DIRECTORY OF PERSONNEL</b> .....                  | <b>44</b> |

## LIST OF TABLES

|                |   |           |
|----------------|---|-----------|
| <b>Table 1</b> | <b>Mortality, Defoliation and Other Damage detected from the air on National Forests, National Parks and Tribal Lands in Montana, during 2009</b> ..... | <b>34</b> |
| <b>Table 2</b> | <b>Acres of Host Type Infested by Bark Beetles on All Ownerships in Montana, during 2009</b> .....  | <b>46</b> |
| <b>Table 3</b> | <b>2009Acres of Host Type Infested by Bark Beetles in Montana, From 2007 Through 2009</b> .....   | <b>47</b> |
| <b>Table 4</b> | <b>Acres with Douglas-fir Beetle-Caused Mortality on All Ownerships in Montana, From 2007 Through 2009</b> .....  | <b>48</b> |
| <b>Table 5</b> | <b>Acres with Mountain Pine Beetle-Caused Mortality on State and Private Lands in Montana, From 2007 Through 2009</b> .....                             | <b>49</b> |
| <b>Table 6</b> | <b>Acres with Mountain Pine Beetle-Caused Mortality on All Federal Ownerships in Montana, From 2007 Through 2009</b> .....                              | <b>50</b> |
| <b>Table 7</b> | <b>Acres with Additional Bark Beetle-Caused Mortality on All Ownerships in Montana, From 2007 Through 2009</b> .....                                    | <b>51</b> |

## LIST OF FIGURES

|                 |   |           |
|-----------------|---|-----------|
| <b>Figure 1</b> | <b>Reporting Areas and County Boundaries in Montana</b> .....   | <b>52</b> |
| <b>Figure 2</b> | <b>Areas Surveyed During the 2009 Forest Health Protection Aerial Detection Survey in Montana</b> ..... | <b>53</b> |
| <b>Figure 3</b> | <b>2009 Mountain Pine Beetle Infestations in Montana</b> .....  | <b>54</b> |
| <b>Figure 4</b> | <b>2009 Douglas-fir Beetle Infestations in Montana</b> .....  | <b>55</b> |
| <b>Figure 5</b> | <b>2009 Fir Engraver Beetle Infestations in Montana</b> .....   | <b>56</b> |
| <b>Figure 6</b> | <b>2009 Subalpine Fir Mortality Complex in Montana</b> .....  | <b>57</b> |
| <b>Figure 7</b> | <b>2009 Western Spruce Budworm Infestations in Montana</b> .....  | <b>58</b> |

## ABBREVIATIONS

The following abbreviations are used throughout this document:

|                    |      |   |  |
|--------------------|------|---|--|
| <b>Beetles</b>     | DFB  | = | Douglas-fir beetle, <i>Dendroctonus pseudotsugae</i> Hopkins           |
|                    | ESB  | = | Spruce beetle, <i>D. rufipennis</i> (Kirby)                            |
|                    | IPS  | = | Pine engraver, <i>Ips pini</i> (Say)                                   |
|                    | MPB  | = | Mountain pine beetle, <i>D. ponderosae</i> Hopkins                     |
|                    | WPB  | = | Western pine beetle, <i>D. brevicomis</i> LeConte                      |
|                    | FE   | = | Fir engraver, <i>Scolytus ventralis</i> LeConte                        |
|                    | WBBB | = | Western balsam bark beetle, <i>Dryocoetes confuses</i> Swaine          |
|                    | RTB  | = | Red turpentine beetle, <i>D. valens</i> LeConte                        |
| <b>Defoliators</b> | WSBW | = | Western spruce budworm, <i>Choristoneura occidentalis</i> Freeman      |
|                    | LCB  | = | Larch casebearer, <i>Coleophora laricella</i> Hübner                   |
|                    | DFTM | = | Douglas-fir tussock moth, <i>Orygia pseudotsugata</i> McDunnough       |
|                    | PTM  | = | Western Pine (Grizzled) tussock moth, <i>Dasychira pinicola</i> (Dyar) |
| <b>Hosts</b>       | LPP  | = | Lodgepole pine   |
|                    | PP   | = | Ponderosa pine   |
|                    | WWP  | = | Western white pine   |
|                    | WBP  | = | Whitebark pine   |
|                    | LP   | = | Limber pine  |
|                    | DF   | = | Douglas-fir  |
|                    | WL   | = | Western larch  |
|                    | GF   | = | Grand fir  |
|                    | SAF  | = | Subalpine fir  |
|                    | ES   | = | Engelmann spruce   |
| <b>Other</b>       | NF   | = | National Forest  |
|                    | RD   | = | Ranger District  |
|                    | IR   | = | Indian Reservation   |
|                    | NP   | = | National Park  |
|                    | BLM  | = | Bureau of Land Management  |
|                    | FIA  | = | Forest Inventory and Analysis  |
|                    | RA   | = | Reporting Area   |

## INTRODUCTION

This report summarizes the major forest insect and disease conditions in Montana during 2009 and was jointly prepared by the Montana Department of Natural Resources and Conservation, Forestry Division (DNRC) and the USDA Forest Service, State and Private Forestry, Forest Health Protection, Northern Region (FHP).

Information for this report was derived from ground and aerial detection surveys (ADS) within RAs across parts of Montana. A RA includes all federal, state, and private land ownerships within a particular geographic boundary (Figure 1).

## SUMMARY OF CONDITIONS

### Bark Beetles

Weather conditions were near normal in 2009. Neither winter nor summer temperatures were extreme; and summer precipitation was only slightly less than normal. We did experience unusually cold temperatures in early October, but it is too early to tell if there were areas in the Region where overwintering beetle larvae were not yet prepared for such cold conditions. It is possible that some areas could experience a brief set-back in beetle population development. However, even should that occur, beetle populations will rebound so long as susceptible stand conditions remain. MPB populations remained high in most infested areas; SAF stands continued to be damaged by a host of factors—one being WBBB; mortality attributed to FE declined markedly; and DFB-infested stands increased in a few areas. Most other bark beetle species were found at relatively low levels throughout the state.

In 2009, we flew most of the forested areas in Montana, with the exception that most wilderness areas were not surveyed. Approximately 27.8 million acres were surveyed.

MPB populations, highest in the west-central portion of Montana, continued to increase in 2009; but not at rates suggested by acreage figures derived from aerial surveys. On parts of the Helena and Deerlodge NFs, beetle populations have begun or continued to decline because of host depletion. Beetle-infested areas increased significantly in a few locations, and populations expanded into some previously un-infested areas—notably

ones on the western Deerlodge, Beaverhead, Lewis and Clark, southern Flathead, western parts of the Lolo, and Gallatin NFs. Considerably more acres of PP with MPB were detected in 2009, especially around Helena. Overall, decreases were recorded in infested stands on portions of the Flathead, Lolo, and Deerlodge NFs. In summary, MPB-infested acres in LPP stands increased dramatically in 2009—up from just over 1.5 million acres recorded in 2008, to slightly more than 2.7 million acres in 2009.

DFB-infested acres remained about the same in western Montana in 2009. At a few locations in northwestern and central Montana, populations remained at low-epidemic levels; in other areas they declined considerably. Decreases were noted in most areas, except in the Pryor Mountains on the Custer NF. Declines were especially noted on the Flathead, Helena, and Lolo NFs. Infested acres increased, but only slightly in 2009 from levels recorded in 2008—from 21,500 acres to 22,500 acres. In most areas, DFB populations remain at nearly endemic levels.

GF mortality attributable to FE increased considerably in 2009, but we believe that is attributed to more acres being flown and more low intensity levels recorded. Infested area increased from 360 acres in 2008 to almost 9,200 acres in 2009, but the severity was less than 1 tree per acre on average.

SAF mortality, likely attributed to WBBB, was mapped on more acres in 2009, likely a result of more infested areas being surveyed. Notable beetle-caused mortality was found in some areas as affected SAF acreage increased from 52,700 acres to just over 79,500 acres.

Mortality in PP stands, attributed to IPS and WPB both decreased in 2009. Both were found at relatively low levels, likely due to nearly normal amounts of moisture received the past couple of years.

Fire activity was once again low during the summer of 2009, so few fire/beetle interactions should be encountered in 2010. WSBW populations, however, increased significantly in many areas and we may observe more severely defoliated trees being killed by bark beetles. While drought effects have ameliorated somewhat, cumulative effects of prolonged dry weather in many parts of the state have taken their toll and bark beetle populations are quite adept at taking advantage of any opportunity presented them. Dependent to a large extent on weather for the remainder of the winter and into spring, some bark beetle populations could rebound in 2010. MPB populations, more reflective of host conditions than weather, likely will continue to expand into and within susceptible host stands and decrease in stands already severely affected.

## **Defoliators**

WSBW continues to be the most significant defoliator in the state. In 2009, 2,554,205 acres were mapped as defoliated by WSBW in Montana. Most of this defoliation was in Beaverhead, Flathead, Gallatin, Judith Basin, Lewis and Clark, Lincoln, Meagher, Powell, and Sanders Counties with each reporting over 100,000 acres. Defoliation was widespread in northwestern and central Montana affecting mainly DF. A few areas with multiple years of defoliation are beginning to see tree mortality in

association with DFB. Continued defoliation, especially with droughty conditions, could cause an increase in DF mortality.

Defoliation by DFTM and LCB were not reported in Montana in 2009. Areas with defoliation by PTM had been reported in recent years.

Cooperative detection monitoring continued for the gypsy moth, *Lymantria dispar*, in 2009 with USDA Animal and Plant Health Inspection Service (APHIS), Montana Department of Agriculture (MDA), Montana Department of Natural Resources and Conservation (DNRC), and USDA Forest Service. A network of more than 1,000 pheromone-baited traps were placed throughout Montana's forests and urban areas most frequented by travelers.

## **Root Diseases**

Root diseases are diseases of the site and do not change drastically from one year to the next. Root disease fungi cause damage and mortality on more than 7 million acres in western Montana, killing more than 30 million trees annually. Root disease-caused mortality is more common west of the Continental Divide. Large areas of root disease can be found east of the Divide, but it tends to occur in more discrete patches, rather than being ubiquitous throughout an area. Also, root diseases can be commonly found in riparian areas east of the Continental Divide, often in ES and SAF.

## **Foliage Diseases**

Over 2,000 acres of Rhabdocline needle cast of DF were identified by ADS in Sanders and Lake Counties on the western edge of the Salish Mountains. About two-thirds of the affected acres were on Tribal lands and one-third was on private lands. ADS also identified WL needle blight on about 60 acres west of Noxon.

## **Dwarf Mistletoes**

Historical assessments of dwarf mistletoe stands in Montana show that about 16% of DF, 33% of LPP, and 30% of WL stands are infested with dwarf mistletoe. More recent assessments using FIA data collected on over 50,000 trees across Montana show that about 1% of DF, 3.2% of LPP, and 3.9% of WL trees were infected. The clumpy nature, and protracted disease-cycle, of dwarf mistletoes allow for impacts across an individual stand to range from none to severe. The clumpy nature of dwarf mistletoe is likely the reason for a discrepancy between these two measurements; the first is measuring infected stands and the second is measuring infected trees.

## **White Pine Blister Rust**

WWP, WBP, and LP are all susceptible to white pine blister rust.

### *Western white pine*

Mature WWP continue to be lost due to a combination of blister rust and MPB. Lack of suitable sites, either man-made or natural, limits natural regeneration, and white pine blister rust may kill a high

proportion of natural seedlings.

Resistant stock is planted operationally on suitable WWP sites on federal, state, industrial, and private forest lands throughout northwestern Montana. In addition to planting rust-resistant stock, pruning of the lower bole is an important tool used in blister rust management. In order to encourage initiation of pruning programs a two-day workshop was held for federal and state foresters in April of 2009 in Libby, Montana, focusing on the operational aspects of how to carry out pruning in WWP plantations.

### *Whitebark pine*

White pine blister rust has been impacting WBP ecosystems for many decades. In addition, recent outbreaks of MPB have caused widespread mortality in many WBP stands already impacted by white pine blister rust. The combination of white pine blister rust, bark beetle outbreaks, and lack of natural regeneration due to fire suppression has raised concerns about the long-term viability of WBP ecosystems.

A recent survey by FHP of WBP stands throughout Idaho and Montana found that of the 42 sampled stands, 69% have less WBP regeneration than that of other species such as SAF and ES. White pine blister rust infection varied from 0% to 81% in these stands, and 40% had infection levels greater than 20%. Taking into account the ongoing impacts of white pine blister rust and bark beetles, the report concludes that approximately 75% of these stands will convert to other species without active restoration efforts or the occurrence of wildfire.

Standardized methodology is now being used to establish monitoring plots in

WBP stands throughout the West. These plots are designed to provide a statistically-based assessment of the incidence of white pine blister rust in the ecosystem and the condition of WBP. A database has been developed (<http://www.fs.fed.us/r1-r4/spf/fhp/prog/programs2.html>) to compile and provide results of surveys in both WBP and LP.

### Limber pine

A report describing the current amount, distribution, and condition of LP in Montana will be available in early 2010. Authored by USFS FHP, the Montana DNRC, and the U.S. Geological Survey, it will also provide information on protection and restoration of this ecologically important species.

LP seeds were collected from three Montana sites in 2009 for inclusion in a West-wide collection being carried out for gene conservation, white pine blister rust resistance screening, and various research projects. Seed will be collected from several additional locations in Montana in 2010 if funding is available and cone yields are adequate.

### 2010 conference in Missoula

In June of 2010 a major conference titled “*The Future of High-Elevation Five-Needle White Pines in Western North America*” will be held in Missoula, Montana. See the following web site for agenda and registration information: <http://www.umt.edu/ce/cps/highfive/>.

### **Abiotic Damage**

Nearly 800 acres of windthrow and 150 acres of avalanche-caused tree damage were mapped in 2009. Most windthrow

occurred in a band that stretches from north-central Montana to southwestern Montana.

### **ANNUAL AERIAL SURVEY**

The annual aerial detection survey in Montana was conducted from June 29 thru September 26, 2009. The survey covered approximately 27.8 million acres of mixed ownership, forested lands, excluding most wilderness areas (Figure 2). Five FHP sketchmappers, using three aircraft, conducted the 2009 aerial survey.

Much of the data summarized in this report is a product of the aerial surveys, as well as ground surveys and biological evaluations. Along with the data summaries, aerial survey maps are available from the Missoula FHP Field Office, in both paper and digital GIS format. Data may also be downloaded at <http://www.fs.fed.us/r1-r4/spf/fhp/aerial/index.html>.

The annual aerial detection survey is an overview survey designed to cover large areas in a relatively short period of time. Aerially detected signatures include tree mortality, defoliation and windthrow. If recent forest disturbance activities are low, other disturbances such as diseases, needle casts, high-water damage and previous fire damage are mapped. The intent of the survey is to cover each area once a year during which time the observer maps as many disturbances as possible. The survey is conducted using single-engine, high-wing aircraft, flying at speeds of approximately 90 to 130 mph, at an average altitude of approximately 1,000 to 2,000 feet above ground level.

The aerial survey data are estimates made from airplanes and though not as many areas were ground checked as we would like, enough were checked to lend confidence to the areas for which we only have aerial survey data. Together, aerial and ground surveys

provide information relative to bark-beetle-caused mortality, as well as other damage agents pertinent to land managers charged with the responsibility of maintaining forest health.

## INSECT AND DISEASE CONDITIONS BY COUNTY

### County Areas in Detail

County summaries follow. For each, damage effects on their respective ownership are noted. To the extent possible, we have indicated areas affected and an estimate of extent. Counties not listed had no reported information. Forestland data in the following are from the annualized surveys performed by USDA Forest Service FIA (<http://www.fia.fs.fed.us>). In some of our tables, you will observe acres of damage on some ownership where there is no forestland reported. Because of the limited forestland in ownerships in some Counties, FIA data may indicate no forestland acres at the County level. This is a result of the intensity of the inventory performed. This discrepancy is within their standard of error. Other Federal lands include tribal ownership.

### Beaverhead County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State  | Total     |
|---------------------------|-----------------|---------------|---------|--------|-----------|
| <b>Forestland</b>         | 1,073,844       | 135,995       | 40,745  | 17,003 | 1,267,587 |
| <b>Aspen Decline</b>      | 0               | 283           | 63      | 0      | 346       |
| <b>DFB</b>                | 51              | 14            | 4       | 0      | 69        |
| <b>ESP</b>                | 2               | 4             | 0       | 0      | 6         |
| <b>MPB-LPP</b>            | 134,180         | 16,526        | 4,909   | 2,710  | 158,325   |
| <b>MPB-High Elevation</b> | 63,854          | 4,803         | 828     | 747    | 70,232    |
| <b>SAF Mortality</b>      | 134             | 6             | 0       | 4      | 144       |
| <b>WSBW</b>               | 54,263          | 31,955        | 15,243  | 9,243  | 110,704   |
| <b>Windthrow</b>          | 25              | 0             | 0       | 0      | 25        |

MPB continues to increase in Beaverhead County, particularly in five-needle pines. New activity in LPP is particularly notable in the Big Hole area and the southern portion of the Pioneer Mountains. Large groups were mapped in the Pioneer and Beaverhead Mountains to the NW and SW of Dillon, respectively. FINDITs plot summaries from recently infested areas show 2-41% of LPP currently dead. Based on previous trends some areas can be expected to have up to 90% mortality of trees >5" diameter. On BLM administered lands in the Centennial Mountain Range, MPB-killed LPP and WBP were at reduced levels from just a few years ago. WSBW caused defoliation continues to be present over much of the County with increases most notable on USFS managed lands. DFB remains low, with only a few acres of newly found ESB activity. Also of note is the high level of SAF mortality attributed to WBBB, with significantly more acres noted in 2009. Schweinitzii root and butt rot is common in DF, causing decay in the butt logs but not acting as an aggressive root pathogen. LPP dwarf mistletoe and LP dwarf mistletoe are present in the County. White pine blister rust is common in WBP. A 25-acre patch of windthrow was mapped on Forest Service administered lands about 12 miles north of Lima Reservoir.

## Big Horn County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State  | Total   |
|---------------------------|-----------------|---------------|---------|--------|---------|
| <b>Forestland</b>         | 0               | 0             | 285,165 | 39,586 | 324,750 |
| <b>Aspen Decline</b>      | 0               | 159           | 66      | 2      | 227     |
| <b>DFB</b>                | 0               | 125           | 0       | 0      | 125     |
| <b>MPB-LPP</b>            | 0               | 4             | 0       | 0      | 4       |
| <b>MPB-PP</b>             | 0               | 113           | 29      | 0      | 142     |
| <b>MPB-High Elevation</b> | 0               | 14            | 0       | 0      | 14      |
| <b>SAF Mortality</b>      | 0               | 4             | 0       | 0      | 4       |
| <b>WSBW</b>               | 0               | 916           | 75      | 0      | 991     |

In 2009, very little bark beetle activity was noted anywhere, with most detected on the Northern Cheyenne IR. Minor amounts of MPB-killed PP and DFB-killed DF were mapped at endemic levels. The most prevalent insect problem was WSBW with activity noted on both tribal and privately managed lands. LPP dwarf mistletoe and LP dwarf mistletoe are present in the County.

## Blaine County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                   | National Forest | Other Federal | Private | State | Total   |
|-------------------|-----------------|---------------|---------|-------|---------|
| <b>Forestland</b> | 0               | 83,825        | 107,216 | 0     | 191,042 |
| <b>MPB-LPP</b>    | 0               | 153           | 0       | 10    | 163     |
| <b>MPB-PP</b>     | 0               | 82            | 2       | 0     | 84      |
| <b>Windthrow</b>  | 0               | 26            | 0       | 0     | 26      |

Scattered LPP and PP mortality from MPB were recorded in the Little Rocky Mountains on the Fort Belknap IR. Likely some of this mortality in PP has occurred in conjunction with IPS. LPP dwarf mistletoe is present in the County. One 26-acre patch of windthrow was noted on the Fort Belknap Reservation, located in the Little People's drainage.

## Broadwater County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State | Total   |
|---------------------------|-----------------|---------------|---------|-------|---------|
| <b>Forestland</b>         | 155,612         | 20,806        | 101,116 | 0     | 277,534 |
| <b>DFB</b>                | 4               | 2             | 4       | 0     | 10      |
| <b>MPB-LPP</b>            | 38,096          | 79            | 7,677   | 10    | 45,862  |
| <b>MPB-PP</b>             | 442             | 744           | 9,166   | 800   | 11,152  |
| <b>MPB-High Elevation</b> | 2,738           | 193           | 1,853   | 0     | 4,784   |
| <b>WSBW</b>               | 27,694          | 1,545         | 17,141  | 270   | 46,650  |

MPB-killed LPP was by far the most significant mortality agent on lands surveyed. Very large groups were mapped in the Big Belt Mountains, near Boulder Mountain, Mount Baldy and southward to Mount Edith. Most stands near Canyon Ferry Lake with host pines show some level of MPB-caused mortality. MPB activity increased significantly in higher elevation WBP forests. MPB activity will probably decrease over the next few years as host is depleted. WSBW increased nearly 4-fold across the County. White pine blister rust has been found in LP. Armillaria root disease is present in the County, and Schweinitzii root and butt rot is quite common, especially causing significant decay in the butt logs of larger, older DF. LPP dwarf mistletoe is present in the County.

## Carbon County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State  | Total   |
|---------------------------|-----------------|---------------|---------|--------|---------|
| <b>Forestland</b>         | 130,122         | 127,055       | 54,590  | 10,092 | 321,860 |
| <b>DFB</b>                | 2,348           | 341           | 138     | 0      | 2,827   |
| <b>MPB-LPP</b>            | 290             | 0             | 2       | 4      | 296     |
| <b>MPB-PP</b>             | 0               | 2             | 8       | 2      | 12      |
| <b>MPB-High Elevation</b> | 7,114           | 771           | 30      | 0      | 7,915   |
| <b>SAF Mortality</b>      | 182             | 0             | 0       | 0      | 182     |
| <b>WSBW</b>               | 643             | 25            | 13      | 0      | 681     |
| <b>Windthrow</b>          | 15              | 0             | 0       | 0      | 15      |

Almost twice as many acres were flown in 2009 as in 2008 in this County. DFB activity increased significantly in 2009 in the Pryor Mountains and elsewhere in the County. However, some of this increase could be in part a result of the increase in acreage flown. MPB activity in LPP increased slightly in 2009 and remained somewhat static in PP stands. However, there was a significant increase in MPB-killed WBP, especially in Rock Creek drainage. White pine blister rust is common in WBP and LP. LPP dwarf mistletoe and LP dwarf mistletoe are present in the County. A small 15-acre patch of blow down was found on Forest Service administered lands at the northeast head of Inferno Canyon in the Pryor Mountains.

## Carter County

Acres of Forestland, Mortality, and Defoliation by Ownership

|               | National Forest | Other Federal | Private | State | Total   |
|---------------|-----------------|---------------|---------|-------|---------|
| Forestland    | 62,418          | 0             | 55,333  | 0     | 117,751 |
| Aspen Decline | 2               | 0             | 4       | 0     | 6       |
| MPB-PP        | 57              | 4             | 32      | 4     | 97      |

Slightly more USFS managed acres were flown in 2009 than 2008. Within these lands MPB and IPS in PP showed slight declines. A few groups of 1-2 impacted trees were recorded in the Slim Buttes, Cave Hills, and Short Pines. A few small groups of MPB-killed PP were recorded in the Ekalaka Hills and in PP stands just west of Camp Crook, but still in minor amounts. It should be noted that at least some tree mortality attributed to MPB may actually be caused by IPS.

## Cascade County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                    | National Forest | Other Federal | Private | State  | Total   |
|--------------------|-----------------|---------------|---------|--------|---------|
| Forestland         | 196,965         | 10,261        | 95,225  | 11,497 | 313,949 |
| DFB                | 6               | 0             | 95      | 0      | 101     |
| MPB-LPP            | 29,922          | 5,014         | 15,659  | 1,079  | 51,674  |
| MPB-PP             | 1,648           | 3,351         | 33,669  | 2,252  | 40,920  |
| MPB-High Elevation | 3,438           | 0             | 777     | 537    | 4,752   |
| SAF Mortality      | 387             | 0             | 0       | 0      | 387     |
| WSBW               | 40,671          | 4,565         | 35,916  | 4,521  | 85,673  |

Number of acres flown almost doubled in 2009 and may thus affect mortality estimates. The 2009 data showed great increases in MPB-caused mortality in LPP, PP and high elevation WBP, although MPB has been active in this area for several years. Many scattered, mostly small groups of MPB-killed PP were mapped throughout the Little Belt Mountains. Groups of MPB-killed WBP were found near Kings Hill as well. DFB activity was low. White pine blister rust is common in LP. Armillaria root disease present in the southeastern portion of the County, and Schweinitzii root and butt rot is quite common, especially causing significant decay in the butt logs of larger, older DF.

## Chouteau County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                   | National Forest | Other Federal | Private | State | Total  |
|-------------------|-----------------|---------------|---------|-------|--------|
| <b>Forestland</b> | 22,295          | 0             | 34,470  | 2,245 | 59,010 |
| <b>DFB</b>        | 0               | 10            | 0       | 0     | 10     |
| <b>MPB-LPP</b>    | 0               | 1,301         | 457     | 253   | 2,011  |
| <b>MBP-PP</b>     | 0               | 400           | 71      | 29    | 500    |
| <b>WSBW</b>       | 0               | 2,599         | 3,122   | 177   | 5,898  |

MPB-killed LPP nearly doubled and was located mostly on tribal lands. The largest increase in MPB activity was in PP stands located on Rocky Boys IR, near Salt Coulee and Sawmill Butte. Most of the MPB-killed LPP stands were near Muddy and Timber Creeks. WSBW activity also significantly increased, again, with the largest increase on the IR in stands with high amounts of DF. DFB activity remained at low levels, but may increase in the future because of WSBW repeated defoliation, especially in stands that have had recent fire activity or have large diameter DF. Armillaria root disease is present in the northern portion of the County, on the Rocky Boys IR, but appears to be uncommon. Butt rot and breakage caused by *Ganoderma applanatum* has been observed in quaking aspen on the Rocky Boys IR.

## Deer Lodge County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State  | Total   |
|---------------------------|-----------------|---------------|---------|--------|---------|
| <b>Forestland</b>         | 213,376         | 9,963         | 45,203  | 27,770 | 296,313 |
| <b>DFB</b>                | 4               | 0             | 5       | 0      | 9       |
| <b>MPB-LPP</b>            | 51,613          | 2,179         | 40,670  | 22,253 | 116,715 |
| <b>MPB-PP</b>             | 0               | 0             | 10      | 0      | 10      |
| <b>MPB-High Elevation</b> | 1,861           | 0             | 576     | 1,054  | 3,491   |
| <b>WSBW</b>               | 3,102           | 0             | 5,530   | 33     | 8,665   |
| <b>Windthrow</b>          | 48              | 0             | 5       | 0      | 53      |

MPB continues to be very active with MPB-caused tree mortality decreasing in the northern portions due to host depletion and increasing in areas south of the Continental Divide. In both 2008 and 2009 similar numbers of acres in each ownership group were surveyed. However, acres of LPP affected by MPB increased by almost 25% with estimated number of trees killed increasing nearly 80%. In LPP stands where MPB activity is increasing, FINDITs surveys show 2-30% total mortality, with up to 60 times more mortality than found in 2008. Far fewer acres of MPB in five-needle pines were noted, especially in private ownership, although the estimate of trees killed actually increased. DNRC personnel hung MPB-baited traps in Anaconda at the MT DNRC Office to determine flight period. Peak collections were on 8/27/09 (36). Acres of

WSBW caused defoliation also increased greatly, with many more acres noted under private ownership. White pine blister rust has been found in LP. Schweinitzii root and butt rot is common in DF, causing decay in the butt logs but not acting as an aggressive root pathogen. LPP dwarf mistletoe and LP dwarf mistletoe are present in the County. One 53-acre patch of windthrow was mapped in just west of Champagne Pass (47 acres on Forest Service administered lands and 5 acres on private land), at the northern edge of the County. There was probably one wind event causing windthrow in Deerlodge, Powell, and Jefferson Counties, very near to where these three counties border each other.

## Fergus County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                              | National Forest | Other Federal | Private | State  | Total   |
|------------------------------|-----------------|---------------|---------|--------|---------|
| <b>Forestland</b>            | 86,611          | 168,352       | 298,106 | 19,491 | 572,560 |
| <b>Aspen Decline</b>         | 0               | 0             | 49      | 0      | 49      |
| <b>DFB</b>                   | 11              | 2             | 23      | 0      | 36      |
| <b>Flooding - High Water</b> | 0               | 0             | 13      | 0      | 13      |
| <b>MPB-LPP</b>               | 1,609           | 446           | 1,278   | 72     | 3,405   |
| <b>MPB-PP</b>                | 5,646           | 588           | 4,216   | 339    | 10,789  |
| <b>MPB-High Elevation</b>    | 6               | 2             | 2       | 0      | 10      |
| <b>SAF</b>                   | 300             | 0             | 0       | 0      | 300     |
| <b>Unidentified Disease</b>  | 43              | 0             | 0       | 0      | 43      |
| <b>WSBW</b>                  | 14,022          | 1,963         | 12,095  | 827    | 28,907  |
| <b>Windthrow</b>             | 0               | 85            | 28      | 24     | 137     |

Nearly 40% more acres were surveyed in 2009, all on BLM, State, and privately managed lands. This may explain some of the increased pest activity on these ownerships. However, data from USFS-managed lands (all acres surveyed both years) suggest nearly a 50-fold and 25-fold increase in LPP and PP acres affected by MPB, respectively. WSBW caused defoliation has also increased substantially (4-fold on USFS managed lands) with the possible exception of BLM managed lands. Root disease patches, assumed to be armillaria root disease, were noted in the Big Snowy Mountains. Armillaria root disease was identified in the North Moccasin Mountains, north of Lewistown. LPP dwarf mistletoe is present in this County. A 43-acre polygon of an unidentified disease was mapped on Forest Service administered lands at the top of Browns Gulch, approximately 8 miles east of the community of Garneill. Also noted were a number of patches of windthrow, ranging in size from 3 acres to 75 acres, and dispersed across BLM (85 acres), State (24 acres) and private lands (28 acres). These were mapped northeast of Lewistown, near the top of Ruby Gulch. All were likely due to one wind event. Reports have been received from Montana DNRC foresters of decline and mortality of mature spruce in the Lewistown area. This is apparently affecting primarily urban and (or) landscape trees. Symptoms include branch dieback, thinning crowns, defoliation and (or) dieback of upper crowns, and tree mortality. Examination of

branch and foliage samples revealed the presence of various insect and disease organisms, but all are known to be mainly secondary problems on trees undergoing some sort of abiotic stress. The problem is as yet undiagnosed, but investigations are examining several possibilities, one being that the underlying cause of damage is pesticide-related. One small 13-acre patch of trees damaged from flooding was noted on private land on the North Fork of McDonald Creek, just west of the confluence with Alkali Creek.

## Flathead County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                                | National Forest | Other Federal | Private | State   | Total     |
|--------------------------------|-----------------|---------------|---------|---------|-----------|
| <b>Forestland</b>              | 1,647,905       | 518,720       | 448,029 | 151,729 | 2,766,383 |
| <b>Avalanche</b>               | 17              | 19            | 0       | 44      | 80        |
| <b>DFB</b>                     | 3,764           | 2,874         | 41      | 53      | 6,732     |
| <b>ESB</b>                     | 18              | 4             | 0       | 2       | 24        |
| <b>FE</b>                      | 271             | 50            | 163     | 92      | 576       |
| <b>Flooding - High Water</b>   | 0               | 2             | 0       | 0       | 2         |
| <b>MPB-LPP</b>                 | 22,492          | 1,630         | 2,669   | 619     | 27,410    |
| <b>MPB-PP</b>                  | 51              | 18            | 176     | 136     | 381       |
| <b>MPB-High Elevation</b>      | 111             | 0             | 0       | 0       | 111       |
| <b>IPS</b>                     | 0               | 0             | 0       | 2       | 2         |
| <b>SAF Mortality</b>           | 10,578          | 711           | 82      | 906     | 12,277    |
| <b>Unidentified Defoliator</b> | 0               | 0             | 100     | 0       | 100       |
| <b>WPB</b>                     | 2               | 0             | 16      | 0       | 18        |
| <b>WSBW</b>                    | 199,108         | 989           | 51,793  | 31,041  | 282,931   |
| <b>WPBR</b>                    | 0               | 0             | 17      | 0       | 17        |
| <b>Windthrow</b>               | 106             | 0             | 0       | 0       | 106       |

WSBW-caused defoliation is still continuing in Glacier NP. Total number of acres defoliated within the County greatly increased in 2009. Except for MPB-caused pine mortality, bark beetle activity was fairly light. MPB activity increased around Hungry Horse Reservoir, especially on the southern end. Scattered groups of MPB-LPP were also found on the southern edge of Glacier Park. FINDITs summaries for MPB in LPP and PP in mixed conifer stands indicate 0-41% total pine mortality. DFB activity remained about the same in 2009, scattered throughout the County. Numerous groups of DFB-caused mortality were mapped in Glacier NP and on the Coram Experimental Forest. In the park, scattered groups of up to several hundred trees were killed by DFB near Logging, Quartz, Bowman and Kintla Lakes. White pine blister rust is common in both WWP and WBP. Root diseases are common in counties west of the Continental Divide. The more common ones known to occur in this County are: s-type annosus root disease, armillaria root disease, and brown cubical root and butt rot. The tree species most affected are DF and true firs. P-type root disease is known to occur in PP. DF dwarf mistletoe, LPP dwarf mistletoe, and WL dwarf mistletoe are present in this

County. Eighty acres of avalanche damage was mapped in the County across Glacier NP (19 acres), State (44 acres) and Forest Service (17 acres) lands. Windthrow was noted on 305 acres, largely on BLM managed lands.

## Gallatin County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State  | Total   |
|---------------------------|-----------------|---------------|---------|--------|---------|
| <b>Forestland</b>         | 562,322         | 87,663        | 183,846 | 0      | 833,831 |
| <b>Aspen Decline</b>      | 0               | 0             | 72      | 0      | 72      |
| <b>DFB</b>                | 979             | 371           | 412     | 80     | 1,842   |
| <b>MPB-LPP</b>            | 157,201         | 11,223        | 71,553  | 10,280 | 250,257 |
| <b>MPB-High Elevation</b> | 67,410          | 2,155         | 9,143   | 166    | 78,874  |
| <b>SAF Mortality</b>      | 13,500          | 48            | 3,984   | 0      | 17,532  |
| <b>WSBW</b>               | 145,533         | 441           | 106,530 | 8,772  | 261,276 |

The most dramatic increase in insect activity on federal lands was in the number of acres defoliated by WSBW; which increased nearly 7-fold across the County. This occurred despite slight decreases in surveyed area. High amounts of WSBW-caused defoliation could result in an increase of DFB activity in the future; currently DFB activity is low in the County. MPB-killed LPP increased nearly 4-fold and almost doubled in high elevation pines (mostly WBP) over the last year. Very large groups of MPB-killed LPP were mapped throughout the Bridger Mountains. South of Bozeman, in the Gallatin Canyon and tributaries of the Gallatin River, LPP and WBP killed by MPB were extensive. MPB continues to attack and kill ornamental pines within the city of Bozeman. FINDITs summaries show some areas decreasing in MPB activity, largely due to host depletion. DNRC personnel hung MPB-baited traps at the Montana State University Marsh Lab and Montana State Trust Lands in Bear Canyon to determine flight period. Peak collections were on 9/1/09 (2 traps), and 8/13/09 (154 traps), respectively. SAF mortality also increased in 2009. White pine blister rust is common in WBP and LP. Tomentosus root disease is known to be significant in some campgrounds in this County. Black stain root disease identified from DF, but is considered of minor importance. LPP dwarf mistletoe and LP dwarf mistletoe are present in the County.

## Glacier County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State | Total   |
|---------------------------|-----------------|---------------|---------|-------|---------|
| <b>Forestland</b>         | 29,112          | 221,590       | 157,535 | 0     | 408,237 |
| <b>DFB</b>                | 0               | 538           | 110     | 0     | 648     |
| <b>ESB</b>                | 0               | 2             | 0       | 0     | 2       |
| <b>FE</b>                 | 0               | 2,382         | 2       | 0     | 2,384   |
| <b>MPB-LPP</b>            | 123             | 3,105         | 136     | 0     | 3,364   |
| <b>MPB-PP</b>             | 0               | 4             | 0       | 0     | 4       |
| <b>MPB-High Elevation</b> | 0               | 6             | 0       | 0     | 6       |
| <b>SAF Mortality</b>      | 0               | 6,905         | 0       | 0     | 6,905   |
| <b>WSBW</b>               | 0               | 18,686        | 395     | 0     | 19,081  |
| <b>Windthrow</b>          | 0               | 2             | 0       | 0     | 2       |
| <b>Winter Injury</b>      | 0               | 27            | 0       | 0     | 27      |

In 2009, the Rocky Mountain Front of the Lewis and Clark NF was flown for the first time in years. The increase in surveyed acres accounts for the new detection of MPB in LPP on USFS managed lands. Tribal lands and Glacier NP, however, had fewer acres surveyed (e.g. only the western portion of the Blackfeet IR was flown). Nevertheless, activity by FE, WBBB, and WSBW greatly increased on both ownerships. WSBW activity was notably high on Glacier Park's east side—especially along Cut Bank Ridge and near both St. Mary and Sherbourne Lakes. Despite detection of similar acres of DFB and MPB activity (in LPP) in 2008 and 2009, estimated number of trees killed suggests a significant drop in overall tree mortality for both beetles on both the NP and IR. However, increasing DFB activity might be anticipated in these areas of high WSBW activity. White pine blister rust is common in WBP and LP. Armillaria root disease known to be a significant pathogen in DF in the western portion of the County on the Blackfeet IR. LPP dwarf mistletoe and WL dwarf mistletoe are present in this County. One small 2-acre patch of windthrow was noted in Glacier NP east of Waterton Lake and one mile south of the Canadian border. Four neighboring polygons of winter injured LP, totaling 27 acres, was mapped on the Blackfeet IR approximately 3 miles east of the community of Saint Mary, east of Glacier NP.

## Golden Valley County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State | Total  |
|---------------------------|-----------------|---------------|---------|-------|--------|
| <b>Forestland</b>         | 9,411           | 0             | 73,038  | 0     | 82,449 |
| <b>MPB-LPP</b>            | 475             | 0             | 6       | 0     | 481    |
| <b>MPB-PP</b>             | 1,305           | 827           | 758     | 117   | 3,007  |
| <b>MPB-High Elevation</b> | 6               | 0             | 0       | 0     | 6      |

A similar number of acres were surveyed on USFS and BLM managed lands in 2009 as in 2008, with MPB activity in LPP showing and estimated 7 to 12 -fold increase. MPB activity in PP was also noted, particularly on USFS lands. WSBW activity, however, appears to have lessened, with no acres meeting the threshold of defoliation noted.

## Granite County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State  | Total   |
|---------------------------|-----------------|---------------|---------|--------|---------|
| <b>Forestland</b>         | 597,814         | 30,748        | 176,283 | 38,303 | 843,147 |
| <b>DFB</b>                | 875             | 44            | 58      | 6      | 983     |
| <b>FE</b>                 | 41              | 0             | 0       | 0      | 41      |
| <b>MPB-LPP</b>            | 144,858         | 5,547         | 21,330  | 1,712  | 173,447 |
| <b>MPB-PP</b>             | 2,163           | 729           | 3,612   | 714    | 7,218   |
| <b>MPB-High Elevation</b> | 5,776           | 0             | 414     | 0      | 6,190   |
| <b>SAF Mortality</b>      | 1,982           | 0             | 5       | 0      | 1,987   |
| <b>WPB</b>                | 0               | 2             | 6       | 4      | 12      |
| <b>WSBW</b>               | 17,638          | 10,518        | 24,739  | 3,751  | 56,646  |

Despite slight decreases in the total numbers of acres surveyed in 2008, 2009 surveys noted a 5-fold increase (or greater) in MPB activity in LPP, PP, and five-needle pines. Large groups of MPB-killed LPP were mapped north and east of Georgetown Lake. Virtually all pine stands east of Philipsburg and north of Anaconda, and between Philipsburg and Deerlodge, have been infested at some level by MPB. Defoliation by WSBW was also noted on approximately 10 times the number of acres, over all ownerships. DFB activity showed only a slight increase in acres while SAF mortality was detected at lower levels. Root diseases are common in counties west of the Continental Divide. The more common ones known to occur in this County are: s-type annosus root disease, armillaria root disease, and brown cubical root and butt rot. The tree species most affected are DF and true firs. P-type root disease is known to occur in PP. Schweinitzii root and butt rot is common in DF, causing decay in the butt logs but not acting as an aggressive root pathogen. Elytroderma needle disease is a significant agent in PP in localized areas in this County. LPP dwarf mistletoe and WL dwarf mistletoe are present in this County.

## Hill County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                   | National Forest | Other Federal | Private | State | Total  |
|-------------------|-----------------|---------------|---------|-------|--------|
| <b>Forestland</b> | 0               | 0             | 40,725  | 0     | 40,725 |
| <b>DFB</b>        | 0               | 2             | 8       | 4     | 14     |
| <b>MPB-LPP</b>    | 0               | 2,930         | 780     | 17    | 3,727  |
| <b>MPB-PP</b>     | 0               | 37            | 28      | 4     | 69     |
| <b>WPB</b>        | 0               | 2             | 0       | 0     | 2      |
| <b>WSBW</b>       | 0               | 1,980         | 2,564   | 12    | 4,556  |

MPB-killed LPP was the most significant mortality agent in the County. Number of acres of MPB-killed LPP, especially on tribal lands, nearly doubled. Very few PP stands were affected by MPB. WSBW activity significantly increased on both tribal and private lands. Armillaria root disease is present in the southern portion of the County, on the Rocky Boys IR, but appears to be uncommon. LPP dwarf mistletoe is present in this County.

## Jefferson County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State | Total   |
|---------------------------|-----------------|---------------|---------|-------|---------|
| <b>Forestland</b>         | 434,471         | 56,442        | 118,934 | 8,099 | 617,947 |
| <b>DFB</b>                | 200             | 31            | 218     | 8     | 457     |
| <b>MPB-LPP</b>            | 273,453         | 4,323         | 21,240  | 105   | 299,121 |
| <b>MPB-PP</b>             | 18,022          | 7,968         | 36,344  | 1,432 | 63,766  |
| <b>MPB-High Elevation</b> | 7,608           | 577           | 675     | 69    | 8,929   |
| <b>WSBW</b>               | 25,790          | 6,527         | 21,791  | 742   | 54,850  |
| <b>Windthrow</b>          | 249             | 0             | 0       | 0     | 249     |

MPB was still the most significant mortality agent in the County this year. MPB-caused mortality increased in some areas and decreased in many others due to the lack of available host trees. Large groups of MPB-killed trees were recorded in both LPP and PP stands, from about Jefferson City west nearly to Castle Rock, and east to nearly Canyon Ferry Lake. There was a significant increase in MPB activity in PP stands especially near Clancy and Montana City. At highest elevations, MPB-killed WBP remains high. Number of acres defoliated by WSBW significantly increased in 2009. DFB-caused mortality remains about the same but may increase in response to an increase of acres repeatedly defoliated by WSBW. White pine blister rust has been found in LP. A large 250-acre patch of windthrow, adjacent to a 53-acre patch of windthrow in Powell County, was mapped on Forest Service administered lands approximately 10 miles due east of the community of Racetrack.

## Judith Basin County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State | Total   |
|---------------------------|-----------------|---------------|---------|-------|---------|
| <b>Forestland</b>         | 269,838         | 0             | 22,039  | 0     | 291,877 |
| <b>DFB</b>                | 46              | 0             | 2       | 0     | 48      |
| <b>ESB</b>                | 10              | 0             | 0       | 0     | 10      |
| <b>MPB-LPP</b>            | 30,604          | 195           | 1,186   | 28    | 32,013  |
| <b>MPB-PP</b>             | 4,684           | 167           | 2,059   | 315   | 7,225   |
| <b>MPB-High Elevation</b> | 14,449          | 17            | 563     | 30    | 15,059  |
| <b>SAF-Mortality</b>      | 1,265           | 0             | 16      | 0     | 1,281   |
| <b>WSBW</b>               | 87,462          | 2,789         | 17,713  | 1,354 | 109,318 |

MPB continues to be the most important pest. With over three times the number of acres surveyed this year, most of it under USFS management, increases in detected MPB activity would be expected. However, nearly seven times the level of MPB affected acres (all hosts) were actually detected, representative of the increasing MPB activity in this County. Increases were notable to the east of King's Hill. Areas of detected WSBW defoliation also increased more than 4 -fold. The lack of increased detection of both DFB activity and mortality of SAF (complex lead by WBBB) with the increased survey area would suggest that actual mortality may have decreased. White pine blister rust is common in LP. Tomentosus root disease is known to be significant in some campgrounds in this County. LPP dwarf mistletoe is present in this County.

## Lake County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                                 | National Forest | Other Federal | Private | State  | Total   |
|---------------------------------|-----------------|---------------|---------|--------|---------|
| <b>Forestland</b>               | 136,743         | 0             | 366,299 | 48,387 | 551,429 |
| <b>Avalanche</b>                | 58              | 0             | 0       | 0      | 58      |
| <b>DFB</b>                      | 130             | 222           | 38      | 30     | 420     |
| <b>ESB</b>                      | 0               | 0             | 4       | 2      | 6       |
| <b>FE</b>                       | 59              | 2             | 49      | 11     | 121     |
| <b>Flooding - High Water</b>    | 0               | 0             | 12      | 0      | 12      |
| <b>MPB-LPP</b>                  | 6,511           | 3,415         | 1,077   | 510    | 11,513  |
| <b>MPB-PP</b>                   | 177             | 1,295         | 421     | 196    | 2,089   |
| <b>MPB-High Elevation</b>       | 0               | 8             | 0       | 0      | 8       |
| <b>IPS</b>                      | 0               | 2             | 2       | 24     | 28      |
| <b>Rhabdocline pseudotsugae</b> | 0               | 335           | 171     | 0      | 506     |
| <b>SAF Mortality</b>            | 1,115           | 1,240         | 45      | 59     | 2,459   |
| <b>Unidentified Defoliator</b>  | 0               | 38            | 13      | 0      | 51      |
| <b>WPB</b>                      | 0               | 12            | 4       | 2      | 18      |
| <b>WSBW</b>                     | 11,950          | 11,185        | 7,189   | 2,959  | 33,283  |
| <b>Winter Injury</b>            | 0               | 13            | 0       | 0      | 13      |

In 2009, aerial survey was conducted on about twice as many acres on tribal lands versus in 2008. MPB activity increased in both LPP and PP. Number of acres of MPB-killed PP significantly increased in the County, especially on tribal lands. MPB-caused mortality was light and scattered on the Island Unit. SAF mortality and DFB-caused mortality was light and scattered throughout the County. In 2009, the MT DNRC continued to trap *Ips* species in management units of PP on Fish, Wildlife, and Parks' Wild Horse Island. A total of 73 traps were set and more than 400,000 *Ips* (all species combined) were trapped between June 15 and August 31. Peak catches were on July 20 and July 27. Over 500 acres of Rhabdocline needle cast was mapped on the Flathead IR (335 acres) and neighboring private land (171 acres). This large area is located on the western edge of the Salish Mountains, east of the Little Bitterroot River, and is adjacent to over 1500 acres of Rhabdocline needle cast in neighboring Sanders County. White pine blister rust is common in both WWP and WBP. Root diseases are common in counties west of the Continental Divide. The more common ones known to occur in this County are: s-type annosus root disease, armillaria root disease, and brown cubical root and butt rot. The tree species most affected are DF and true firs. P-type root disease is known to occur in PP. Elytroderma needle disease is a significant agent in PP in localized areas in this County. DF dwarf mistletoe, LPP dwarf mistletoe, and WL dwarf mistletoe are present in this County. A notable problem with crown dieback of PP has occurred for the last several years along the Rocky Point area of Flathead Lake. Work by the Montana DNRC has shown that, for the most part, the damage is apparently due to Diplodia shoot blight, a native disease of PP in the West. A report titled "Ponderosa Pine Crown Dieback along the Rocky Point Area of Flathead Lake, Montana" has been completed that details the damage, diagnosis, biology, and

recommended management for this disease, both for the Rocky Point area and any other locations where similar problems may arise. Contact Amy Gannon ([agannon@mt.gov](mailto:agannon@mt.gov)) of the Montana DNRC for a copy of this report. One 58-acre avalanche was mapped on Forest Service administered lands at the headwaters of Lost Creek, just south of Swan Lake. This is very near a 17-acre avalanche in Flathead County. Two small polygons of flood damage were noted on private lands. A 13-acre patch of snow damaged trees was mapped at the head of Sullivan Gulch, near the Flathead and Lake County line, southwest of Big Arm.

## Lewis and Clark County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                             | National Forest | Other Federal | Private | State  | Total     |
|-----------------------------|-----------------|---------------|---------|--------|-----------|
| <b>Forestland</b>           | 982,490         | 30,326        | 350,445 | 42,855 | 1,406,116 |
| <b>Avalanche</b>            | 11              | 0             | 0       | 0      | 11        |
| <b>DFB</b>                  | 643             | 2             | 1,383   | 280    | 2,308     |
| <b>MPB-LPP</b>              | 234,107         | 10,237        | 61,556  | 11,308 | 317,208   |
| <b>MPB-PP</b>               | 52,489          | 15,417        | 82,689  | 11,020 | 161,615   |
| <b>MPB (High Elevation)</b> | 3,035           | 14            | 5       | 57     | 3,111     |
| <b>SAF Mortality</b>        | 53              | 0             | 0       | 0      | 53        |
| <b>WSBW</b>                 | 145,970         | 11,551        | 77,018  | 11,396 | 245,935   |
| <b>Windthrow</b>            | 16              | 0             | 0       | 0      | 16        |

More acres of aerial survey were flown in 2009 on all lands. In 2009, total acres of MPB-killed pines once again increased significantly. In many areas with previous high infestations, intensity began to decline because the majority of the host trees had already been killed by MPB. WSBW and MPB activity is increasing near Rogers Pass. Number of WSBW-defoliated acres greatly increased in the County. Some of the acres defoliated also had increasing levels of DFB activity, such as near Flesher Pass. Many large groups of MPB-killed LPP were once again mapped in nearly every direction from Lincoln. Other bark beetle activity was virtually at endemic levels, except DFB. DNRC personnel hung MPB-baited traps at Mount Helena and the MT DNRC Office to determine flight period. Peak collections were on 8/5/09 (212) and 8/12/09 (216), respectively. White pine blister rust is common in WBP and LP. Armillaria root disease is present in the southeastern portion of the County, and Schweinitzii root and butt rot is quite common, especially causing significant decay in the butt logs of larger, older DF. LPP dwarf mistletoe is present in this County. One 11-acre patch of avalanche damage was noted on Forest Service administered lands just east of the Continental Divide, at the headwaters for Little Prickly Pear Creek, north of Black Mountain. A 16-acre patch of windthrow was mapped on Forest Service administered lands in the headwaters of Deadman Creek, just west of Stemple Pass on the Continental Divide.

## Lincoln County

### Acres of Forestland, Mortality, and Defoliation by Ownership

|                              | National Forest | Other Federal | Private | State  | Total     |
|------------------------------|-----------------|---------------|---------|--------|-----------|
| <b>Forestland</b>            | 1,762,707       | 0             | 358,595 | 39,206 | 2,160,508 |
| <b>Diplodia Canker</b>       | 39              | 0             | 0       | 0      | 39        |
| <b>DFB</b>                   | 312             | 0             | 49      | 6      | 367       |
| <b>ESB</b>                   | 8               | 0             | 0       | 4      | 12        |
| <b>FE</b>                    | 157             | 0             | 150     | 11     | 318       |
| <b>Flooding - High Water</b> | 2               | 0             | 0       | 0      | 2         |
| <b>MPB-LPP</b>               | 4,528           | 0             | 979     | 20     | 5,527     |
| <b>MPB-PP</b>                | 633             | 51            | 86      | 14     | 784       |
| <b>MBP-WP</b>                | 16              | 0             | 2       | 0      | 18        |
| <b>MPB-High Elevation</b>    | 10              | 0             | 0       | 0      | 10        |
| <b>IPS</b>                   | 2               | 2             | 2       | 0      | 6         |
| <b>SAF Mortality</b>         | 6,186           | 0             | 0       | 112    | 6,298     |
| <b>WPB</b>                   | 74              | 0             | 52      | 2      | 128       |
| <b>WSBW</b>                  | 149,938         | 0             | 27,794  | 3,120  | 180,852   |
| <b>WPBR</b>                  | 963             | 0             | 57      | 0      | 1,020     |
| <b>Windthrow</b>             | 2               | 0             | 0       | 2      | 4         |

Aerial survey was conducted in most of the County in 2009. MPB-killed LPP and PP increased across the County; however, Lincoln County is still one of the least impacted in the State. MPB-killed LPP was light and scattered throughout the Yaak River drainage, and east and south of Libby. Endemic levels of WPB and IPS were found scattered throughout the County. Defoliation from WSBW was recorded at high levels throughout the County, especially west of Highway 2, south of Libby. Two male gypsy moths were caught in Lincoln County. Molecular analysis confirmed both were European. One 39-acre patch of PP infested with Diplodia twig blight was mapped on Forest Service administered lands on the north side of Seventeenmile Creek of the Yaak River. Aerial Detection Survey is not a reliable method for recording the presence of Diplodia twig blight, but the detection from ADS is worth noting in this report. Just over 1,000 acres of white pine blister rust was mapped on Forest Service (963 acres) and private lands (57 acres) in the County. Most polygons were mapped in west and southern portions of the County. Aerial Detection Survey is not a reliable method for recording the presence of white pine blister rust, but the detection from ADS is worth noting in this report. White pine blister rust is common in both WWP and WBP. Root diseases are common in counties west of the Continental Divide. The more common ones known to occur in this County are: s-type annosus root disease, armillaria root disease, and brown cubical root and butt rot. The tree species most affected are DF and true firs. P-type root disease is known to occur in PP. Armillaria root disease was found affecting large PP near Rocky Gorge Campground on Lake Koocanusa. It is possible annosus root disease may have also been involved in the decline of these trees, but was not positively identified (MFO-TR-09-44). DF dwarf mistletoe, LPP dwarf mistletoe, and WL dwarf mistletoe are present in this County. A small 2-acre patch of flood

damage on Forest Service administered lands was noted. Two small, 2-acre polygons of windthrow were observed on Forest Service and State lands.

## Madison County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State | Total   |
|---------------------------|-----------------|---------------|---------|-------|---------|
| <b>Forestland</b>         | 602,613         | 120,988       | 112,579 | 0     | 836,180 |
| <b>Aspen Decline</b>      | 13              | 0             | 26      | 0     | 39      |
| <b>DFB</b>                | 14              | 6             | 4       | 0     | 24      |
| <b>MPB-LPP</b>            | 107,474         | 8,295         | 23,803  | 1,210 | 140,782 |
| <b>MPB-PP</b>             | 92              | 0             | 2       | 0     | 94      |
| <b>MPB-High Elevation</b> | 55,400          | 1,978         | 5,627   | 308   | 63,313  |
| <b>SAF Mortality</b>      | 5,609           | 6             | 720     | 2     | 6,337   |
| <b>WSBW</b>               | 44,677          | 18,253        | 25,232  | 2,581 | 90,743  |

MPB has been active in Madison County for several years, with activity continuing to show some increases in LPP but, overall, beginning to level off as host is depleted. Mortality of WBP continues throughout high elevations of both the Tobacco Root and Snowcrest Ranges. Past WBP mortality in the Gravelly Range has left few remaining large WBP, many of which are dying in small scattered spots. Mortality of LPP continues along the eastern fronts of the Tobacco and Gravelly Ranges, and in the Snowcrest Range where LPP are found. Both DFB activity and SAF mortality remain similar to 2008 levels, approximating endemic conditions. The few small polygons of DFB-caused DF mortality were noted along the Warm Springs drainage of the Gravelly Range and the headwaters of the Ruby River. Area affected by WSBW defoliation showed modest increases over all ownerships at lower elevations around all three mountain ranges. If defoliation continues at high levels or drought conditions return, increases in DFB activity may be expected. White pine blister rust is common in WBP and LP. Schweinitzii root and butt rot is common in DF, causing decay in the butt logs but not acting as an aggressive root pathogen. LP dwarf mistletoe is present in this County.

## Meagher County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State  | Total   |
|---------------------------|-----------------|---------------|---------|--------|---------|
| <b>Forestland</b>         | 452,100         | 0             | 221,586 | 17,765 | 691,450 |
| <b>DFB</b>                | 10              | 0             | 20      | 4      | 34      |
| <b>MPB-LPP</b>            | 221,893         | 413           | 28,501  | 1,037  | 251,844 |
| <b>MPB-PP</b>             | 7,690           | 986           | 20,998  | 892    | 30,566  |
| <b>MPB-High Elevation</b> | 18,824          | 39            | 651     | 0      | 19,514  |
| <b>SAF Mortality</b>      | 54              | 0             | 0       | 0      | 54      |
| <b>WSBW</b>               | 165,095         | 4,282         | 150,491 | 5,455  | 325,323 |
| <b>Windthrow</b>          | 40              | 0             | 0       | 0      | 40      |

In 2009, most of the County was flown, which partly explains significant increase in number of acres reported to be killed by bark beetles. Despite this, number of acres killed by MPB increased substantially throughout the Castle Mountains. MPB activity in LPP increased by nearly 2-fold and by nearly 10-fold in PP. WSBW activity significantly increased throughout the County on private and Forest Service administered lands.

Although DFB activity decreased, increases in the future can be anticipated in stands heavily defoliated by WSBW. White pine blister rust has been found in LP. Armillaria root disease is present in this County. Schweinitzii root and butt rot is quite common, especially causing significant decay in the butt logs of larger, older DF. LPP dwarf mistletoe is present in this County. One 40-acre patch of windthrow on Forest Service administered lands was noted in Moonshine Gulch of Antelope Creek.

## Mineral County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State  | Total   |
|---------------------------|-----------------|---------------|---------|--------|---------|
| <b>Forestland</b>         | 632,710         | 0             | 85,080  | 21,305 | 739,095 |
| <b>DFB</b>                | 201             | 0             | 14      | 0      | 215     |
| <b>ESB</b>                | 2               | 0             | 0       | 0      | 2       |
| <b>FE</b>                 | 589             | 0             | 53      | 2      | 644     |
| <b>MPB-LPP</b>            | 106,969         | 0             | 1,916   | 60     | 108,945 |
| <b>MPB-PP</b>             | 3,574           | 0             | 180     | 27     | 3,781   |
| <b>MPB-WP</b>             | 4               | 0             | 0       | 0      | 4       |
| <b>MPB-High Elevation</b> | 4               | 0             | 0       | 0      | 4       |
| <b>SAF Mortality</b>      | 468             | 0             | 6       | 0      | 474     |
| <b>WPB</b>                | 373             | 0             | 15      | 20     | 408     |
| <b>WPBR</b>               | 25              | 0             | 4       | 0      | 29      |

There was an increase of about 20% in the number of acres flown in the County. MPB-killed LPP was by far the most significant mortality agent on lands surveyed, especially south of I-90 from just above Trout Creek west to Lookout Pass. Fader groups just east of Lookout Pass were large and intensely infested. Most intensely infested stands were near Lookout Pass and in the Dry and Cedar Creek drainages. North of I-90 and the Clark Fork River, beetle activity was much less frequently encountered. There was also a modest increase from 172 to 378 acres of lower-elevation PP killed by MPB in 2009. Minor amounts of WPB-killed PP and SAF mortality were also recorded in the County. White pine blister rust is common in both WWP and WBP. Root diseases are common in counties west of the Continental Divide. The more common ones known to occur in this County are: s-type annosus root disease, armillaria root disease, laminated root disease, and brown cubical root and butt rot. The tree species most affected are DF and true firs. P-type root disease is known to occur in PP. DF dwarf mistletoe, LPP dwarf mistletoe, and WL dwarf mistletoe are present in this County.

## Missoula County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                                | National Forest | Other Federal | Private | State   | Total     |
|--------------------------------|-----------------|---------------|---------|---------|-----------|
| <b>Forestland</b>              | 641,553         | 10,194        | 600,301 | 144,223 | 1,396,272 |
| <b>DFB</b>                     | 1,084           | 4             | 111     | 10      | 1,209     |
| <b>Drought</b>                 | 0               | 0             | 2       | 0       | 2         |
| <b>ESB</b>                     | 0               | 0             | 2       | 0       | 2         |
| <b>FE</b>                      | 7               | 0             | 29      | 0       | 36        |
| <b>Flooding - High Water</b>   | 0               | 2             | 0       | 0       | 2         |
| <b>MPB-LPP</b>                 | 87,299          | 14,651        | 23,591  | 11,475  | 137,016   |
| <b>MPB-PP</b>                  | 6,335           | 1,164         | 18,195  | 9,259   | 34,953    |
| <b>MPB-High Elevation</b>      | 4               | 149           | 0       | 0       | 153       |
| <b>IPS</b>                     | 0               | 0             | 2       | 0       | 2         |
| <b>SAF Mortality</b>           | 149             | 683           | 11      | 66      | 909       |
| <b>Unidentified Defoliator</b> | 0               | 177           | 0       | 0       | 177       |
| <b>WPB</b>                     | 6               | 2             | 17      | 18      | 43        |
| <b>WSB</b>                     | 1,235           | 4,484         | 6,459   | 4,966   | 17,144    |

MPB activity significantly increased around Missoula, Lolo and the Ninemile Creek drainage. The most significant increase was in the acres of MPB-killed PP followed by MPB-killed LPP. Large groups of MPB-killed LPP were mapped east of the Rattlesnake Wilderness, especially towards Sheep Mountain and Hidden Lake. MPB-killed LPP was also common north and east of Missoula. MPB-killed PP stands were extensively found throughout the Rattlesnake Creek drainage, from Mount Jumbo, north into the Wilderness and northwestward to Grant Creek. Other PP stands were infested to a lesser extent in the Butler Creek drainage. Acres of MPB-killed PP greatly increased on private lands in 2009. DNRC personnel hung MPB-baited traps at Condon, Clearwater Junction, Boy Scout Road, and the MT DNRC Office to determine flight period. Peak

collections were on 9/17/09 (3), 8/14/09 (24), 8/26/09 (266), and 7/15/09 (33), respectively. DFB activity remained about the same in 2009, with 1,848 acres affected across the County. There were minor amounts of SAF mortality recorded in the Fish Creek drainage. WSBW defoliated stands were also detected, especially in the eastern part of the County. WBP and IPS activity slightly increased and was scattered. White pine blister rust is common in both WWP and WBP. Root diseases are common in counties west of the Continental Divide. The more common ones known to occur in this County are: s-type annosus root disease, armillaria root disease, laminated root disease, and brown cubical root and butt rot. The tree species most affected are DF and true firs. P-type root disease is known to occur in PP. An unidentified species of black stain root disease was noted in LPP in the Hemlock-Elk Analysis Area on Forest Service administered lands west of Condon (MFO-TR-09-11). Elytroderma needle disease is a significant agent in PP in localized areas in this County. DF dwarf mistletoe, LPP dwarf mistletoe, and WL dwarf mistletoe are present in this County. One 2-acre polygon of drought damage was mapped on private lands in the Garnet Range, near the headwaters of Ashby Creek.

## Musselshell County

Acres of Forestland, Mortality, and Defoliation by Ownership

|            | National Forest | Other Federal | Private | State  | Total   |
|------------|-----------------|---------------|---------|--------|---------|
| Forestland | 0               | 28,858        | 276,685 | 15,962 | 321,505 |
| MPB-PP     | 0               | 0             | 4       | 1      | 5       |

Only 5 acres of MPB in PP were detected along the County border, with one acre being on federally managed lands (Fish and Wildlife Service) and four acres on privately managed lands. At least 240 acres of PTM activity were observed by DNRC field staff. Defoliation was severe throughout the PP stand.

## Park County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                       | National Forest | Other Federal | Private | State  | Total     |
|-----------------------|-----------------|---------------|---------|--------|-----------|
| Forestland            | 751,119         | 81,903        | 171,212 | 31,636 | 1,035,870 |
| Aspen Decline         | 0               | 71            | 0       | 0      | 71        |
| DFB                   | 446             | 32            | 146     | 0      | 624       |
| ESB                   | 0               | 2             | 0       | 0      | 2         |
| Flooding - High Water | 0               | 180           | 0       | 0      | 180       |
| MPB-LPP               | 53,801          | 35,501        | 48,911  | 108    | 138,321   |
| MPB-High Elevation    | 32,059          | 36,640        | 14,219  | 0      | 82,918    |
| SAF Mortality         | 6,004           | 302           | 2,432   | 0      | 8,738     |
| WSBW                  | 32,850          | 35,807        | 47,591  | 459    | 116,707   |

MPB-killed LPP increased almost 4-fold in the County in 2009. Large polygons of MPB-killed LPP were found in the Gallatin, Absaroka and Crazy Mountain ranges. MPB activity in five-needle pine almost doubled. SAF mortality increased significantly from 1,591 to 8,444 acres. WSBW-caused defoliation almost doubled with most of the increase on private lands. White pine blister rust is common in WBP and LP. Tomentosus root disease is known to be significant in some campgrounds in this County. LPP dwarf mistletoe is present in this County.

## Petroleum County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                   | National Forest | Other Federal | Private | State  | Total   |
|-------------------|-----------------|---------------|---------|--------|---------|
| <b>Forestland</b> | 0               | 69,956        | 12,613  | 24,076 | 106,644 |
| <b>MPB-PP</b>     | 0               | 2             | 0       | 0      | 2       |

Two acres of MPB-killed PP were detected.

## Phillips County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                   | National Forest | Other Federal | Private | State | Total   |
|-------------------|-----------------|---------------|---------|-------|---------|
| <b>Forestland</b> | 0               | 74,920        | 58,478  | 0     | 133,397 |
| <b>DFB</b>        | 0               | 6             | 0       | 0     | 6       |
| <b>MPB-LPP</b>    | 0               | 73            | 0       | 0     | 73      |
| <b>MPB-PP</b>     | 0               | 417           | 19      | 2     | 438     |
| <b>WSBW</b>       | 0               | 43            | 0       | 0     | 43      |

Fewer acres were flown in the County in 2009. Scattered groups of LPP and PP mortality were detected throughout the County, including Fort Belknap IR. Defoliation from WSBW was recorded on 560 acres. LPP dwarf mistletoe is present in this County.

## Pondera County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State  | Total   |
|---------------------------|-----------------|---------------|---------|--------|---------|
| <b>Forestland</b>         | 93,289          | 0             | 0       | 11,661 | 104,950 |
| <b>DFB</b>                | 13              | 0             | 6       | 0      | 19      |
| <b>MPB-LPP</b>            | 2,304           | 27            | 21      | 242    | 2,594   |
| <b>MPB-High Elevation</b> | 58              | 0             | 4       | 0      | 62      |
| <b>SAF Mortality</b>      | 2,030           | 0             | 0       | 0      | 2,030   |
| <b>WSBW</b>               | 0               | 308           | 1,569   | 455    | 2,332   |
| <b>Windthrow</b>          | 75              | 0             | 0       | 0      | 75      |

In 2009 the Rocky Mountain Front of the Lewis & Clark was surveyed for the first time in many years. Increased pest activity is due to this nearly 40 -fold increase in survey area. On Forest Service administered lands, mortality of LPP due to MPB and SAF due to a complex of agents were the most important pest agents. On other ownerships, WSBW was the most prevalent agent. White pine blister rust is common in WBP and LP. A 72-acre patch of windthrow was mapped on Forest Service administered lands at the headwaters of the South Fork of Two Medicine River, southeast of East Glacier Park.

## Powder River County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                   | National Forest | Other Federal | Private | State  | Total   |
|-------------------|-----------------|---------------|---------|--------|---------|
| <b>Forestland</b> | 253,254         | 71,096        | 156,632 | 10,052 | 491,033 |
| <b>MPB-PP</b>     | 99              | 0             | 8       | 2      | 109     |
| <b>WSBW</b>       | 137             | 0             | 167     | 0      | 304     |

Beetle activity was recorded at endemic levels with only a minor amount of MPB-killed PP mapped in forests, mostly to the east and south of Ashland. No IPS activity was noted, although it is likely that MPB and IPS are acting together in current, endemic pockets of PP mortality. The PTM activity (642 acres) noted in 2008 has subsided.

## Powell County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State  | Total   |
|---------------------------|-----------------|---------------|---------|--------|---------|
| <b>Forestland</b>         | 629,866         | 69,322        | 242,905 | 13,704 | 955,797 |
| <b>DFB</b>                | 183             | 27            | 58      | 8      | 276     |
| <b>MPB-LPP</b>            | 183,149         | 51,805        | 45,667  | 7,642  | 288,263 |
| <b>MPB-PP</b>             | 5,808           | 2,958         | 24,921  | 2,936  | 36,623  |
| <b>MPB-High Elevation</b> | 2,346           | 0             | 13      | 0      | 2,359   |
| <b>IPS</b>                | 0               | 0             | 135     | 0      | 135     |
| <b>SAF Mortality</b>      | 102             | 0             | 0       | 0      | 102     |
| <b>WPB</b>                | 0               | 0             | 20      | 6      | 26      |
| <b>WSBW</b>               | 55,046          | 41,451        | 114,034 | 13,286 | 223,817 |
| <b>Windthrow</b>          | 52              | 0             | 0       | 0      | 52      |

WSBW caused defoliation significantly increased across many parts of the County. MPB activity increased in both LPP and PP, on private and federal lands. Around MacDonald pass, mortality has peaked and is declining. On Forest Service administered lands, large polygons of MPB-killed LPP were detected near Ovando Mountain. Endemic levels of DFB, WPB and IPS were lightly scattered across the County.

White pine blister rust is common in WBP and LP. Root diseases are common in counties west of the Continental Divide. The more common ones known to occur in this County are: s-type annosus root disease, armillaria root disease, and brown cubical root and butt rot. The tree species most affected are DF and true firs. P-type root disease is known to occur in PP. LPP dwarf mistletoe is present in this County. One 52-acre patch of windthrow was mapped very near a 250-acre patch of windthrow mapped on Forest Service administered lands in Jefferson County. Both patches are likely due to the same wind event, and are located approximately 10 miles east of the community of Racetrack.

## Ravalli County

### Acres of Forestland, Mortality, and Defoliation by Ownership

|                                   | National Forest | Other Federal | Private | State  | Total     |
|-----------------------------------|-----------------|---------------|---------|--------|-----------|
| <b>Forestland</b>                 | 1,062,377       | 0             | 139,434 | 29,246 | 1,231,057 |
| <b>DFB</b>                        | 219             | 0             | 6       | 0      | 225       |
| <b>Dwarf Mistletoe</b>            | 0               | 0             | 2       | 0      | 2         |
| <b>Elytroderma</b>                | 21              | 0             | 1       | 0      | 22        |
| <b>FE</b>                         | 18              | 0             | 0       | 0      | 18        |
| <b>MPB-LPP</b>                    | 12,204          | 0             | 308     | 66     | 12,578    |
| <b>MPB-PP</b>                     | 698             | 0             | 355     | 437    | 1,490     |
| <b>MPB-High Elevation</b>         | 2,959           | 0             | 0       | 0      | 2,959     |
| <b>Pine Engraver Beetle (LPP)</b> | 7               | 0             | 0       | 0      | 7         |
| <b>IPS</b>                        | 0               | 0             | 2       | 2      | 4         |
| <b>SAF Mortality</b>              | 5,546           | 0             | 43      | 0      | 5,589     |
| <b>WSBW</b>                       | 1,329           | 0             | 61      | 444    | 1,834     |

In 2008, MPB activity was the most common bark beetle-caused mortality in the County, and that was also the case in 2009. Activity in LPP appears to be leveling off with the 2009 survey detecting more acres affected but fewer trees killed. Large groups of MPB-killed LPP were mapped in the Sapphire Mountains east of Stevensville, areas east of Hamilton, and around Painted Rocks Reservoir. MPB activity (acres and numbers) increased in both PP and five-needle pines. Several patches of PP mortality were noted east of Darby. Other bark beetle-caused mortality was recorded at endemic levels. Modest increases in WSBW-caused defoliation were detected. It should be noted that despite nearly double the acres of SAF mortality mapped in 2009, only around half the number of trees were estimated to have been killed.

Small patches of DF dwarf mistletoe and Elytroderma needle disease were mapped in County. Aerial Detection Survey is not a reliable method for recording the presence of these two diseases. Both diseases are quite common in the forests of Ravalli County. White pine blister rust is common in WBP. Root diseases are common in counties west of the Continental Divide. The more common ones known to occur in this County are: s-type annosus root disease, armillaria root disease, laminated root disease, and brown cubical root and butt rot. The tree species most affected are DF and true firs. P-type root disease is known to occur in PP. A yet-to-be-identified root disease was noted in young DF in a 20- to 25-year old plantation on Forest Service administered lands east of Stevensville near Ambrose Saddle. The decay was noted in several trees across the plantation (MFO-TR-09-35). Identification of the responsible agent is forthcoming. Elytroderma needle disease is a significant agent in PP in localized areas in this County. DF dwarf mistletoe, LPP dwarf mistletoe, and WL dwarf mistletoe are present in this County.

## Rosebud County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State  | Total   |
|---------------------------|-----------------|---------------|---------|--------|---------|
| <b>Forestland</b>         | 76,525          | 38,671        | 223,688 | 32,981 | 371,865 |
| <b>MPB-PP</b>             | 20              | 61            | 2       | 0      | 83      |
| <b>MPB-High Elevation</b> | 0               | 2             | 0       | 0      | 2       |
| <b>WSBW</b>               | 0               | 491           | 0       | 0      | 491     |

Pest activity remains at low levels, with limited WSBW and minimal MPB (likely in combination with IPS) noted on the Northern Cheyenne IR.

## Sanders County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                                 | National Forest | Other Federal | Private | State  | Total     |
|---------------------------------|-----------------|---------------|---------|--------|-----------|
| <b>Forestland</b>               | 880,116         | 11,339        | 477,381 | 61,432 | 1,430,268 |
| <b>DFB</b>                      | 207             | 0             | 46      | 0      | 253       |
| <b>FE</b>                       | 1,891           | 2             | 2,894   | 261    | 5,048     |
| <b>Larch Needle Blight</b>      | 60              | 0             | 0       | 0      | 60        |
| <b>MPB-LPP</b>                  | 98,114          | 462           | 4,947   | 1,547  | 105,070   |
| <b>MPB-PP</b>                   | 494             | 314           | 163     | 38     | 1,009     |
| <b>MPB-WP</b>                   | 2               | 0             | 2       | 0      | 4         |
| <b>IPS</b>                      | 616             | 2             | 10      | 0      | 628       |
| <b>Rhabdocline pseudotsugae</b> | 0               | 997           | 557     | 0      | 1,554     |
| <b>SAF Mortality</b>            | 209             | 20            | 54      | 0      | 283       |
| <b>WPB</b>                      | 102             | 4             | 47      | 8      | 161       |
| <b>WSBW</b>                     | 158,932         | 0             | 15,323  | 2,173  | 176,428   |
| <b>WPBR</b>                     | 34              | 0             | 3       | 0      | 37        |

Many groups of MPB-killed LPP were mapped surrounding Plains and Thompson Falls. Fader groups increased significantly throughout the Prospect Creek drainage from Thompson Pass to Thompson Falls. South of Plains and south of the Clark Fork River, small groups of MPB-killed LPP, WPB-killed PP, and FE-impacted GF were found widely scattered. Elsewhere in the County, bark beetle activity was light and scattered. WSBW activity significantly increased in some parts of the County and could lead to increases in DFB activity in the future.

One 60-acre polygon of severe WL needle blight was mapped on Forest Service administered lands in Pew Gulch, just west of Noxon. Over 1500 acres of Rhabdocline needle cast was mapped on the Flathead IR (997 acres) and neighboring private land (557 acres). This large area is located on the western edge of the Salish Mountains, east of the Little Bitterroot River. White pine blister rust is common in both WWP and WBP. Root diseases are common in counties west of the Continental Divide. The more common ones known to occur in this County are: s-type annosus root disease,

armillaria root disease, laminated root disease, and brown cubical root and butt rot. The tree species most affected are DF and true firs. P-type root disease is known to occur in PP. Severe Armillaria root disease was found affecting large PP in the Pilgrim Creek drainage (MFO-TR-09-21). Tomentosus root disease significantly impacted LP in a campground near Trout Creek (MFO-TR-09-09). Elytroderma needle disease is a significant agent in PP in localized areas in this County. DF dwarf mistletoe, LPP dwarf mistletoe, and WL dwarf mistletoe are present in this County.

## Silver Bow County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State | Total   |
|---------------------------|-----------------|---------------|---------|-------|---------|
| <b>Forestland</b>         | 178,528         | 24,440        | 39,153  | 4,067 | 246,189 |
| <b>Aspen Decline</b>      | 0               | 0             | 56      | 0     | 56      |
| <b>DFB</b>                | 13              | 8             | 10      | 6     | 37      |
| <b>MPB-LPP</b>            | 102,577         | 5,712         | 7,774   | 4,191 | 120,254 |
| <b>MPB-High Elevation</b> | 11,528          | 2             | 36      | 212   | 11,778  |
| <b>WSBW</b>               | 23,687          | 11,070        | 17,552  | 2,666 | 54,975  |

Over 80% of the County was surveyed in both 2008 and 2009. MPB continues to kill large numbers of trees over large areas; nearly doubling activity in high elevations (LPP and five-needle pines). However, mortality in LPP (trees per acre and numbers of trees) is down, due largely to depletion of adequate host. WSBW-caused defoliation has increased over two-fold. Although DFB activity remains at low, endemic levels, increased defoliation may cause greater DFB activity in the future. White pine blister rust is common in WBP and LP.

## Stillwater County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                                | National Forest | Other Federal | Private | State | Total   |
|--------------------------------|-----------------|---------------|---------|-------|---------|
| <b>Forestland</b>              | 122,032         | 0             | 114,463 | 0     | 236,496 |
| <b>DFB</b>                     | 22              | 0             | 2       | 0     | 24      |
| <b>MPB-LPP</b>                 | 301             | 0             | 4       | 0     | 305     |
| <b>MPB-PP</b>                  | 0               | 0             | 2       | 0     | 2       |
| <b>MPB-High Elevation</b>      | 18              | 0             | 2       | 0     | 20      |
| <b>SAF Mortality</b>           | 12              | 0             | 0       | 0     | 12      |
| <b>Unidentified Defoliator</b> | 11              | 0             | 0       | 0     | 11      |
| <b>WSBW</b>                    | 6               | 0             | 24      | 0     | 30      |

MPB-caused mortality was observed in LPP and WBP stands administered by the Custer NF and in the Beartooth Mountains. Several groups of MPB-killed LPP were

noted near Washbourn Mountains, west of Absarokee. MPB caused mortality in PP slightly declined in the County and SAF mortality also declined. Minor amounts of DFB-caused mortality were also noted. Damage caused by PTM was not mapped by ADS and landowners confirm that populations appear to be in decline. White pine blister rust has been found in WBP and LP.

## Sweet Grass County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State | Total   |
|---------------------------|-----------------|---------------|---------|-------|---------|
| <b>Forestland</b>         | 243,865         | 16,803        | 117,302 | 0     | 377,970 |
| <b>DFB</b>                | 96              | 0             | 2       | 0     | 98      |
| <b>MPB-LPP</b>            | 11,712          | 284           | 3,292   | 0     | 15,288  |
| <b>MPB-High Elevation</b> | 11,714          | 0             | 470     | 0     | 12,184  |
| <b>SAF Mortality</b>      | 6,095           | 0             | 1       | 0     | 6,096   |
| <b>WSBW</b>               | 3,826           | 138           | 2,704   | 53    | 6,721   |

DFB activity was low and scattered in 2009. There was a 3-fold increase in MPB-killed LPP and five-needle pines mostly in the Beartooth and eastern Crazy Mountains and in the Boulder Range. Number of acres of WSBW defoliation almost doubled in 2009. White pine blister rust has been found in WBP and LP.

## Teton County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                              | National Forest | Other Federal | Private | State  | Total   |
|------------------------------|-----------------|---------------|---------|--------|---------|
| <b>Forestland</b>            | 216,012         | 10,294        | 20,237  | 10,294 | 256,837 |
| <b>DFB</b>                   | 2,260           | 112           | 60      | 43     | 2,475   |
| <b>ESB</b>                   |                 | 16            |         |        | 16      |
| <b>Flooding - High Water</b> |                 | 67            |         |        | 67      |
| <b>MPB-LPP</b>               | 9,845           | 13,987        | 1,812   | 1,322  | 26,966  |
| <b>MPB-High Elevation</b>    | 2,388           | 2,477         | 32      | 5      | 4,902   |
| <b>SAF Mortality</b>         | 102             | 1,971         | 0       | 0      | 2,073   |
| <b>WSBW</b>                  | 191             | 3,652         | 3,061   | 1,198  | 8,102   |
| <b>Windthrow</b>             |                 | 71            |         |        | 71      |

Increased efforts to survey Lewis and Clark NF in 2009 provided some of the first aerial pest survey data for the Rocky Mountain Front, in many years. MPB in LPP was the greatest pest problem noted on all ownerships, although MPB in five-needle pines and DFB in DF was both notable on Forest Service administered lands. DFB was particularly notable near edges of the 2007 fires, due largely to DFB's affinity for scorched DF where populations can build to outbreak numbers. WBBB affected SAF

were also mapped over 70 acres. WSBW caused defoliation was also very active on both private and state lands. Increases in defoliation, coupled with locally high DFB populations may result in continued loss of DF. White pine blister rust is common in WBP and LP.

## Valley County

Although no ADS were conducted in Valley County, FHP personnel did visit Fort Peck Corps of Engineer managed lands to evaluate reported tree mortality. Ground surveys found scattered small pockets of PP being attacked and killed by secondary bark beetles, particularly *Ips* species.

## Wheatland County

Acres of Forestland, Mortality, and Defoliation by Ownership

|                           | National Forest | Other Federal | Private | State | Total  |
|---------------------------|-----------------|---------------|---------|-------|--------|
| <b>Forestland</b>         | 41,083          | 0             | 25,677  | 0     | 66,760 |
| <b>ESB</b>                | 2               | 0             | 0       | 0     | 2      |
| <b>MPB-LPP</b>            | 12,842          | 0             | 591     | 0     | 13,433 |
| <b>MPB-PP</b>             | 1,656           | 15            | 4,050   | 198   | 5,919  |
| <b>MPB-High Elevation</b> | 481             | 0             | 321     | 0     | 802    |
| <b>WSBW</b>               | 36,198          | 64            | 2,655   | 125   | 39,042 |

A 15 -fold increase in surveyed area, largely on Forest Service administered lands, explains much of the increased pest activity noted in 2009. In pines, MPB remains the most significant pest; killing significant numbers of LPP, PP, and five-needle pines (includes LP). Over 25% of the surveyed lands showed signs of defoliation by WSBW. WSBW defoliates principally DF, but also affecting ES and SAF. White pine blister rust has been found in LP.

**Table 1. Mortality, Defoliation and Other Damage detected from the air on National Forests, National Parks and Tribal Lands in Montana, during 2009.**

| <b>BEAVERHEAD NF</b>           |              |              |                                |              |              |
|--------------------------------|--------------|--------------|--------------------------------|--------------|--------------|
| <b>Dillon RD</b>               | <b>Acres</b> | <b>Trees</b> | MPB (High Elev 5-needle Pines) | 925          | 1,545        |
| Douglas-fir Beetle             | 14           | 42           | <b>Stevensville RD</b>         | <b>Acres</b> | <b>Trees</b> |
| Mountain Pine Beetle (LPP)     | 68,975       | 228,697      | Douglas-fir Beetle             | 26           | 70           |
| Subalpine Fir Mortality        | 30           | 209          | Mountain Pine Beetle (PP)      | 835          | 1,179        |
| MPB (High Elev 5-needle Pines) | 50,794       | 191,143      | Mountain Pine Beetle (LPP)     | 3,832        | 16,936       |
| Western Spruce Budworm         | 43,405       | 0            | Subalpine Fir Mortality        | 1,949        | 1,953        |
| <b>Madison RD</b>              | <b>Acres</b> | <b>Trees</b> | MPB (High Elev 5-needle Pines) | 1,831        | 2,672        |
| Douglas-fir Beetle             | 12           | 58           | Western Spruce Budworm         | 1,787        | 0            |
| Engelmann Spruce Beetle        | 2            | 6            | <b>Sula RD</b>                 | <b>Acres</b> | <b>Trees</b> |
| Mountain Pine Beetle (LPP)     | 46,398       | 175,268      | Douglas-fir Beetle             | 36           | 76           |
| Subalpine Fir Mortality        | 334          | 1,144        | Pine Engraver Beetle (PP)      | 4            | 10           |
| MPB (High Elev 5-needle Pines) | 12,695       | 38,610       | Mountain Pine Beetle (PP)      | 69           | 86           |
| Western Spruce Budworm         | 18,478       | 0            | Mountain Pine Beetle (LPP)     | 631          | 1,057        |
| <b>Sheridan RD</b>             | <b>Acres</b> | <b>Trees</b> | Fir Engraver Beetle            | 8            | 10           |
| Douglas-fir Beetle             | 12           | 50           | Subalpine Fir Mortality        | 580          | 1,022        |
| Mountain Pine Beetle (LPP)     | 21,540       | 72,689       | Pine Engraver Beetle (LPP)     | 7            | 16           |
| Subalpine Fir Mortality        | 79           | 185          | MPB (High Elev 5-needle Pines) | 175          | 345          |
| MPB (High Elev 5-needle Pines) | 27,821       | 116,308      | Elytroderma                    | 22           | 0            |
| Western Spruce Budworm         | 15,673       | 0            | <b>West Fork RD</b>            | <b>Acres</b> | <b>Trees</b> |
| Windthrow                      | 25           | 0            | Douglas-fir Beetle             | 96           | 172          |
| Aspen Decline                  | 12           | 0            | Mountain Pine Beetle (PP)      | 89           | 56           |
| <b>Wisdom RD</b>               | <b>Acres</b> | <b>Trees</b> | Mountain Pine Beetle (LPP)     | 8,079        | 13,122       |
| Douglas-fir Beetle             | 22           | 194          | Fir Engraver Beetle            | 6            | 10           |
| Mountain Pine Beetle (LPP)     | 9,576        | 17,599       | Subalpine Fir Mortality        | 1,955        | 1,952        |
| Subalpine Fir Mortality        | 36           | 252          | MPB (High Elev 5-needle Pines) | 22           | 45           |
| MPB (High Elev 5-needle Pines) | 2,571        | 4,652        |                                |              |              |
| Western Spruce Budworm         | 1,204        | 0            |                                |              |              |
| <b>Wise River RD</b>           | <b>Acres</b> | <b>Trees</b> | <b>CUSTER NF</b>               |              |              |
| Douglas-fir Beetle             | 6            | 33           | <b>Ashland RD</b>              | <b>Acres</b> | <b>Trees</b> |
| Mountain Pine Beetle (LPP)     | 92,961       | 346,439      | Mountain Pine Beetle (PP)      | 123          | 130          |
| MPB (High Elev 5-needle Pines) | 16,609       | 58,187       | Western Spruce Budworm         | 304          | 0            |
| Western Spruce Budworm         | 8,406        | 0            | <b>Beartooth RD</b>            | <b>Acres</b> | <b>Trees</b> |
|                                |              |              | Douglas-fir Beetle             | 2,461        | 6,751        |
|                                |              |              | Mountain Pine Beetle (LPP)     | 823          | 1,624        |
|                                |              |              | Subalpine Fir Mortality        | 782          | 1,698        |
|                                |              |              | MPB (High Elev 5-needle Pines) | 7,906        | 28,269       |
|                                |              |              | Western Spruce Budworm         | 683          | 0            |
|                                |              |              | Unidentified Defoliator        | 11           | 0            |
|                                |              |              | Windthrow                      | 15           | 0            |
|                                |              |              | <b>Sioux RD</b>                | <b>Acres</b> | <b>Trees</b> |
|                                |              |              | Mountain Pine Beetle (PP)      | 77           | 93           |
|                                |              |              | Aspen Decline                  | 2            | 0            |
|                                |              |              |                                |              |              |
| <b>BITTERROOT NF</b>           |              |              |                                |              |              |
| <b>Darby RD</b>                | <b>Acres</b> | <b>Trees</b> |                                |              |              |
| Douglas-fir Beetle             | 65           | 112          |                                |              |              |
| Mountain Pine Beetle (PP)      | 145          | 193          |                                |              |              |
| Mountain Pine Beetle (LPP)     | 461          | 976          |                                |              |              |
| Fir Engraver Beetle            | 4            | 4            |                                |              |              |
| Subalpine Fir Mortality        | 1,102        | 903          |                                |              |              |



|                                |              |              |  |                            |              |              |
|--------------------------------|--------------|--------------|--|----------------------------|--------------|--------------|
| <b>Gardiner RD</b>             | <b>Acres</b> | <b>Trees</b> |  |                            |              |              |
| Douglas-fir Beetle             | 221          | 455          |  | Mountain Pine Beetle (PP)  | 297          | 303          |
| Mountain Pine Beetle (LPP)     | 17,944       | 79,125       |  | Mountain Pine Beetle (LPP) | 41,492       | 46,035       |
| Subalpine Fir Mortality        | 1,675        | 6,167        |  | Fir Engraver Beetle        | 16           | 20           |
| MPB (High Elev 5-needle Pines) | 13,831       | 55,543       |  | Subalpine Fir Mortality    | 16           | 28           |
| Western Spruce Budworm         | 22,707       | 0            |  | Western Spruce Budworm     | 89,634       | 0            |
| <b>Hebgen Lake RD</b>          | <b>Acres</b> | <b>Trees</b> |  | Larch Needle Blight        | 60           | 0            |
| Douglas-fir Beetle             | 4            | 15           |  | <b>Fisher River RD</b>     | <b>Acres</b> | <b>Trees</b> |
| Mountain Pine Beetle (LPP)     | 72,556       | 383,529      |  | Douglas-fir Beetle         | 43           | 44           |
| Subalpine Fir Mortality        | 3,830        | 11,032       |  | Engelmann Spruce Beetle    | 6            | 6            |
| MPB (High Elev 5-needle Pines) | 35,555       | 106,032      |  | Mountain Pine Beetle (WP)  | 4            | 4            |
| Western Spruce Budworm         | 38,343       | 0            |  | Mountain Pine Beetle (PP)  | 197          | 202          |
| <b>Livingston RD</b>           | <b>Acres</b> | <b>Trees</b> |  | Mountain Pine Beetle (LPP) | 2,850        | 7,467        |
| Douglas-fir Beetle             | 373          | 1,371        |  | Western Pine Beetle        | 72           | 59           |
| Mountain Pine Beetle (LPP)     | 67,691       | 558,102      |  | Fir Engraver Beetle        | 154          | 96           |
| Subalpine Fir Mortality        | 10,369       | 21,500       |  | Subalpine Fir Mortality    | 12           | 30           |
| MPB (High Elev 5-needle Pines) | 46,399       | 218,300      |  | Western Spruce Budworm     | 52,637       | 0            |
| Western Spruce Budworm         | 22,475       | 0            |  | White Pine Blister Rust    | 239          | 29           |
|                                |              |              |  | <b>Fortine RD</b>          | <b>Acres</b> | <b>Trees</b> |
|                                |              |              |  | Douglas-fir Beetle         | 197          | 284          |
|                                |              |              |  | Engelmann Spruce Beetle    | 2            | 4            |
| <b>HELENA NF</b>               |              |              |  | Mountain Pine Beetle (PP)  | 4            | 4            |
| <b>Helena RD</b>               | <b>Acres</b> | <b>Trees</b> |  | Mountain Pine Beetle (LPP) | 1,711        | 5,138        |
| Douglas-fir Beetle             | 6            | 52           |  | Western Pine Beetle        | 2            | 2            |
| Mountain Pine Beetle (PP)      | 72,839       | 208,090      |  | Fir Engraver Beetle        | 4            | 11           |
| Mountain Pine Beetle (LPP)     | 219,161      | 2,445,219    |  | Subalpine Fir Mortality    | 3,956        | 13,597       |
| MPB (High Elev 5-needle Pines) | 4,474        | 35,238       |  | Western Spruce Budworm     | 13,773       | 0            |
| Western Spruce Budworm         | 102,597      | 0            |  | Windthrow                  | 2            | 0            |
| Avalanche                      | 11           | 0            |  | <b>Libby RD</b>            | <b>Acres</b> | <b>Trees</b> |
| <b>Lincoln RD</b>              | <b>Acres</b> | <b>Trees</b> |  | Douglas-fir Beetle         | 77           | 89           |
| Douglas-fir Beetle             | 46           | 88           |  | Mountain Pine Beetle (WP)  | 4            | 4            |
| Mountain Pine Beetle (PP)      | 11,380       | 1,742        |  | Mountain Pine Beetle (PP)  | 143          | 169          |
| Mountain Pine Beetle (LPP)     | 180,918      | 764,496      |  | Mountain Pine Beetle (LPP) | 134          | 114          |
| Subalpine Fir Mortality        | 20           | 67           |  | Western Pine Beetle        | 46           | 33           |
| MPB (High Elev 5-needle Pines) | 741          | 1,494        |  | Fir Engraver Beetle        | 119          | 88           |
| Western Spruce Budworm         | 138,642      | 0            |  | Subalpine Fir Mortality    | 40           | 83           |
| Windthrow                      | 16           | 0            |  | Western Spruce Budworm     | 97,091       | 0            |
| <b>Townsend RD</b>             | <b>Acres</b> | <b>Trees</b> |  | White Pine Blister Rust    | 161          | 44           |
| Douglas-fir Beetle             | 55           | 336          |  | <b>Rexford RD</b>          | <b>Acres</b> | <b>Trees</b> |
| Mountain Pine Beetle (PP)      | 4,208        | 45,605       |  | Douglas-fir Beetle         | 6            | 6            |
| Mountain Pine Beetle (LPP)     | 81,985       | 819,187      |  | Pine Engraver Beetle (PP)  | 2            | 10           |
| MPB (High Elev 5-needle Pines) | 9,851        | 56,460       |  | Mountain Pine Beetle (WP)  | 4            | 4            |
| Western Spruce Budworm         | 89,802       | 0            |  | Mountain Pine Beetle (PP)  | 129          | 167          |
| Windthrow                      | 40           | 0            |  | Mountain Pine Beetle (LPP) | 325          | 344          |
|                                |              |              |  | Fir Engraver Beetle        | 25           | 16           |
|                                |              |              |  | Subalpine Fir Mortality    | 259          | 545          |
| <b>KOOTENAI NF</b>             |              |              |  | Western Spruce Budworm     | 460          | 0            |
| <b>Cabinet RD</b>              | <b>Acres</b> | <b>Trees</b> |  | White Pine Blister Rust    | 19           | 0            |
| Douglas-fir Beetle             | 97           | 186          |  |                            |              |              |
| Mountain Pine Beetle (WP)      | 4            | 4            |  |                            |              |              |



|                                |              |              |  |                                |              |              |
|--------------------------------|--------------|--------------|--|--------------------------------|--------------|--------------|
| Pine Engraver Beetle (PP)      | 616          | 1,500        |  | <b>FORT BELKNAP IR</b>         | <b>Acres</b> | <b>Trees</b> |
| Mountain Pine Beetle (PP)      | 158          | 202          |  | Douglas-fir Beetle             | 6            | 6            |
| Mountain Pine Beetle (LPP)     | 45,573       | 66,508       |  | Mountain Pine Beetle (LPP)     | 236          | 615          |
| Western Pine Beetle            | 50           | 98           |  | Mountain Pine Beetle (PP)      | 522          | 1,620        |
| Fir Engraver Beetle            | 852          | 927          |  | Western Spruce Budworm         | 43           | 0            |
| Subalpine Fir Mortality        | 151          | 450          |  | Windthrow                      | 26           | 0            |
| Western Spruce Budworm         | 68,107       | 0            |  |                                |              |              |
|                                |              |              |  | <b>NORTHERN CHEYENNE IR</b>    | <b>Acres</b> | <b>Trees</b> |
| <b>BLACKFEET IR</b>            | <b>Acres</b> | <b>Trees</b> |  | Mountain Pine Beetle (PP)      | 115          | 126          |
| Douglas-fir Beetle             | 582          | 61           |  | MPB (High Elev 5-needle Pines) | 2            | 2            |
| Engelmann Spruce Beetle        | 2            | 2            |  | Western Spruce Budworm         | 491          | 0            |
| Fir Engraver Beetle            | 801          | 370          |  |                                |              |              |
| Mountain Pine Beetle (LPP)     | 584          | 731          |  | <b>ROCKY BOYS IR</b>           | <b>Acres</b> | <b>Trees</b> |
| Mountain Pine Beetle (PP)      | 4            | 6            |  | Douglas-fir Beetle             | 24           | 60           |
| Subalpine Fir Mortality        | 107          | 72           |  | Mountain Pine Beetle (LPP)     | 5,738        | 74,887       |
| Western Spruce Budworm         | 4,334        | 0            |  | Mountain Pine Beetle (PP)      | 569          | 3,145        |
| Winter Injury                  | 27           | 0            |  | Western Pine Beetle            | 2            | 2            |
|                                |              |              |  | Western Spruce Budworm         | 10,454       | 0            |
| <b>CROW IR</b>                 | <b>Acres</b> | <b>Trees</b> |  |                                |              |              |
| Aspen Decline                  | 227          | 0            |  | <b>GLACIER NP</b>              | <b>Acres</b> | <b>Trees</b> |
| Douglas-fir Beetle             | 125          | 96           |  | Avalanche                      | 19           | 0            |
| Mountain Pine Beetle (LPP)     | 4            | 14           |  | Douglas-fir Beetle             | 2,939        | 3,598        |
| Mountain Pine Beetle (PP)      | 89           | 152          |  | Engelmann Spruce Beetle        | 4            | 3            |
| MPB (High Elev 5-needle Pines) | 14           | 40           |  | Fir Engraver Beetle            | 1,632        | 1,744        |
| Subalpine Fir Mortality        | 4            | 8            |  | Flooding - High Water          | 2            | 0            |
| Western Spruce Budworm         | 991          | 0            |  | Mountain Pine Beetle (LPP)     | 4,450        | 6,503        |
|                                |              |              |  | Mountain Pine Beetle (PP)      | 2            | 2            |
| <b>FLATHEAD IR</b>             | <b>Acres</b> | <b>Trees</b> |  | MPB (High Elev 5-needle Pines) | 6            | 10           |
| Douglas-fir Beetle             | 259          | 254          |  | Subalpine Fir Mortality        | 7,509        | 16,164       |
| Fir Engraver Beetle            | 11           | 19           |  | Western Spruce Budworm         | 15,716       | 0            |
| Flooding - High Water          | 2            | 0            |  | Windthrow                      | 2            | 0            |
| Mountain Pine Beetle (LPP)     | 16,933       | 95,919       |  |                                |              |              |
| Mountain Pine Beetle (PP)      | 2,706        | 2,236        |  | <b>YELLOWSTONE NP</b>          | <b>Acres</b> | <b>Trees</b> |
| MPB (High Elev 5-needle Pines) | 160          | 467          |  | Aspen Decline                  | 71           | 0            |
| Pine Engraver Beetle (PP)      | 40           | 71           |  | Douglas-fir Beetle             | 79           | 240          |
| Rhabdocline pseudotsugae       | 2,060        | 0            |  | Engelmann Spruce Beetle        | 18           | 20           |
| Subalpine Fir Mortality        | 2,017        | 4,773        |  | Flooding - High Water          | 248          | 0            |
| Unidentified Defoliator        | 228          | 0            |  | Mountain Pine Beetle (LPP)     | 58,393       | 166,269      |
| Western Pine Beetle            | 20           | 20           |  | Mountain Pine Beetle (PP)      | 462          | 922          |
| Western Spruce Budworm         | 12,296       | 0            |  | MPB (High Elev 5-needle Pines) | 41,001       | 112,264      |
| Winter Injury                  | 13           | 0            |  | Subalpine Fir Mortality        | 2,343        | 4,270        |
|                                |              |              |  | Western Spruce Budworm         | 41,660       | 0            |
|                                |              |              |  | Windthrow                      | 71           | 0            |

## COMMON AND SCIENTIFIC NAMES

### Pathogens

|                                |  |                                |
|--------------------------------|--|--------------------------------|
| Annosus root disease           | <i>Heterobasidion annosum</i> (Fr.:Fr.) Bref.              | Primary hosts: DF, GF, PP, SAF |
| Armillaria root disease        | <i>Armillaria ostoyae</i> (Romagn.) Herink                 | DF, GF, SAF, sapling pines     |
| Black stain root disease       | <i>Leptographium wageneri</i> (Kendrick) M.J. Wingfield    | DF, PP                         |
| Brown cubical butt rot         | <i>Phaeolus schweinitzii</i> (Fr.:Fr.) Pat.                | DF                             |
| Diplodia shoot blight          | <i>Diplodia pinea</i> (Desmaz.) J. Kickx fil.              | PP                             |
| Dothistroma needle cast        | <i>Dothistroma septospora</i> (Doroguine) Morelet          | LP, PP, WWP, LPP, WBP          |
| Dwarf mistletoes               | <i>Arceuthobium</i> spp.                                   | LPP, LP, DF, WL                |
| Brown stringy rot              | <i>Echinodontium tinctorium</i> (Ell. & Ev.) Ell. & Ev.    | GF, WH                         |
| Elytroderma needle cast        | <i>Elytroderma deformans</i> (Weir) Darker                 | PP                             |
| Laminated root rot             | <i>Phellinus weirii</i> (Murrill) R.L. Gilbertson.         | DF, GF, WH, SAF                |
| Larch casebearer               | <i>Coleophora laricella</i> (Hubner)                       | WL                             |
| Larch needle blight            | <i>Hypodermella laricis</i> Tub.                           | WL                             |
| Larch needle cast              | <i>Meria laricis</i> Vuill.                                | WL                             |
| Lodgepole pine needle cast     | <i>Lophodermella concolor</i> (Dearn.) Darker              | LPP                            |
| Rhabdocline needle cast        | <i>Rhabdocline</i> spp.                                    | DF                             |
| Schweinitzii root and butt rot | <i>Phaeolus schweinitzii</i> (Fr. :Fr.) Pat.               | DF                             |
| Tomentosus root disease        | <i>Inonotus tomentosus</i> (Fr.) Teng.                     | ES, DF, LPP                    |
| Western gall rust              | <i>Endocronartium harknessii</i> (J.P. Moore) Y. Hiratsuka | LPP, PP                        |
| White pine blister rust        | <i>Cronartium ribicola</i> J.C. Fisch.                     | WWP, WBP, LP                   |

### Insects

|                                      |   |                           |
|--------------------------------------|---|---------------------------|
| Douglas-fir beetle                   | <i>Dendroctonus pseudotsugae</i> Hopkins      | DF                        |
| Douglas-fir tussock moth             | <i>Orygia pseudotsugata</i> (McDunnough)      | DF, TF, ES                |
| Gypsy moth                           | <i>Lymantria dispar</i> (Linnaeus)            | Most hardwoods            |
| Mountain pine beetle                 | <i>Dendroctonus ponderosae</i> Hopkins        | All pines                 |
| Pine engraver beetle                 | <i>Ips pini</i> (Say)                         | PP, LPP                   |
| Spruce beetle                        | <i>Dendroctonus rufipennis</i> Swaine         | ES                        |
| Western balsam bark beetle           | <i>Dryocoetes confuses</i> Swaine             | SAF                       |
| Western pine (Grizzled) tussock moth | <i>Dasychira pinicola</i> (Dyar)              | PP (DF,ES,SAF,GF,WL, LPP) |
| Western pine beetle                  | <i>Dendroctonus brevicomis</i> LeConte        | PP                        |
| Fir engraver beetle                  | <i>Scolytis ventralis</i> LeConte             | GF, SAF                   |
| Hemlock looper                       | <i>Lambdina fiscellaria lugubrosa</i> (Hulst) | DF                        |
| False hemlock looper                 | <i>Nepytia canosaria</i> (Walker)             | DF                        |

DF = Douglas-fir; GF = Grand fir; TF = True fir; SAF = Subalpine fir; PP = Ponderosa pine; LP = Limber pine; LPP = Lodgepole pine; WWP = Western white pine; ES = Engelmann spruce; WH = Western hemlock; WL = Western larch; WBP = Whitebark pine

## **PUBLICATIONS 2009**

DeNitto, Gregg; Gibson, Ken. 2009. Evaluation of Several Insect/Disease Concerns on the Three Rivers and Libby RDs, July 7-9, 2009. Missoula Field Office. R1. MFO-TR-09-13. July 30, 2009.

Gibson, Ken. 2009. Bark Beetle Potential in Trees Cut and Left on Site, BPA Power Line Right-of-Way, Libby Ranger District, April 22, 2009. Missoula Field Office. R1. MFO-TR-09-05. May 7, 2009.

Gibson, Ken. 2009. Down Trees in BPA Power Line Right-of-Way, Follow-up, Libby Ranger District, May 13, 2009. Missoula Field Office. R1. MFO-TR-09-08. May 21, 2009.

Gibson, Ken. 2009. Western Spruce Budworm and Mountain Pine Beetle Three Saddle Analysis Area Stevensville Ranger District, July 31, 2009. Missoula Field Office. R1. MFO-TR-09-16. August 17, 2009.

Gibson, Ken. 2009. Mountain Pine Beetle Activity, Hebgen Lake Ranger District, August 17-18, 2009. Missoula Field Office. R1. MFO-TR-09-18. August 27, 2009.

Gibson, Ken. 2009. Bark Beetle Threats in Windthrown and Fire-Damaged Trees Beartooth Ranger District August 20, 2009. Missoula Field Office. R1. MFO-TR-09-19. August 27, 2009.

Gibson, Ken. 2009. Assistance to DNRC Service Forester, Polson, MT, August 28, 2009. Missoula Field Office. R1. MFO-TR-09-22. September 4, 2009.

Gibson, Ken. 2009. Vegetation Management Plan-Impacts of Forest Insects and Diseases, Bridger Bowl Ski Area, Bozeman, MT, October 15, 2009. Missoula Field Office. R1. MFO-TR-09-34. October 22, 2009.

Gibson, Ken. 2009. Bark Beetle Conditions at Rocky Gorge Campground, Rexford Ranger District, October 22, 2009. Missoula Field Office. R1. MFO-TR-09-37. October 22, 2009.

Gibson, Ken; DeNitto, Gregg. 2009. Hazard Trees in Warm Springs Campground, Sula Ranger District, May 5, 2009. Missoula Field Office. R1. MFO-TR-09-07. May 7, 2009.

Gibson, Ken; Jackson, Marcus. Management Alternatives, Wise River Restoration Area, Bureau of Land Management, Butte Field Office, March 20, 2009. Missoula Field Office. R1. MFO-TR-09-04. March 20, 2009.

Gibson, Ken; Jackson, Marcus. 2009. Bigfork Tree Improvement Area, Swan Lake RD Disease and Insect Concerns, July 6, 2009. Missoula Field Office. R1. MFO-TR-09-12.

Gibson, Ken; Steed, Brytten. 2009. Bark Beetle Status and Management, Fort Belknap Reservation 1 September 2009. Missoula Field Office. R1. MFO-TR-09-24. September 10, 2009.

Gibson, Ken; Steed, Brytten. 2009. Right-of-Way clearing, West Fork Highway, West Fork Ranger District, October 2, 2009. Missoula Field Office. R1. MFO-TR-09-28. October 8, 2009.

Jackson, Marcus. 2009. North Shore and Bull River Campgrounds, Kootenai NF Disease and Insect Evaluation, June 8, 2009. Missoula Field Office. R1. MFO-TR-09-09. July 6, 2009.

Jackson, Marcus. 2009. Tally Lake Campground October 13, 2009. Missoula Field Office. R1. MFO-TR-09-32. October 28, 2009.

Jackson, Marcus. 2009. Dwarf Mistletoe Suppression Work on the Flathead Indian Reservation, December 3, 2009. Missoula Field Office. R1. MFO-TR-09-43. December 15, 2009.

Jackson, Marcus; Cramer, William. 2009. Ft. Belknap Limber Pine October 15, 2009. Missoula Field Office. R1. MFO-TR-09-33. October 28, 2009.

Jackson, Marcus; Sturdevant, Nancy. 2009. Ashley-Herrig Resource management Project, Disease and Insect Evaluation, June 15, 2009. Missoula Field Office. R1. MFO-TR-09-10. June 24, 2009.

Jackson, Marcus; Sturdevant, Nancy. 2009. Spotted Bear River Project, September 4, 2009. Missoula Field Office. R1. MFO-TR-09-26. October 8, 2009.

Kegley, Sandy; Gibson, Ken. 2009. Individual-Tree Tests of Verbenone and Green-leaf Volatiles to Protect Lodgepole, Whitebark and Ponderosa Pines, 2004-2007. Missoula, MT: USDA Forest Service, Region 1, Forest Health Protection. Numbered Report 09-03. 12p.

Kegley, Sandy; Schwandt, John; Gibson, Ken. 2009. Mountain Pine Beetle and Blister Rust Impacts, Yellowstone Club, Big Sky, Montana, August 11, 2009. Missoula Field Office. R1. MFO-TR-09-17. September 22, 2009.

Lockman, Blakey. 2009. Three Saddle Project Area, Stevensville RD, Bitterroot NF, October 8, 2009. Missoula Field Office. R1. MFO-TR-09-35. October 22, 2009.

Lockman, Blakey. 2009. Stump Treatments at Rocky gorge CG and Ten Allen Sale Area Rexford RD, Kootenai NF December 10, 2009. Missoula Field Office. R1. MFO-TR-09-44. January 5, 2010.

Lockman, Blakey; DeNitto, Gregg. 2009. WBP in Cedar-Thom Analysis Area, Superior RD, LoLo NF, August 18, 2009. Missoula Field Office. R1. MFO-TR-09-23. October 9, 2009.

Lockman, Blakey; Gibson, Ken. 2009. CEEM Certification Stand Review and Blowdown, Pierce Fuel Reduction Unit 2, Swan Lake Ranger District, June 26, 2009. Missoula Field Office. R1. MFO-TR-09-11. August 6, 2009.

Lockman, Blakey; Gibson, Ken. 2009. PSR Proposals, Superior Ranger District, August 26, 2009. Missoula Field Office. R1. MFO-TR-09-20. September 4, 2009.

Lockman, Blakey; Jackson, Marcus. 2009. Root Disease in Cedar-Thom Analysis Area, Superior RD. Missoula Field Office. R1. MFO-TR-09-06. May 7, 2009.

Lockman, Blakey; Sturdevant, Nancy. 2009. Pilgrim Creek Analysis Area, Cabinet Ranger District, June 16, 2009. Missoula Field Office. R1. MFO-TR-09-21. September 10, 2009.

Steed, Brytten. 2009. Lewis and Clark NF, Proposed FY2010 PSR Projects, August 3-7, 2009. Missoula Field Office. R1. MFO-TR-09-29. October 9, 2009.

Steed, Brytten. 2009. Beaverhead-Deerlodge NF Review of FY 2009 Funded PSR projects and FY 2010 proposals, June 3, 2009; August 10-13 & 17-18, 2009. Missoula Field Office. R1. MFO-TR-09-30. October 9, 2009.

Steed, Brytten. 2009. Mountain Pine Beetle at Piney and Spring Hill Campgrounds: Mortality of trees marked for spray. Missoula Field Office. R1. MFO-TR-09-38. October 29, 2009.

Steed, Brytten. 2009. Weather-related Mountain Pine Beetle Brood Mortality on the B-D NF. Missoula Field Office. R1. MFO-TR-09-40. November 24, 2009.

Steed, Brytten. 2009. 2009 FINDIT Survey Summary for the Beaverhead-Deerlodge NF. Missoula Field Office. R1. MFO-TR-09-41. December 2, 2009.

Steed, Brytten; Gibson, Ken. 2009. Barton Springs-Mountain Pine Beetle, Follow-up, Pintler Ranger District, Beaverhead-Deerlodge National Forest, February 19, 2009. Missoula Field Office. R1. MFO-TR-09-02. 2009.

Steed, Brytten; Gibson, Ken. 2009. Pine Engraver Beetle-Caused Mortality CMR National Wildlife Refuge and Fort Peck Recreation Area Fort Peck MT September 2, 2009. Missoula Field Office. R1. MFO-TR-09-25. September 10, 2009.

Steed, Brytten; Lockman, Blakey; Jackson, Marcus. 2009. Lewis and Clark NF Disease Review, June 8-9, 2009. Missoula Field Office. R1. MFO-TR-09-31. October 13, 2009.

Sturdevant, Nancy. 2009. Bark Beetle Activity in the Grasshopper Salvage and Fuels Management Project, BLM, January 8, 2008. Missoula Field Office. R1. MFO-TR-09-01. February 11, 2009.

Sturdevant, Nancy. 2009. Potential for Bark Beetle Activity in Blowdown, Resulting from the Mountain Wave Wind Event on the Fortine R. D., Kootenai N. F, July 1, 2009. Missoula Field Office. R1. MFO-TR-09-14. July 20, 2009.

Sturdevant, Nancy. 2009. Mountain Pine Beetle Suppression Review and Pine Engraver Activity at Great Divide and Deep Creek. Missoula Field Office. R1. MFO-TR-09-15. August 5, 2009.

Sturdevant, Nancy. 2009. Flathead PSR Service Trip, August 11, 2009. Missoula Field Office. R1. MFO-TR-09-27. October 7, 2009.

Sturdevant, Nancy. 2009. Beetle Trends in the Yank Swamp Project Area, BLM, Billon Field Office. Missoula Field Office. R1. MFO-TR-09-36. November 3, 2009.

Sturdevant, Nancy. 2009. Mountain Pine Beetle Activity in the Little Gravelies Fuels Reduction Project November 24, 2009. Missoula Field Office. R1. MFO-TR-09-42. January 5, 2010.

Sturdevant, Nancy; Cramer, William. 2009. Douglas-fir Beetle Activity Following Thinning and Burning on Private, State and Forest Service Lands in the Centennial Valley, March 3, 2009. Missoula Field Office. R1. MFO-TR-09-03, 2009.

Sturdevant, Nancy; Gibson, Ken; Cramer, William. 2009. Protecting Lodgepole Pine from Mountain Pine Beetle Attack Using Verbenone and Green-Leaf Volatiles Rocky Boys Indian Reservation, 2008. Missoula, MT: USDA Forest Service, Region 1, Forest Health Protection. Numbered Report 09-05. 5 p.

Sturdevant, Nancy; Jackson, Marcus. 2009. Insect and Disease Conditions on Peter's Ridge. Swan Lake Ranger District, Flathead NF, December 21, 2009. Missoula Field Office. R1. MFO-TR-09-39, January 13, 2010.

## DIRECTORY OF PERSONNEL

### Montana Department of Natural Resources and Conservation

Phone: (406) 542-4300

|                  |                                   |                                     |                |
|------------------|-----------------------------------|-------------------------------------|----------------|
| Bob Harrington   | State Forester                    | bharrington@state.mt.us             | 4301           |
| Rob Ethridge     | Chief, Service Forestry Bureau    | rethridge@state.mt.us               | 4303           |
| Amy Gannon       | Forest Pest Management Specialist | agannon@state.mt.us                 | 4283           |
| Brennan Ferguson | Contract Forest Pathologist       | brennan@fergusonforestpathology.com | (406) 239-7761 |

Montana Department of Natural Resources and Conservation, Forestry Division  
2705 Spurgin Road, Missoula, Montana 59804



### USDA Forest Service Regional Office

Phone: (406) 329-3308

|                          |                                      |                     |                |
|--------------------------|--------------------------------------|---------------------|----------------|
| Mike Dudley <sup>1</sup> | Director, S&PF                       | mdudley@fs.fed.us   | (801) 625-5253 |
| Peggy Polichio           | Deputy Director, S&PF                | ppolichio@fs.fed.us | (406) 329-3280 |
| Rob Cruz <sup>1</sup>    | Forest Health Monitoring Coordinator | rcruz@fs.fed.us     | (801) 625-5162 |
| Janet Valle <sup>1</sup> | Pesticide Use Coordinator            | jvalle@fs.fed.us    | (801) 625-5258 |

USDA Forest Service, Northern Region, Federal Building,  
200 East Broadway, P.O. Box 7669, Missoula, Montana 59807

<sup>1</sup> USDA Forest Service, Intermountain Region  
324 25<sup>th</sup> Street, Ogden, UT 84401



**DIRECTORY OF PERSONNEL, continued**

**USDA Forest Service Missoula Field Office**

Phone: (406) 329-3277

|                   |                                  |                       |      |
|-------------------|----------------------------------|-----------------------|------|
| Gregg DeNitto     | Group Leader                     | gdenitto@fs.fed.us    | 3637 |
| Mark Casey        | Bio. Sci. Tech., Aerial Surveyor | mcasey@fs.fed.us      | 3136 |
| William Cramer    | Bio. Science Technician          | wcramer@fs.fed.us     | 3130 |
| Ken Gibson        | Entomologist                     | kgibson@fs.fed.us     | 3278 |
| Marcus Jackson    | Plant Pathologist                | mbjackson@fs.fed.us   | 3282 |
| Gary Little       | Bio. Sci. Tech., Aerial Surveyor |                       | 3051 |
| Blakey Lockman    | Plant Pathologist                | blockman@fs.fed.us    | 3180 |
| William O'Donnell | GIS Coordinator                  | wodonnell@fs.fed.us   | 3502 |
| Scott Sontag      | Bio. Sci. Tech., Aerial Surveyor | ssontag@fs.fed.us     | 3323 |
| Brytten Steed     | Entomologist                     | bsteed@fs.fed.us      | 3142 |
| Nancy Sturdevant  | Entomologist                     | nsturdevant@fs.fed.us | 3281 |

USDA Forest Service, Northern Region, Federal Building,  
200 East Broadway, P.O. Box 7669, Missoula, Montana 59807

|  |  |                    |              |
|--|--|--------------------|--------------|
| Contract Digitizing<br>and Map<br>Production | International Forestry<br>Consultants Inc. | www.inforestry.com | 425-820-3420 |
|--|--|--------------------|--------------|



**Table 2. Acres of Host Type Infested by Bark Beetles  
on all Ownerships in Montana, during 2009**

|                             | USFS      | Other Fed | State   | Private | Total     |
|-----------------------------|-----------|-----------|---------|---------|-----------|
| <b>DFB<sup>1</sup></b>      | 14,143    | 4,773     | 537     | 3,075   | 22,528    |
| <b>ESB<sup>1</sup></b>      | 45        | 28        | 8       | 4       | 85        |
| <b>FE<sup>1</sup></b>       | 3,033     | 2,436     | 378     | 3,340   | 9,187     |
| <b>IPS<sup>1</sup></b>      | 618       | 6         | 28      | 153     | 805       |
| <b>MPB-LPP<sup>1</sup></b>  | 2,140,525 | 199,518   | 79,888  | 444,303 | 2,864,234 |
| <b>MPB-PP<sup>1</sup></b>   | 113,809   | 38,180    | 31,162  | 242,282 | 425,433   |
| <b>MPB-5-NP<sup>1</sup></b> | 315,197   | 49,848    | 3,185   | 36,241  | 404,471   |
| <b>MPB-WWP<sup>1</sup></b>  | 22        | 0         | 4       | 0       | 26        |
| <b>MPB-All<sup>1</sup></b>  | 2,569,553 | 287,546   | 114,239 | 722,826 | 3,694,164 |
| <b>SAF<sup>1</sup></b>      | 62,060    | 11,898    | 1,149   | 7,398   | 82,505    |
| <b>WPB<sup>1</sup></b>      | 557       | 22        | 61      | 176     | 816       |
| <b>WSBW<sup>1</sup></b>     | 1,446,987 | 228,746   | 112,613 | 813,586 | 2,601,932 |

<sup>1</sup>DFB = Douglas-fir beetle; ESB = Spruce beetle; FE = Fir engraver beetle; IPS = Pine engraver beetle;  
 MPB LPP = Mountain pine beetle in Lodgepole pine; MPB PP = Mountain Pine Beetle in Ponderosa pine;  
 MPB 5-NP = Mountain pine beetle in 5-Needle pines (WBP & LP); MPB WP = Mountain pine beetle in Western white pine;  
 MPB All = Mountain pine beetle in all pines; SAF = Subalpine fir mortality complex; WPB = Western pine beetle;  
 WSBW = Western spruce budworm.

**Table 3. Acres of Host Type Infested by Bark Beetles in Montana, From 2007 Through 2009**

| Insect           | 2007 <sup>2,3</sup> |                  | 2008 <sup>2,3</sup> |                   | 2009 <sup>2,3</sup> |                   |
|------------------|---------------------|------------------|---------------------|-------------------|---------------------|-------------------|
|                  | Acres               | Trees            | Acres               | Trees             | Acres               | Trees             |
| DFB <sup>1</sup> | 22,285              | 58,773           | 21,558              | 39,587            | 22,528              | 52,873            |
| ESB <sup>1</sup> | 403                 | 572              | 54                  | 60                | 75                  | 100               |
| FE <sup>1</sup>  | 1,657               | 1,634            | 359                 | 803               | 9,187               | 7,842             |
| IPS <sup>1</sup> | 1,187               | 2,299            | 10,778              | 36,669            | 805                 | 1,751             |
| MPB <sup>1</sup> | 807,334             | 2,559,859        | 1,819,245           | 16,903,542        | 3,694,164           | 22,259,733        |
| SAF <sup>1</sup> | 115,250             | 286,757          | 53,194              | 149,802           | 82,505              | 245,595           |
| WPB <sup>1</sup> | 401                 | 243              | 167                 | 182               | 816                 | 1,302             |
|                  |                     |                  |                     |                   |                     |                   |
| <b>TOTAL</b>     | <b>948,517</b>      | <b>2,910,137</b> | <b>1,905,355</b>    | <b>17,130,645</b> | <b>3,810,080</b>    | <b>22,569,196</b> |

<sup>1</sup>DFB = Douglas-fir beetle; ESB = Spruce beetle; FE = Fir engraver beetle; IPS = Pine engraver beetle; MPB = Mountain pine beetle; SAF = Subalpine fir mortality complex; WPB = Western pine beetle.

<sup>2</sup>Not all areas were flown due to fires, inclement weather or seasonal limitations.

<sup>3</sup>Yellowstone NP includes MT, ID and WY acres.

**Table 4. Acres with Douglas-fir Beetle-Caused Mortality on All Ownerships in Montana, From 2007 Through 2009**

| Reporting Area  | 2007          |               | 2008          |               | 2009          |               |
|-----------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                 | Acres         | Trees         | Acres         | Trees         | Acres         | Trees         |
| Beaverhead      | 4,197*        | 11,138*       | 376           | 1,766         | 87            | 480           |
| Bitterroot      | 1,835*        | 5,430*        | 674*          | 1,933*        | 229           | 436           |
| Custer          | 1,086         | 4,302         | 85*           | 163*          | 2,941         | 7,552         |
| Deerlodge       | 107*          | 386*          | 411           | 1,219         | 539           | 2,175         |
| Flathead        | 5,450         | 9,020         | 7,899         | 12,844        | 4,969         | 8,797         |
| Gallatin        | 2,058*        | 9,803*        | 722*          | 3,515*        | 2,446         | 10,302        |
| Garnets         | 47*           | 154*          | 283           | 650           | 122           | 236           |
| Helena          | 419*          | 1442*         | 5,166*        | 3,859*        | 1,707         | 1,062         |
| Kootenai        | 724*          | 805*          | 205*          | 289*          | 466*          | 684*          |
| Lewis & Clark   | 203*          | 630*          | 937*          | 2,075*        | 3,382         | 14,418        |
| Lolo            | 1,261*        | 3,029*        | 2,364         | 6,772         | 1,626*        | 2,416*        |
| Blackfeet IR    | ★             | ★             | 55            | 68            | 582           | 61            |
| Crow IR         | 348           | 1,371         | ★             | ★             | 125           | 96            |
| Flathead IR     | 552           | 1,102         | 51*           | 73*           | 259           | 254           |
| Fort Belknap IR | 13            | 57            | 0             | 0             | 6*            | 6*            |
| No. Cheyenne IR | 0             | 0             | ★             | ★             | 0             | 0             |
| Rocky Boy's IR  | 2             | 25            | 18            | 56            | 24            | 60            |
| Glacier NP      | 3,604*        | 8,814*        | 2,146         | 3,950         | 2,939*        | 3,598*        |
| Yellowstone NP  | 379           | 1,265         | 166           | 355           | 79            | 240           |
|                 |               |               |               |               |               |               |
| <b>TOTAL</b>    | <b>22,285</b> | <b>58,773</b> | <b>21,718</b> | <b>39,587</b> | <b>22,528</b> | <b>52,873</b> |

★ = Not surveyed \* = Partially surveyed

Yellowstone NP includes acres in MT, ID and WY

**Table 5. Acres with Mountain Pine Beetle-Caused Mortality on State and Private Lands in Montana, From 2007 Through 2009**

| Reporting Area  | 2007             |                 |                   |                  | 2008             |                 |                   |                  | 2009             |                 |                   |                  |
|-----------------|------------------|-----------------|-------------------|------------------|------------------|-----------------|-------------------|------------------|------------------|-----------------|-------------------|------------------|
|                 | LPP <sup>1</sup> | PP <sup>1</sup> | 5-NP <sup>1</sup> | WWP <sup>1</sup> | LPP <sup>1</sup> | PP <sup>1</sup> | 5-NP <sup>1</sup> | WWP <sup>1</sup> | LPP <sup>1</sup> | PP <sup>1</sup> | 5-NP <sup>1</sup> | WWP <sup>1</sup> |
| Beaverhead      | 6,074*           | 4*              | 918*              | 0*               | 26,874           | 0               | 3,064             | 0                | 31,580           | 0               | 4,095             | 0                |
| Bitterroot      | 6*               | 45*             | 0*                | 0*               | 36*              | 61*             | 0*                | 0*               | 424              | 890             | 0                 | 0                |
| Custer          | 217              | 145             | 6                 | 0                | 63*              | 527*            | 3*                | 0*               | 10               | 67              | 32                | 0                |
| Deerlodge       | 16,529*          | 21*             | 302*              | 0*               | 79,935           | 834             | 5,435             | 0                | 97,179           | 2,602           | 2,289             | 0                |
| Flathead        | 2,971            | 1,009           | 0                 | 0                | 2,619            | 795             | 2                 | 0                | 4,172            | 752             | 0                 | 0                |
| Gallatin        | 9,641*           | 56*             | 7,971*            | 0*               | 32,811*          | 21*             | 16,467*           | 0*               | 152,091          | 0               | 27,297            | 0                |
| Garnets         | 1,430*           | 84*             | 0*                | 0*               | 25,712           | 1,922           | 105               | 0                | 35,560           | 28,315          | 0                 | 0                |
| Helena          | 14,886*          | 6,312*          | 0*                | 0*               | 55,354*          | 24,009*         | 105*              | 0*               | 136,546          | 178,360         | 2,836             | 0                |
| Kootenai        | 4*               | 4*              | 0*                | 0*               | 951*             | 148*            | 0*                | 0*               | 1,894*           | 217*            | 0*                | 4*               |
| Lewis & Clark   | 735*             | 2,922*          | 55*               | 0*               | 4,191*           | 1,899*          | 183*              | 0*               | 30,605           | 50,617          | 2,877             | 0                |
| Lolo            | 5,309*           | 608*            | 0*                | 0*               | 11,079           | 531             | 0                 | 0                | 31,478*          | 10,777*         | 0*                | 0*               |
| Blackfeet IR    | ★                | ★               | ★                 | ★                | 44               | 0               | 0                 | 0                | 132              | 0               | 0                 | 0                |
| Crow IR         | 2                | 36              | 0                 | 0                | ★                | ★               | ★                 | ★                | 0                | 29              | 0                 | 0                |
| Flathead IR     | 1,399            | 109             | 0                 | 0                | 431*             | 203*            | 0*                | 0*               | 987              | 662             | 0                 | 0                |
| Fort Belknap IR | 0                | 2               | 0                 | 0                | 2                | 30              | 0                 | 0                | 10*              | 22*             | 0*                | 0*               |
| No. Cheyenne IR | 0                | 8               | 0                 | 0                | ★                | ★               | ★                 | ★                | 0                | 2               | 0                 | 0                |
| Rocky Boy's IR  | 593              | 62              | 0                 | 0                | 561              | 258             | 0                 | 0                | 1,507            | 132             | 0                 | 0                |
| Glacier NP      | 4*               | 0*              | 0*                | 0*               | 190              | 0               | 0                 | 0                | 16*              | 0*              | 0*                | 0*               |
| Yellowstone NP  | 0*               | 0*              | 0*                | 0*               | 0                | 0               | 0                 | 0                | 0                | 0               | 0                 | 0                |
|                 |                  |                 |                   |                  |                  |                 |                   |                  |                  |                 |                   |                  |
| <b>TOTAL</b>    | 59,800           | 11,427          | 9,252             | 0                | 240,853          | 31,238          | 25,364            | 0                | 524,191          | 273,444         | 39,426            | 4                |

<sup>1</sup>LPP = Lodgepole pine; PP = Ponderosa pine; 5-NP = 5-needle pines (WBP & LP); WWP = Western white pine

★ = Not surveyed; \* = Partially surveyed; Yellowstone NP includes MT, ID, and WY acres

**Table 6. Acres with Mountain Pine Beetle-Caused Mortality on All Federal Ownerships  
in Montana, From 2007 Through 2009**

| Reporting Area  | 2007             |                 |                   |                  | 2008             |                 |                   |                  | 2009             |                 |                   |                  |
|-----------------|------------------|-----------------|-------------------|------------------|------------------|-----------------|-------------------|------------------|------------------|-----------------|-------------------|------------------|
|                 | LPP <sup>1</sup> | PP <sup>1</sup> | 5-NP <sup>1</sup> | WWP <sup>1</sup> | LPP <sup>1</sup> | PP <sup>1</sup> | 5-NP <sup>1</sup> | WWP <sup>1</sup> | LPP <sup>1</sup> | PP <sup>1</sup> | 5-NP <sup>1</sup> | WWP <sup>1</sup> |
| Beaverhead      | 55,673*          | 6*              | 43,257*           | 0*               | 220,579          | 0               | 71,414            | 0                | 265,158          | 0               | 116,697           | 0                |
| Bitterroot      | 2,491*           | 334*            | 180*              | 0*               | 8,083*           | 297*            | 1,384*            | 0*               | 12,703           | 705             | 2,959             | 0                |
| Custer          | 1,151            | 476             | 3,334             | 0                | 3,277*           | 476*            | 1,110*            | 0*               | 819              | 196             | 8,538             | 0                |
| Deerlodge       | 160,671*         | 78*             | 2,800*            | 0*               | 458,193          | 745             | 18,733            | 0                | 539,967          | 2,935           | 19,120            | 0                |
| Flathead        | 40,148           | 62              | 16                | 0                | 19,058           | 39              | 21                | 0                | 31,567           | 398             | 111               | 0                |
| Gallatin        | 20,659*          | 42*             | 44,989*           | 0*               | 66,374*          | 5*              | 67,135*           | 0*               | 248,320          | 0               | 127,601           | 0                |
| Garnets         | 3,830*           | 21*             | 0*                | 0*               | 44,374           | 82              | 0                 | 0                | 53,578           | 2,057           | 0                 | 0                |
| Helena          | 91,317*          | 5,235*          | 526*              | 0*               | 263,767*         | 13,054*         | 4,088*            | 0*               | 459,157          | 104,332         | 12,471            | 0                |
| Kootenai        | 3,405*           | 48*             | 14*               | 4*               | 5,956*           | 116*            | 0*                | 0*               | 45,284*          | 869*            | 10*               | 18*              |
| Lewis & Clark   | 9,766*           | 3,331*          | 2,529*            | 0*               | 38,528*          | 2,248*          | 9,961*            | 0*               | 305,791          | 24,741          | 36,305            | 0                |
| Lolo            | 123,274*         | 2,562*          | 58*               | 2*               | 119,425          | 775             | 44                | 0                | 294,013*         | 12,134*         | 50*               | 4*               |
| Blackfeet IR    | ★                | ★               | ★                 | ★                | 623              | 0               | 0                 | 0                | 452              | 4               | 0                 | 0                |
| Crow IR         | 286              | 127             | 26                | 0                | ★                | ★               | ★                 | ★                | 4                | 60              | 14                | 0                |
| Flathead IR     | 26,413           | 929             | 16                | 0                | 2,371*           | 537*            | 0*                | 0*               | 15,946           | 2,044           | 160               | 0                |
| Fort Belknap IR | 28               | 104             | 0                 | 0                | 137              | 598             | 0                 | 0                | 226*             | 500*            | 0*                | 0*               |
| No. Cheyenne IR | 0                | 302             | 0                 | 0                | ★                | ★               | ★                 | ★                | 0                | 113             | 2                 | 0                |
| Rocky Boy's IR  | 1,281            | 75              | 0                 | 0                | 1,718            | 190             | 0                 | 0                | 4,231            | 437             | 0                 | 0                |
| Glacier NP      | 10,028*          | 4*              | 24*               | 0*               | 20,786           | 2               | 0                 | 0                | 4,434*           | 2*              | 6*                | 0*               |
| Yellowstone NP  | 28,085           | 0               | 36,838            | 0                | 25,535           | 2               | 29,950            | 0                | 58,393           | 462             | 41,001            | 0                |
|                 |                  |                 |                   |                  |                  |                 |                   |                  |                  |                 |                   |                  |
| <b>TOTAL</b>    | 578,506          | 13,736          | 134,607           | 6                | 1,298,784        | 19,166          | 203,840           | 0                | 2,340,043        | 151,989         | 365,045           | 22               |

<sup>1</sup>LPP = Lodgepole pine; PP = Ponderosa pine; 5-NP = 5-needle pines (WBP & LP); WWP = Western white pine

★ = Not surveyed; \* = Partially surveyed; Yellowstone NP includes MT, ID, and WY acres

**Table 7. Acres with Additional Bark Beetle-Caused Mortality on All Ownerships  
in Montana, From 2007 Through 2009**

| Reporting Area  | Engelmann Spruce |      |      | Fir Engraver Beetle |      |        | Pine Engraver Beetle |        |      | Subalpine fir Mortality |         |        | Western Pine Beetle |      |      |
|-----------------|------------------|------|------|---------------------|------|--------|----------------------|--------|------|-------------------------|---------|--------|---------------------|------|------|
|                 | 2007             | 2008 | 2009 | 2007                | 2008 | 2009   | 2007                 | 2008   | 2009 | 2007                    | 2008    | 2009   | 2007                | 2008 | 2009 |
| Beaverhead      | 0*               | 0    | 6    | 0*                  | 0    | 0      | 0*                   | 0      | 0    | 21862*                  | 464     | 554    | 10*                 | 0    | 0    |
| Bitterroot      | 0*               | 4*   | 0    | 10*                 | 0*   | 18     | 0*                   | 2*     | 4    | 9547*                   | 2,838   | 5,589  | 0*                  | 24*  | 0    |
| Custer          | 0                | 0*   | 0    | 0                   | 0*   | 0      | 10                   | 431*   | 0    | 5,485                   | 2,449*  | 771    | 0                   | 0*   | 0    |
| Deerlodge       | 0*               | 2    | 0    | 0*                  | 0    | 0      | 0*                   | 0      | 0    | 120*                    | 8,400   | 2,328  | 0*                  | 2    | 2    |
| Flathead        | 12               | 24   | 32   | 1,559               | 174  | 494    | 0                    | 0      | 2    | 6,987                   | 16,393  | 13,038 | 124                 | 0    | 20   |
| Gallatin        | 0*               | 0*   | 0    | 0*                  | 0*   | 0      | 5*                   | 4*     | 0    | 33350*                  | 14,054* | 36,384 | 0*                  | 0*   | 0    |
| Garnets         | 0*               | 0    | 0    | 0*                  | 31   | 0      | 33*                  | 0      | 137  | 334*                    | 9       | 0      | 0*                  | 2    | 67   |
| Helena          | 0*               | 4*   | 0    | 0*                  | 0*   | 0      | 0*                   | 9,995* | 0    | 6*                      | 16*     | 20     | 0*                  | 0*   | 2    |
| Kootenai        | 5*               | 10*  | 8*   | 66*                 | 36*  | 333*   | 0*                   | 12*    | 6*   | 6732*                   | 394*    | 6,139* | 0*                  | 4*   | 128* |
| Lewis & Clark   | 0*               | 2*   | 13   | 0*                  | 0*   | 0      | 40*                  | 176*   | 0    | 1022*                   | 1,341*  | 4,173  | 2*                  | 0*   | 0    |
| Lolo            | 8*               | 0    | 2*   | 18*                 | 48   | 5,898* | 470*                 | 45     | 616* | 457*                    | 2,261   | 1529*  | 265*                | 107  | 575* |
| Blackfeet IR    | ★                | 0    | 2    | ★                   | 4    | 801    | ★                    | 0      | 0    | ★                       | 2       | 107    | ★                   | 0    | 0    |
| Crow IR         | 0                | ★    | 0    | 0                   | ★    | 0      | 0                    | ★      | 0    | 4,736                   | ★       | 4      | 0                   | ★    | 0    |
| Flathead IR     | 0                | 0*   | 0    | 2                   | 2*   | 11     | 610                  | 4*     | 40   | 1,045                   | 99*     | 2,017  | 124                 | 28*  | 20   |
| Fort Belknap IR | 0                | 0    | 0*   | 0                   | 0    | 0*     | 19                   | 65     | 0*   | 0                       | 0       | 0*     | 0                   | 0    | 0*   |
| No. Cheyenne IR | 0                | ★    | 0    | 0                   | ★    | 0      | 0                    | ★      | 0    | 0                       | ★       | 0      | 0                   | ★    | 0    |
| Rocky Boy's IR  | 0                | 0    | 0    | 0                   | 0    | 0      | 0                    | 44     | 0    | 0                       | 0       | 0      | 0                   | 0    | 2    |
| Glacier NP      | 4*               | 4    | 4*   | 2*                  | 64   | 1,632* | 0*                   | 0      | 0*   | 1580*                   | 3,624   | 7,509* | 0*                  | 0    | 0*   |
| Yellowstone NP  | 374              | 4    | 18   | 0                   | 0    | 0      | 0                    | 0      | 0    | 21,987                  | 850     | 2,343  | 0                   | 0    | 0    |
| <b>TOTAL</b>    | 403              | 54   | 85   | 1,657               | 359  | 9,187  | 1,187                | 10,778 | 805  | 115,250                 | 53,194  | 82,505 | 401                 | 182  | 816  |

★ = Not surveyed    \* = Partially surveyed

Yellowstone NP includes MT, ID and WY acres