

CARIBOU-TARGHEE NATIONAL FOREST

Westside Ranger District

2014 Aerial Insect and Disease Detection Survey




Disclaimer
 Insect and disease data should be used only as an indicator of insect and disease activity, and should be ground-truthed for actual location and causal agent. Polygons indicate locations of tree mortality, defoliation, and/or other damage. Intensity of damage is variable, and not all trees and areas indicated are dead or damaged. The joint cooperators reserve the right to correct, modify, update, or replace the data as necessary. Using this data for purposes other than those for which it was intended may yield inaccurate or misleading results. The topographic map image is included to provide a visual point of reference.

Aerial Insect & Disease Detection Surveys

Aerial insect and disease detection surveys are conducted annually to detect and monitor annual, visible, vegetation damage primarily caused by insects. Aerial detection surveys are intended to detect new activity, to monitor the trend of ongoing activity, to provide general location information, and to subjectively rate levels of defoliation. These flights are conducted in a joint partnership between the USDA Forest Service, Idaho Department of Lands, and The Nevada Division of Forestry.

Data represented on this map are based on trees visibly affected by forest insects, as detected by aerial observers. Most bark beetle-killed trees are not typically symptomatic (leaved foliage that is yellow, orange, or brown) until nearly a year following beetle attack. Therefore, the numbers of trees killed by bark beetles, as indicated on this map, are a reflection of last year's mortality. The numbers do not reflect the current year's beetle population or number of currently attacked trees.

Observers have just a few seconds to recognize, identify, and document observed activity. Air turbulence, cloud shadow, haze, smoke, and observer experience can all affect the quality of the survey.



INSECT & DISEASE ACTIVITY

BARK BEETLES

Species	Host	Mortality
Mountain Pine Beetle (MPB)	Lodgepole pine	1-4
Douglas-fir Beetle (DFB)	Douglas-fir	5-14
Spruce Beetle (SB)	Spruce	15+ Trees
Ips confusus	Pinyon pine	

MORTALITY & DISEASE

Subalpine Fir Mortality	Subalpine fir	0/20
Decline - Aspen	Stand decline	Light <50% Heavy >50%

DEFOLIATORS

Western Spruce Budworm - Douglas-fir, True Fir	Defoliation	Light ~ 50% Heavy > 50%
Unknown defoliation - Asp, Oak, DF, Willow	Not Rated	Light, Heavy

ADDITIONAL SYMBOLS AND DAMAGE AGENTS

# - Trees affected	Survey	Boundary
HA - Trees/Acre affected	Area	
Aspen	Asp	Ponderosa pine
Douglas-fir	DF	Spruce
Larch	L	Subalpine fir
Lodgepole pine	LP	Whitebark pine