


# 2010 Aerial Insect and Disease Survey Farson, Wyoming USGS 100K TOPO!: 42109-A1




1:100,000

**Legend**

Use of the Number System  
Example: 5-25 = The first number before the dash is the causal agent code. The number after the dash is the number of dead "ladder" trees in the polygon or point. When recent dead trees are not counted, an intensity code of L-Light, M-Moderate, and H-High may be used after the causal agent code. Periodically, trees per acre estimates are used after the causal agent code instead of number of dead "ladder" trees or an intensity code. For example: 5-10A = The first number before the dash is the causal agent code. The number after the dash is an estimation of the number of dead "ladder" trees in the polygon per acre. In this case it would be an estimation that, on the average, one tree per every two acres would be dead "ladder" trees. In another example: 5-3A = that on the average, an estimated three trees per acre are dead "ladder" trees. A "T" is used as a separator when a point/polygon has more than one causal agent code.

Code	Causal Agent	Primary Host	Code	Causal Agent	Primary Host	Code	Causal Agent	Primary Host
1	Douglas-fir beetle	Douglas-fir	60	Arthropods	Lodgepole Pine	100	fox squirrel ragging	Cottonwood/Poplar
2	Engelmann spruce beetle	Engelmann Spruce	61	White pine blister rust	Sitka Spruce	101	fall webworm	Cottonwood/Poplar
3	Mountain pine beetle	Ponderosa Pine	62	Dwarf mistlebe	Softwoods	102	road salt	Softwoods
4	Mountain pine beetle	Lodgepole Pine	63	Yucca dieback	Ponderosa Pine	103	pinewood nematode	Scots Pine
5	Mountain pine beetle	5-Needle Pine	64	Indus 605, 60 & 68	All Tree Species	110	oak wilt	Oak
6	Mountain pine beetle	Ponderosa Pine	65	Air pollutants	All Tree Species	111	foliar disease	All Tree Species
7	Western pine beetle	White Fir	66	Chemical damage	All Tree Species	112	spurge ice	White Spruce
8	Western pine beetle	Douglas-fir	67	Lophodermium pinastri	Softwoods	113	twined chestnut borer	Oak
9	Western pine beetle	Subalpine Fir	68	Rhabdocline pseudotsugae	Douglas-fir	114	anthracnose like leaf disease	Bir Oak
10	Douglas-fir engraver beetle	Softwoods	69	Lophodermium arcuta	Softwoods	115	Dieback	All Tree Species
11	Western balsam bark beetle	Softwoods	70	Lecanosticella aculeata	Softwoods	116	Mortality	All Tree Species
12	Unidentified bark beetle	Ponderosa Pine	71	Lophodermium concolor	Softwoods	117	Discoloration	All Tree Species
13	Pine engraver	Lodgepole Pine	72	Dothiorella per	Softwoods	118	Herbicide	All Tree Species
14	Pine engraver	Ponderosa Pine	73	Needle cast (Hymenoptera)	Softwoods	119	Flagging	All Tree Species
15	Ponderosa pine needle miner	Lodgepole Pine	74	Root Rot	All Tree Species	120	aspen tortrix	Quaking Aspen
16	Jack pine budworm	Jack Pine	75	Unidentified disease	Softwoods	121	Mammalian blight	Quaking Aspen
17	Jack pine budworm	Douglas-fir	76	Winter damage light	All Tree Species	200	Dieback (ash)	Ash
18	Sproue budworm, medium defol.	Douglas-fir	77	Winter damage medium	All Tree Species	201	Dieback (cottonwood)	Cottonwood/Poplar
19	Sproue budworm, heavy defol.	Douglas-fir	78	Winter damage heavy	All Tree Species	202	Dieback (hardwood)	Hardwoods
20	Douglas-fir sawfly	Douglas-fir	79	Winter damage heavy	All Tree Species	203	Dieback (oak)	Oak
21	Pine butterfly	Ponderosa Pine	80	Diabrotica	Softwoods	204	Dieback (oak)	Oak
22	Pine looper	Ponderosa Pine	81	Prionus bark stain	Common Pinon	210	Mortality (oak cottonwood)	Cottonwood/Poplar
23	Pine tortrix	Ponderosa Pine	82	Fire	All Tree Species	211	Mortality (eastern cedar)	Eastern Red Cedar
24	Teat caterpillars	Hardwoods	83	Windthrow	Softwoods	212	Mortality (hardwood)	Hardwoods
25	Leaf beetles	Hardwoods	84	High water damage	All Tree Species	213	Mortality (oak)	Oak
26	Oak leaf roller	Hardwoods	85	Arthropods	All Tree Species	214	Mortality (spruce)	Spruce
27	Pine needle-shaft miner	Ponderosa Pine	86	Quaking Aspen	Quaking Aspen	220	Discoloration (ash)	Softwoods
28	Pine bark beetle	Ponderosa Pine	87	Unidentified defolator	Common Pinon	221	Discoloration (cottonwood)	Cottonwood/Poplar
29	Leaf beetles	Hardwoods	88	Juniper mortality-unknown agents)	Juniper	222	Discoloration (eastern cedar)	Eastern Red Cedar
30	Leaf beetles	Hardwoods	89	Quaking Aspen	Quaking Aspen	223	Discoloration (hardwood)	Hardwoods
31	Leaf beetles	Hardwoods	90	Limber Pine	Limber Pine	224	Discoloration (oak)	Oak
32	Leaf beetles	Hardwoods	91	Limber Pine	Limber Pine	225	Discoloration (spruce)	Spruce
33	Leaf beetles	Hardwoods	92	Limber Pine	Limber Pine	230	Herbicide (oak)	Eastern Red Cedar
34	Leaf beetles	Hardwoods	93	Limber Pine	Limber Pine	231	Herbicide (eastern cedar)	Eastern Red Cedar
35	Leaf beetles	Hardwoods	94	Limber Pine	Limber Pine	232	Flagging (hardwood)	Hardwoods
36	Leaf beetles	Hardwoods	95	Limber Pine	Limber Pine	233	Flagging (hardwood)	Hardwoods
37	Leaf beetles	Hardwoods	96	Limber Pine	Limber Pine	234	Unidentified defolator (cottonwood)	Cottonwood/Poplar
38	Leaf beetles	Hardwoods	97	Limber Pine	Limber Pine	251	Unidentified defolator (oak)	Oak
39	Leaf beetles	Hardwoods	98	Limber Pine	Limber Pine	252	Unidentified defolator (hardwood)	Hardwoods
40	Leaf beetles	Hardwoods	99	Limber Pine	Limber Pine	300	Mortality (pine)	Pine
41	Leaf beetles	Hardwoods	100	Limber Pine	Limber Pine			
42	Leaf beetles	Hardwoods	101	Limber Pine	Limber Pine			
43	Leaf beetles	Hardwoods	102	Limber Pine	Limber Pine			
44	Leaf beetles	Hardwoods	103	Limber Pine	Limber Pine			
45	Leaf beetles	Hardwoods	104	Limber Pine	Limber Pine			
46	Leaf beetles	Hardwoods	105	Limber Pine	Limber Pine			
47	Leaf beetles	Hardwoods	106	Limber Pine	Limber Pine			
48	Leaf beetles	Hardwoods	107	Limber Pine	Limber Pine			



**How Aerial Surveys Are Conducted**

Data represented on this map are based on aerial observations manually recorded onto a map. This procedure is considered both an art form and a form of scientific data collection, and is highly subjective. An observer only has a few seconds to recognize the color difference between healthy and damaged trees of different species; diagnose causal agents correctly; estimate intensity; delineate the extent of damage; and precisely record this information on a georeferenced map. Air turbulence, cloud shadows, distance from aircraft, haze, smoke, and observer experience can all affect the quality of the survey. These data summaries provide an estimate of conditions on the ground and may differ from estimates derived by other methods.

Aerial surveys provide information on the current status for many causal agents, and are important when examining insect activity trends by comparing historical and current survey data over large areas.

Overview surveys are a 'snap shot' in time and therefore may not be timed to accurately capture the true extent or severity of a particular disturbance activity. Aerial surveys can be thought of as the first stage in a multi-stage sampling design. Other remote sensing approaches, including aerial photography, electro-optical sensors, and specially designed aerial surveys with modified flight patterns, can be used to more accurately delineate the extent and severity of a particular disturbance agent. The preceding methods are often more costly than overview surveys, and are generally reserved to address situations of sufficient environmental, economic, or political importance.

**Map Created November 4 2010**  
**Projection: UTM NAD83 Zone 13**  
**Author: J. Ross, USDA Forest Service**

A data dictionary and digital copies of this map and the insect and disease data are available at: <http://www.fs.fed.us/r2/resources/ihm/aerialsurveys/>

**DIRECT ALL INQUIRIES TO:**

**Wyoming State Forestry Division**  
 1100 West 22nd Street  
 Cheyenne, Wyoming 82002

**USDA Forest Service, Region 2**  
 Renewable Resources  
 Forest Health Management  
 PO Box 25127  
 Lakewood, Colorado 80225

\*\*\*\*DISCLAIMER\*\*\*\*  
 Forest Health Protection (FHP) and its partners strive to maintain an accurate Aerial Detection Survey (ADS) Dataset, but due to the conditions under which the data are collected, FHP and its partners shall not be held responsible for missing or inaccurate data. ADS are not intended to replace more specific information. An accuracy assessment has not been done for this dataset; however, ground checks are completed in accordance with local and national guidelines <http://www.fs.fed.us/foresthealth/aviationqualityassurance.shtml>. Maps and data may be updated without notice. Please cite "USDA Forest Service, Forest Health Protection and its partners" as the source of this data in maps and publications.

Due to the nature of aerial surveys, the data on this map will only provide rough estimates of location, intensity and the resulting trend information for agents detectable from the air. Many of the most destructive diseases are not represented on this map because these agents are not detectable from aerial surveys. The data presented on this map should only be used as a partial indicator of insect and disease activity, and should be validated on the ground for actual location and causal agent. Shaded areas show locations where tree mortality or defoliation were apparent from the air. Intensity of damage is variable and not all trees in shaded areas are dead or defoliated.

The insect and disease data represented on this map are available digitally from the USDA Forest Service, Region Two Forest Health Management group. The cooperators reserve the right to correct, update, modify or replace GIS products. Using this map for purposes other than those for which it was intended may yield inaccurate or misleading results.