

**Appendix F
Disposition of
1986 Forest Plan
Management Indicator
Species**

Appendix F – Disposition of 1986 Management Indicator Species

Management Indicator Species	Representative Habitat	Disposition
Robbins' cinquefoil	Alpine	Habitat tracked by 2 alpine community Ecological Indicators (EI); Species effects presented in Rare and Unique Features section of EIS and in Biological Evaluation (BE)
Canada lynx	Low Disturbance Forested	Habitat tracked by high elevation spruce-fir bird suite EI; Species effects discussed in Rare and Unique Features section of EIS and in BA
Blackpoll warbler	High Disturbance Forested; young or resembling young	Habitat tracked by high elevation spruce-fir bird suite EI, which includes Blackpoll warbler
Gray-cheeked (now Bicknell's) thrush	High Disturbance Forested; mature, overmature or old growth	Habitat tracked by high elevation spruce-fir bird suite EI, which includes Bicknell's thrush; Species effects discussed in Rare and Unique Features section of EIS and in BA
Northern goshawk	Mature/overmature hardwoods and mixed woods	Habitat tracked by scarlet tanager MIS
Chestnut-sided warbler	Regenerating hardwoods and mixed woods	Maintained as MIS for regenerating/young hardwoods
Broad-winged hawk	Mature/overmature paper birch and aspen	Habitat tracked by ruffed grouse MIS; Habitat effects also discussed in Vegetation and Habitats section of EIS
Ruffed grouse	All ages of aspen and regenerating/young birch	Maintained as MIS
Gray squirrel	Mature/overmature oak and oak-pine	Habitat tracked in Vegetation and Habitats section of EIS
Rufous-sided (now Eastern) towhee	Regenerating/young oak and oak-pine	Habitat tracked in Vegetation and Habitats and Rare and Unique Features sections of EIS
Pine warbler	Mature/overmature pine	Habitat tracked in Vegetation and Habitats and Rare and Unique Features sections of EIS
Northern (now Dark-eyed) junco	Regenerating/young pine	Habitat tracked in Vegetation and Habitats and Rare and Unique Features sections of EIS
White-tailed deer	All ages of hemlock	Habitat tracked in Vegetation and Habitats section of EIS; Deer evaluated indirectly in BA
Cape May warbler	Mature/overmature spruce and fir	Mature spruce and fir category retained, but blackburnian warbler used instead as MIS

Management Indicator Species	Representative Habitat	Disposition
Snowshoe hare	Regenerating/young spruce and fir	Habitat type retained, but tracked instead by magnolia warbler MIS
Eastern kingbird and Eastern bluebird	Upland openings — grass, forb, or orchard	Opening habitat tracked in the Vegetation and Habitats section of EIS
Mourning warbler	Upland openings — shrub	Opening habitat tracked in Vegetation and Habitats and Rare and Unique Features sections of EIS
Black duck	Wetlands	Habitat effects discussed in Rare and Unique Features and Watershed sections of EIS
Brook trout	Permanent lakes, ponds, and streams	Habitat effects discussed in Riparian and Aquatics, Watershed, and Rare and Unique Features sections of EIS
Peregrine falcon	Cliffs and talus	Retained as an EI; Additional effects also discussed in BA
Pine (now American) marten	State threatened species	Marten dropped as MIS because it did not represent a particular habitat type or condition; however, now used to evaluate fragmentation and landscape corridor conditions in the Wildlife section, Chapter 3
Osprey	State threatened species	Osprey dropped as MIS because it did not represent a particular habitat type or condition; wetland and pond habitats are evaluated in the Watershed and Riparian and Aquatics effects section, Chapter 3
Common loon	State threatened species	Common loon dropped as MIS because it did not represent a particular habitat type or condition; however, it is specifically tracked as a sensitive species in the BA.
Sunapee trout	State endangered	Sunapee trout dropped as MIS because it did not represent a particular habitat type or condition; it is no longer believed to be present on the Forest.