



2010 Aerial Insect and Disease Survey Newcastle, Wyoming USGS 100K TOPO! 43104-E1



1:100,000

Legend		Causal Agent(s)	Not Flown
<p>Use of the Number System Example: 5-25 = The first number before the dash is the causal agent code. The number after the dash is the number of dead "faded" trees in the polygon or point. When recent dead trees are not counted, an intensity code of L-light, M-medium, and H-high may be used after the causal agent code. Periodically, trees per acre estimates are used after the causal agent code instead of number of dead "faded" trees (or an intensity code). For example: 5-12A = The first number before the dash is the causal agent code. The number after the dash is an estimation of the number of dead "faded" trees in the polygon per acre. In this case it would be an estimation that, on the average, one tree per every two acres would be a dead "faded" tree. In another example: 5-3A = that on the average, an estimated three trees per acre are dead "faded" trees. A "P" is used as a separator when a point/polygon has more than one causal agent code.</p>			
Code	Causal Agent	Primary Host	Code
1	Douglas fir beetle	Douglas fir	100
2	Engelmann spruce beetle	Engelmann spruce	101
3	Mountain pine beetle	Ponderosa pine	102
4	Mountain pine beetle	Lodgepole pine	103
5	Western pine beetle	White fir	104
6	Western pine beetle	White fir	105
7	Western pine beetle	White fir	106
8	Western pine beetle	White fir	107
9	Western pine beetle	White fir	108
10	Western pine beetle	White fir	109
11	Western pine beetle	White fir	110
12	Western pine beetle	White fir	111
13	Western pine beetle	White fir	112
14	Western pine beetle	White fir	113
15	Western pine beetle	White fir	114
16	Western pine beetle	White fir	115
17	Western pine beetle	White fir	116
18	Western pine beetle	White fir	117
19	Western pine beetle	White fir	118
20	Western pine beetle	White fir	119
21	Western pine beetle	White fir	120
22	Western pine beetle	White fir	121
23	Western pine beetle	White fir	122
24	Western pine beetle	White fir	123
25	Western pine beetle	White fir	124
26	Western pine beetle	White fir	125
27	Western pine beetle	White fir	126
28	Western pine beetle	White fir	127
29	Western pine beetle	White fir	128
30	Western pine beetle	White fir	129
31	Western pine beetle	White fir	130
32	Western pine beetle	White fir	131
33	Western pine beetle	White fir	132
34	Western pine beetle	White fir	133
35	Western pine beetle	White fir	134
36	Western pine beetle	White fir	135
37	Western pine beetle	White fir	136
38	Western pine beetle	White fir	137
39	Western pine beetle	White fir	138
40	Western pine beetle	White fir	139
41	Western pine beetle	White fir	140
42	Western pine beetle	White fir	141
43	Western pine beetle	White fir	142
44	Western pine beetle	White fir	143
45	Western pine beetle	White fir	144
46	Western pine beetle	White fir	145
47	Western pine beetle	White fir	146
48	Western pine beetle	White fir	147



How Aerial Surveys Are Conducted

Data represented on this map are based on aerial observations manually recorded onto a map. This procedure is considered both an art form and a form of scientific data collection, and is highly subjective. An observer only has a few seconds to recognize the color difference between healthy and damaged trees of different species; diagnose causal agents correctly; estimate intensity; delineate the extent of damage; and precisely record this information on a georeferenced map. Air turbulence, cloud shadows, distance from aircraft, haze, smoke, and observer experience can all affect the quality of the survey. These data summaries provide an estimate of conditions on the ground and may differ from estimates derived by other methods.

Overview surveys are a 'snap shot' in time and therefore may not be timed to accurately capture the true extent or severity of a particular disturbance activity. Aerial surveys can be thought of as the first stage in a multi-stage sampling design. Other remote sensing approaches, including aerial photography, electro-optical sensors, and specially designed aerial surveys with modified flight patterns, can be used to more accurately delineate the extent and severity of a particular disturbance agent. The preceding methods are often more costly than overview surveys, and are generally reserved to address situations of sufficient environmental, economic, or political importance.

Map Created November 1 2010
Projection: UTM NAD83 Zone 13
Author: J. Ross, USDA Forest Service

A data dictionary and digital copies of this map and the insect and disease data are available at: <http://www.fs.fed.us/r2/resources/hm/aerialsurvey/>

DIRECT ALL INQUIRIES TO:

Wyoming State Forestry Division
 1100 West 22nd Street
 Cheyenne, Wyoming 82002

USDA Forest Service, Region 2
 Renewable Resources
 Forest Health Management
 PO Box 25127
 Lakewood, Colorado 80225

*******DISCLAIMER*******

Forest Health Protection (FHP) and its partners strive to maintain an accurate Aerial Detection Survey (ADS) Dataset, but due to the conditions under which the data are collected, FHP and its partners shall not be held responsible for missing or inaccurate data. ADS are not intended to replace more specific information. An accuracy assessment has not been done for this dataset; however, ground checks are completed in accordance with local and national guidelines <http://www.fs.fed.us/foresthealth/assessment/qualityassurance.shtml>. Maps and data may be updated without notice. Please cite "USDA Forest Service, Forest Health Protection and its partners" as the source of this data in maps and publications.

Due to the nature of aerial surveys, the data on this map will only provide rough estimates of location, intensity and the resulting trend information for agents detectable from the air. Many of the most destructive diseases are not represented on this map because these agents are not detectable from aerial surveys. The data presented on this map should only be used as a partial indicator of insect and disease activity, and should be validated on the ground for actual location and causal agent. Shaded areas show locations where tree mortality or defoliation were apparent from the air. Intensity of damage is variable and not all trees in shaded areas are dead or defoliated.

The insect and disease data represented on this map are available digitally from the USDA Forest Service, Region Two Forest Health Management group. The cooperators reserve the right to correct, update, modify or replace GIS products. Using this map for purposes other than those for which it was intended may yield inaccurate or misleading results.