

## R2 Individual Species Recommendations

<b>TAXON</b>	<b>Scientific Name</b>			<b>Common Name</b>		
PD	ERIGERON RADICATUS			TAPROOTED FLEABANE		
<b>FWS</b>	<b>FS</b>	<b>BLM</b>				
<b>Grank</b>	<b>CO S Rank</b>	<b>WY S Rank</b>	<b>SD S Rank</b>	<b>NE S Rank</b>	<b>KS S Rank</b>	
G3 N3		S2				
	<b>CO Status</b>	<b>WY Status</b>	<b>SD Status</b>	<b>NE Status</b>	<b>KS Status</b>	
<b>IUCN</b>	<b>PIF – Area Importance</b>		<b>PIF – Population Trend</b>			
<b>FS Experts assigned Species Evaluation</b>			<b>Non-NFS Experts assigned Species Evaluations</b>			
			Scott Laursen, Bonnie Heidel, and Joy Hanley.			
<b>Sensitive Species Recommendation</b> <span style="float: right;">(Please check 1 box below)</span>						
<input type="checkbox"/>	R2 Sensitive Species (SS)	<input checked="" type="checkbox"/>	Not R2 SS, but should be considered for other Emphasis Species Lists			
<input type="checkbox"/>	Not R2 SS and Not of Concern Now	<input type="checkbox"/>	Insufficient Information Available to make a recommendation			
<b>Recommendation Rationale</b>						
<p><b>In Region 2 only known from the Shoshone NF (8 known occurrences with no population numbers in evaluation). Threats to this species are low due to the rugged alpine habitat it occupies above timberline in krummholz (low confidence rank) and no sheep grazing occurring in its habitat on the Shoshone NF according to Shoshone NF Botanist. Populations are presumed to be stable (high confidence rank). Recreational livestock may be a threat to this species on the Shoshone NF.</b></p>						
<p><b>John Proctor and Gay Austin</b></p>						
<b>Name/Title</b>			<b>Date</b>			