

## R2 Individual Species Recommendations

<b>TAXON</b>	<b>Scientific Name</b>		<b>Common Name</b>		
PD	Viola renifolia var. brainerdii		Kidney-leaf white violet		
<b>FWS</b>	<b>FS</b>	<b>BLM</b>			
<b>Grank</b>	<b>CO S Rank</b>	<b>WY S Rank</b>	<b>SD S Rank</b>	<b>NE S Rank</b>	<b>KS S Rank</b>
G5T5		S1	S4		
	<b>CO Status</b>	<b>WY Status</b>	<b>SD Status</b>	<b>NE Status</b>	<b>KS Status</b>
<b>IUCN</b>	<b>PIF – Area Importance</b>		<b>PIF – Population Trend</b>		
<b>FS Experts assigned Species Evaluation</b>			<b>Non-NFS Experts assigned Species Evaluations</b>		
			Joy Handley, Bonnie Heidel, Scott Laursen		
<b>Sensitive Species Recommendation</b> (Please check 1 box below)					
<input type="checkbox"/>	<b>R2 Sensitive Species (SS)</b>		<b>Not R2 SS, but should be considered for other Emphasis Species Lists</b>		
<input checked="" type="checkbox"/>	<b>Not R2 SS and Not of Concern Now</b>		<b>Insufficient Information Available to make a recommendation</b>		
<b>Recommendation Rationale</b>					
<p>Plants database considers the variety as a synonym with the species. The species has a very wide distribution across Canada and the northern tier of states. Occurs in moist, shaded coniferous forest habitats. Populations could be impacted by recreation use and timber harvesting. Not of concern due to its broad distribution and relatively broad habitat associations.</p>					

<b>Name/Title</b>	Nancy Warren & Jeff Redders	<b>Date</b>	Oct. 3, 2002
-------------------	-----------------------------	-------------	--------------