


2011 Aerial Insect and Disease Survey Cody, Wyoming USGS 100K TOPO!: 44109-E1




1:100,000

Legend

Causal Agent(s) (Orange box) **Not Flown** (Grey box)

Use of the Number System
Example: 5-25 = The first number before the dash is the causal agent code. The number after the dash is the number of dead "fader" trees in the polygon or point. When recent dead trees are not counted, an intensity code of L=light, M=moderate, and H=high may be used after the causal agent code. Periodically, trees per acre estimates are used after the causal agent code instead of number of dead "fader" trees (or an intensity code). For example: 5-12A = The first number before the dash is the causal agent code. The number after the dash is an estimation of the number of dead "fader" trees in the polygon per acre. In this case it would be an estimation that, on the average, one tree per every two acres would be a dead "fader" tree. In another example: 5-3A = that on the average, an estimated three trees per acre are dead "fader" trees. A "17" is used as a separator when a point polygon has more than one causal agent code.

| Code | Causal Agent | Primary Host | Code | Causal Agent | Primary Host |
|------|--------------------------------------|------------------|------|---|-----------------------|
| 1 | Causal Agent | | 48 | Albopitca | Lodgepole Pine |
| 2 | Engelmann spruce beetle | Engelmann Spruce | 49 | Stactoloma rust | Lodgepole Pine |
| 3 | Blue spruce tip beetle | Blue Spruce | 50 | White pine sister rust | Cottwood Poplar |
| 4 | Mountain pine beetle | Ponderosa Pine | 51 | Dwarf mistletoe | Cottwood Poplar |
| 5 | Mountain pine beetle | Lodgepole Pine | 52 | Elysiodes | Scotch Pine |
| 6 | Western pine beetle | Ponderosa Pine | 53 | Includes #65, #66 & #68 | All Tree Species |
| 7 | Fire Enginer | White Fir | 54 | Air pollutants | All Tree Species |
| 8 | Douglas fir engraver beetle | Douglas Fir | 55 | Chemical damage | All Tree Species |
| 9 | Western balsam bark beetle | Subalpine Fir | 56 | Lophodermium pinastri | Softwoods |
| 10 | Unidentified bark beetle | Softwoods | 57 | Rhododendron pseudotsugae | Douglas Fir |
| 11 | Pine engraver | Lodgepole Pine | 58 | Lophodermium arcuta | Softwoods |
| 12 | Pine engraver | Lodgepole Pine | 59 | Lecanora acicola | Softwoods |
| 13 | Ponderosa pine needle miner | Lodgepole Pine | 60 | Lophodermium concolor | Softwoods |
| 14 | Ponderosa pine needle miner | Ponderosa Pine | 61 | Dothistroma pini | Softwoods |
| 15 | Jack pine budworm | Jack Pine | 62 | Needle cast (Hypodermatomyces) | Softwoods |
| 16 | Spruce budworm, light defol. | Douglas Fir | 63 | Rust Red | All Tree Species |
| 17 | Spruce budworm, medium defol. | Douglas Fir | 64 | Unidentified disease | Softwoods |
| 18 | Spruce budworm, heavy defol. | Douglas Fir | 65 | Winter damage light | All Tree Species |
| 19 | Douglas fir bark moth | Douglas Fir | 66 | Winter damage medium | All Tree Species |
| 20 | Pine butterfly | Ponderosa Pine | 67 | Winter damage heavy | All Tree Species |
| 21 | Pine looper | Ponderosa Pine | 68 | Dipodops | Softwoods |
| 22 | Pine tortrix | Ponderosa Pine | 69 | Pinon black stain | Common Pinon |
| 23 | Text caterpillars | Hardwoods | 70 | Fire | All Tree Species |
| 24 | Leaf beetles | Hardwoods | 71 | Porcupine | Softwoods |
| 25 | Aspen defoliation | Quaking Aspen | 72 | Windthrow | All Tree Species |
| 26 | Oak leaf roller | Hardwoods | 73 | High water damage | All Tree Species |
| 27 | Pine needle-shaft miner | Ponderosa Pine | 74 | Avalanche | Quaking Aspen |
| 28 | Pine sawflies | Ponderosa Pine | 75 | Aspen decline-multiple agents) | All Tree Species |
| 29 | Vanilla oak leaf caterpillar | Hardwoods | 76 | Pinon pine mortality | Common Pinon |
| 30 | Unidentified defoliator | All Tree Species | 77 | Juniper mortality (unknown agents) | Juniper |
| 31 | Cottwood Decline/Mortality | Cottwood | 78 | Camelid oak decline-unknown agents) | Camelid Oak |
| 32 | Heterobasium annosum (Fomes annosus) | Softwoods | 79 | Liner tree decline-multiple agents) | Liner Tree |
| 33 | Amelaria setyae (Amelaria melles) | Softwoods | 80 | Hail damage | All Tree Species |
| 34 | Thromopsis | Softwoods | 81 | Unknown polygon | Unknown |
| 35 | Cystospora | All Tree Species | 82 | Unknown Pinon | Common Pinon |
| 36 | Western gall rust | Unknown | 83 | 100 yd pinon mosaic | Lodgepole Pine |
| 37 | Domatidra rust | Unknown | 84 | 101 rust salt tip | Lodgepole Pine |
| 38 | | | 85 | 102 insect etia disease | Elm |
| 39 | | | 86 | 103 dipodops blight | Ponderosa Pine |
| 40 | | | 87 | 104 drought killed narrow leaf cottonwood | Narrowleaf Cottonwood |
| 41 | | | 88 | | |
| 42 | | | 89 | | |
| 43 | | | 90 | | |
| 44 | | | 91 | | |
| 45 | | | 92 | | |
| 46 | | | 93 | | |
| 47 | | | 94 | | |
| 48 | | | 95 | | |
| 49 | | | 96 | | |
| 50 | | | 97 | | |
| 51 | | | 98 | | |
| 52 | | | 99 | | |
| 53 | | | 100 | | |
| 54 | | | 101 | | |
| 55 | | | 102 | | |
| 56 | | | 103 | | |
| 57 | | | 104 | | |
| 58 | | | 105 | | |
| 59 | | | 106 | | |
| 60 | | | 107 | | |
| 61 | | | 108 | | |
| 62 | | | 109 | | |
| 63 | | | 110 | | |
| 64 | | | 111 | | |
| 65 | | | 112 | | |
| 66 | | | 113 | | |
| 67 | | | 114 | | |
| 68 | | | 115 | | |
| 69 | | | 116 | | |
| 70 | | | 117 | | |
| 71 | | | 118 | | |
| 72 | | | 119 | | |
| 73 | | | 120 | | |
| 74 | | | 121 | | |
| 75 | | | 122 | | |
| 76 | | | 123 | | |
| 77 | | | 124 | | |
| 78 | | | 125 | | |
| 79 | | | 126 | | |
| 80 | | | 127 | | |
| 81 | | | 128 | | |
| 82 | | | 129 | | |
| 83 | | | 130 | | |
| 84 | | | 131 | | |
| 85 | | | 132 | | |
| 86 | | | 133 | | |
| 87 | | | 134 | | |
| 88 | | | 135 | | |
| 89 | | | 136 | | |
| 90 | | | 137 | | |
| 91 | | | 138 | | |
| 92 | | | 139 | | |
| 93 | | | 140 | | |
| 94 | | | 141 | | |
| 95 | | | 142 | | |
| 96 | | | 143 | | |
| 97 | | | 144 | | |
| 98 | | | 145 | | |
| 99 | | | 146 | | |
| 100 | | | 147 | | |
| 101 | | | 148 | | |
| 102 | | | 149 | | |
| 103 | | | 150 | | |
| 104 | | | 151 | | |
| 105 | | | 152 | | |
| 106 | | | 153 | | |
| 107 | | | 154 | | |
| 108 | | | 155 | | |
| 109 | | | 156 | | |
| 110 | | | 157 | | |
| 111 | | | 158 | | |
| 112 | | | 159 | | |
| 113 | | | 160 | | |
| 114 | | | 161 | | |
| 115 | | | 162 | | |
| 116 | | | 163 | | |
| 117 | | | 164 | | |
| 118 | | | 165 | | |
| 119 | | | 166 | | |
| 120 | | | 167 | | |
| 121 | | | 168 | | |
| 122 | | | 169 | | |
| 123 | | | 170 | | |
| 124 | | | 171 | | |
| 125 | | | 172 | | |
| 126 | | | 173 | | |
| 127 | | | 174 | | |
| 128 | | | 175 | | |
| 129 | | | 176 | | |
| 130 | | | 177 | | |
| 131 | | | 178 | | |
| 132 | | | 179 | | |
| 133 | | | 180 | | |
| 134 | | | 181 | | |
| 135 | | | 182 | | |
| 136 | | | 183 | | |
| 137 | | | 184 | | |
| 138 | | | 185 | | |
| 139 | | | 186 | | |
| 140 | | | 187 | | |
| 141 | | | 188 | | |
| 142 | | | 189 | | |
| 143 | | | 190 | | |
| 144 | | | 191 | | |
| 145 | | | 192 | | |
| 146 | | | 193 | | |
| 147 | | | 194 | | |
| 148 | | | 195 | | |
| 149 | | | 196 | | |
| 150 | | | 197 | | |
| 151 | | | 198 | | |
| 152 | | | 199 | | |
| 153 | | | 200 | | |



How Aerial Surveys Are Conducted

Data represented on this map are based on aerial observations manually recorded onto a map. This procedure is considered both an art form and a form of scientific data collection, and is highly subjective. An observer only has a few seconds to recognize the color difference between healthy and damaged trees of different species; diagnose causal agents correctly; estimate intensity; delineate the extent of damage; and precisely record this information on a georeferenced map. Air turbulence, cloud shadows, distance from aircraft, haze, smoke, and observer experience can all affect the quality of the survey. These data summaries provide an estimate of conditions on the ground and may differ from estimates derived by other methods.

Aerial surveys provide information on the current status for many causal agents, and are important when examining insect activity trends by comparing historical and current survey data over large areas.

Overview surveys are a "snap shot" in time and therefore may not be timed to accurately capture the true extent or severity of a particular disturbance activity. Aerial surveys can be thought of as the first stage in a multi-stage sampling design. Other remote sensing approaches, including aerial photography, electro-optical sensors, and specially designed aerial surveys with modified flight patterns, can be used to more accurately delineate the extent and severity of a particular disturbance agent. The preceding methods are often more costly than overview surveys, and are generally reserved to address situations of sufficient environmental, economic, or political importance.

Map Created December 1 2011
Projection: UTM NAD83 Zone 13
Author: J. Ross, USDA Forest Service

A data dictionary and digital copies of this map and the insect and disease data are available at: <http://www.fs.fed.us/r2/resources/fhm/aerialsurvey/>

DIRECT ALL INQUIRIES TO:

Wyoming State Forestry Division
 1100 West 22nd Street
 Cheyenne, Wyoming 82002

USDA Forest Service, Region 2
 Renewable Resources
 Forest Health Management
 PO Box 25127
 Lakewood, Colorado 80225

*****DISCLAIMER*****
 Forest Health Protection (FHP) and its partners strive to maintain an accurate Aerial Detection Survey (ADS) Dataset, but due to the conditions under which the data are collected, FHP and its partners shall not be held responsible for missing or inaccurate data. ADS are not intended to replace more specific information. An accuracy assessment has not been done for this dataset; however, ground checks are completed in accordance with local and national guidelines <http://www.fs.fed.us/foresthealth/aviation/qualityassurance.shtml>. Maps and data may be updated without notice. Please cite "USDA Forest Service, Forest Health Protection and its partners" as the source of this data in maps and publications.

Due to the nature of aerial surveys, the data on this map will only provide rough estimates of location, intensity and the resulting trend information for agents detectable from the air. Many of the most destructive diseases are not represented on this map because these agents are not detectable from aerial surveys. The data presented on this map should only be used as a partial indicator of insect and disease activity, and should be validated on the ground for actual location and causal agent. Shaded areas show locations where tree mortality or defoliation were apparent from the air. Intensity of damage is variable and not all trees in shaded areas are dead or defoliated.

The insect and disease data represented on this map are available digitally from the USDA Forest Service, Region Two Forest Health Management group. The cooperators reserve the right to correct, update, modify or replace GIS products. Using this map for purposes other than those for which it was intended may yield inaccurate or misleading results.