

**Aquatic Species Biological Assessment for the
Seneca, Deadhorse, Hanscomb, and Fields Peak Allotments**

BLUE MOUNTAIN RANGER DISTRICT

MALHEUR NATIONAL FOREST

GRANT COUNTY, OREGON

Prepared by:

Ian Reid

Acting Malheur National Forest Threatened and Endangered Species Coordinator

Signature: /s/ Ian Reid

Date: October 14, 2011

AQUATIC BIOLOGICAL ASSESSMENT

for

Endangered Species Act Section 7 and
Magnuson-Stevens Act Consultation

TABLE 1. ESA ACTION AREA HYDROLOGIC UNIT CODE NAMES AND NUMBERS

Seneca Allotment

4 th Fields	4 th Field HUC Numbers	6 th Fields	6 th Field HUC Numbers
Upper John Day	17070201	Middle Canyon Creek	170702010703
		Laycock Creek	170702010901
5 th Fields	5 th Field HUC Numbers		
Canyon Creek	1707020107		
Laycock Creek-John Day River	1707020109		

Deadhorse Allotment

4 th Fields	4 th Field HUC Numbers	6 th Fields	6 th Field HUC Numbers
Upper John Day	17070201	Clark Creek-John Day River	170702010904
		Riley Creek	170702010903
5 th Fields	5 th Field HUC Numbers		
Laycock Creek-John Day River	1707020109		

Hanscomb Allotment

4 th Fields	4 th Field HUC Numbers	6 th Fields	6 th Field HUC Numbers
Upper John Day	17070201	Laycock Creek	170702010901
5 th Fields	5 th Field HUC Numbers		
Laycock Creek-John Day River	1707020109		

Fields Peak Allotment

4 th Fields	4 th Field HUC Numbers	6 th Fields	6 th Field HUC Numbers
Upper John Day	17070201	Dry Creek-John Day River	170702010906
5 th Fields	5 th Field HUC Numbers	Headwaters Murderers Creek	170702010301
Murderers Creek	1707020103	Fields Creek	170702011002
Fields Creek-John Day River	1707020110		
Laycock Creek-John Day River	1707020109		

Name and Location of Administrative Unit: Blue Mountain Ranger District, Malheur National Forest, P.O. Box 909, John Day, OR 97845

TABLE 2. ESA & EFH EFFECT DETERMINATIONS (ENDANGERED (E) AND THREATENED (T) SPECIES)

Seneca Allotment

Common Name	Scientific Name	Known to Occur?	Potential to Occur?	ESA ¹ Status	Critical Habitat Status	Essential Fish Habitat Status	ESA ¹ Effect Determination Species/CH ²	EFH ³ Effect Determination
MCR steelhead	<i>Oncorhynchus mykiss</i>	Yes	Yes	Threatened	Designated	Not applicable	NLAA/NLAA ⁴	Not applicable
Spring Chinook	<i>Oncorhynchus tshawytscha</i>	No	No	Not warranted	Not applicable	Established	Not applicable/ Not applicable	Will not adversely affect
CR bull trout	<i>Salvelinus confluentus</i>	No	No	Threatened	Designated	Not applicable	No effect / No effect	Not applicable

Deadhorse Allotment

Common Name	Scientific Name	Known to Occur?	Potential to Occur?	ESA Status	Critical Habitat Status	Essential Fish Habitat Status	ESA Effect Determination Species/CH	EFH Effect Determination
MCR steelhead	<i>Oncorhynchus mykiss</i>	Yes	Yes	Threatened	Designated	Not applicable	NLAA/NLAA ⁴	Not applicable
Spring Chinook	<i>Oncorhynchus tshawytscha</i>	No	No	Not warranted	Not applicable	Established	Not applicable/ Not applicable	Will not adversely affect
CR bull trout	<i>Salvelinus confluentus</i>	No	No	Threatened	Designated	Not applicable	No effect / No effect	Not applicable

Hanscomb Allotment

Common Name	Scientific Name	Known to Occur?	Potential to Occur?	ESA Status	Critical Habitat Status	Essential Fish Habitat Status	ESA Effect Determination Species/CH	EFH Effect Determination
MCR steelhead	<i>Oncorhynchus mykiss</i>	Yes	Yes	Threatened	Designated	Not applicable	LAA/LAA ⁴	Not applicable
Spring Chinook	<i>Oncorhynchus tshawytscha</i>	No	No	Not warranted	Not applicable	Established	Not applicable/ Not applicable	Will not adversely affect
CR bull trout	<i>Salvelinus confluentus</i>	No	No	Threatened	Designated	Not applicable	No effect / No effect	Not applicable

¹ Endangered Species Act

² Critical Habitat

³ Essential Fish Habitat under the Magnuson-Stevens Fishery Conservation and Management Act

⁴ May Affect, Likely to Adversely Affect

Fields Peak Allotment

Common Name	Scientific Name	Known to Occur?	Potential to Occur?	ESA Status	Critical Habitat Status	Essential Fish Habitat Status	ESA Effect Determination Species/CH	EFH Effect Determination
MCR steelhead	<i>Oncorhynchus mykiss</i>	Yes	Yes	Threatened	Designated	Not applicable	LAA/LAA ⁴	Not applicable
Spring Chinook	<i>Oncorhynchus tshawytscha</i>	No	No	Not warranted	Not applicable	Established	Not applicable/ Not applicable	Will not adversely affect
CR bull trout	<i>Salvelinus confluentus</i>	No	No	Threatened	Designated	Not applicable	No effect / No effect	Not applicable

TABLE OF CONTENTS

1	INTRODUCTION	1
1.1	<i>Consultation History</i>	1
1.1.1	Informal Consultations (Recent and ongoing)	1
1.1.1.1	Livestock Grazing	1
1.1.1.2	Blue Mountain Expedited Section 7 Consultation Process.....	1
1.1.2	Formal Consultation (Recent and ongoing)	2
1.1.2.1	Aquatic and Riparian Restoration Programmatic Consultation	2
1.1.2.2	Livestock Grazing	3
2	DESCRIPTION OF PROJECT AREA	3
2.1	<i>Seneca Allotment</i>	3
2.2	<i>Deadhorse Allotment</i>	7
2.3	<i>Hanscomb Allotment</i>	10
2.4	<i>Fields Peak Allotment</i>	13
3	FOREST DIRECTION AND POLICIES GUIDING DEVELOPMENT OF PROPOSED ACTION.....	17
3.1	<i>Malheur National Forest LRMP</i>	17
3.1.1	LRMP Amendment 29 Desired Future Conditions	18
3.1.2	PACFISH LRMP Amendment.....	19
3.1.2.1	PACFISH Riparian Management Objectives	21
3.1.2.2	PACFISH Riparian Habitat Conservation Areas and Standards	21
3.1.2.3	PACFISH Monitoring	22
3.1.2.4	PACFISH Enclosure B: Livestock Grazing Guidelines	22
3.2	<i>Malheur National Forest Riparian Monitoring Strategy</i>	23
4	PROPOSED ACTION AND ESA ACTION AREA	27
4.1	<i>Proposed Action</i>	28
4.1.1	Permit Information and Grazing Systems	28
4.1.2	Project Design Criteria.....	29
4.1.3	Grazing Use Indicators and Supporting Rationale	30
4.1.4	Monitoring	40
4.1.5	Adaptive Management	41
4.1.5.1	Compliance Strategy for the Streambank Alteration Endpoint Indicator	42
4.1.6	Coordination and Reporting.....	43
4.2	<i>Interrelated Actions</i>	43
4.3	<i>Project Elements</i>	44

4.4	<i>ESA Action Area</i>	45
5	STATUS OF THE SPECIES AND DESIGNATED CRITICAL HABITAT	45
5.1	<i>Middle Columbia River Steelhead Distinct Population Segment</i>	46
5.1.1	Listing History and Location	46
5.1.2	Life History and Habitat Requirements	46
5.1.3	MCR Steelhead Populations	46
5.1.4	MCR Steelhead DPS Viability Status	47
5.1.5	John Day River MPG Population Status	47
5.1.6	Population Limiting Factors	49
5.2	<i>Critical Habitat for Middle Columbia River Steelhead DPS</i>	49
5.2.1	Designation History	49
5.2.2	Primary Constituent Elements	50
5.2.3	Status of Middle Columbia River Steelhead Critical Habitat	50
6	ENVIRONMENTAL BASELINE	52
6.1	<i>NMFS Matrix of Pathways and Indicators</i>	52
6.1.1	Middle Columbia River Steelhead Recovery Plan	56
6.1.2	Malheur National Forest Roads Analysis Report	57
6.1.3	Malheur National Forest Water Temperature Monitoring	61
6.2	<i>PIBO Monitoring</i>	62
6.2.1	Effectiveness Monitoring	62
6.2.1.1	Evaluation of Existing Conditions to PIBO Managed and Reference Means	66
6.2.2	Implementation Monitoring	69
7	EFFECTS OF THE PROPOSED ACTION	69
7.1	<i>Project Element and Interrelated Action Evaluation</i>	69
7.1.1	Project Elements Dropped from Further Analysis	70
7.1.2	Project Elements and Interrelated Actions with Entirely Beneficial Effects	70
7.1.3	Project Elements Remaining for Analysis	70
7.1.3.1	PE1: Livestock Use of Allotment/Pastures	70
7.1.3.2	PE2: Permittee Management of Livestock and Infrastructure Maintenance	70
7.1.3.3	PE5: Monitoring	71
7.2	<i>Analysis of Effects to Designated Critical Habitat</i>	71
7.3	<i>Analysis of Effects to Listed Species</i>	99
7.3.1	General Effects	99
7.3.2	Direct Effects to Species	99
7.3.3	Direct and Indirect Effects to Aquatic and Riparian Habitat	100
7.3.3.1	Effects on Water Temperature	101
7.3.3.2	Effects on Sediment/Turbidity, Substrate and Substrate Embeddedness	105

	7.3.3.3	Effects on Refugia.....	107
	7.3.3.4	Effects on large woody debris.....	107
8		ESA EFFECT DETERMINATIONS.....	108
	8.1	<i>Rationale</i>	108
9		ESA CUMULATIVE EFFECTS.....	110
	9.1	<i>ODFW State lands</i>	110
	9.2	<i>ODFW Elk and Deer Management</i>	110
	9.3	<i>Unauthorized Livestock Grazing</i>	112
	9.4	<i>Actions on Private Property</i>	112
10		ESSENTIAL FISH HABITAT FOR CHINOOK SALMON.....	112
11		REFERENCES.....	114

LIST OF TABLES

TABLE 1. ESA ACTION AREA HYDROLOGIC UNIT CODE NAMES AND NUMBERS	2
TABLE 2. ESA & EFH EFFECT DETERMINATIONS (ENDANGERED (E) AND THREATENED (T) SPECIES)	3
TABLE 3. STREAM MILES OF MIDDLE COLUMBIA RIVER STEELHEAD CRITICAL HABITAT IN THE SENECA ALLOTMENT	4
TABLE 4. STREAM MILES OF MIDDLE COLUMBIA RIVER STEELHEAD CRITICAL HABITAT IN THE DEADHORSE ALLOTMENT	8
TABLE 5. STREAM MILES OF MIDDLE COLUMBIA RIVER STEELHEAD CRITICAL HABITAT IN THE HANSCOMB ALLOTMENT	11
TABLE 6. STREAM MILES OF MIDDLE COLUMBIA RIVER STEELHEAD CRITICAL HABITAT IN THE FIELDS PEAK ALLOTMENT	14
TABLE 7. IDENTIFICATION OF MORE STRINGENT HABITAT INDICATOR NUMERIC VALUES OR CRITERIA BETWEEN AMENDMENT 29 DESIRED FUTURE CONDITIONS OR PACFISH RIPARIAN MANAGEMENT OBJECTIVES.....	18
TABLE 8. PERMIT INFORMATION FOR THE SENECA, DEADHORSE, HANSCOMB, FIELDS PEAK ALLOTMENT.....	28
TABLE 9. PERMIT INFORMATION - ALLOTMENT SPECIFIC	28
TABLE 10. SENECA ALLOTMENT LIVESTOCK MOVE TRIGGER AND END-POINT INDICATORS BY PASTURE.....	31
TABLE 11. DEADHORSE ALLOTMENT LIVESTOCK MOVE TRIGGER AND END-POINT INDICATORS BY PASTURE.....	32
TABLE 12. HANSCOMB ALLOTMENT LIVESTOCK MOVE TRIGGER AND END-POINT INDICATORS BY PASTURE.....	33
TABLE 13. FIELDS PEAK ALLOTMENT LIVESTOCK MOVE TRIGGER AND END-POINT INDICATORS BY PASTURE.....	34
TABLE 14. MCR STEELHEAD JOHN DAY RIVER MPG SUMMARY OF ABUNDANCE, PRODUCTIVITY, RISK RATINGS, AND MINIMUM ABUNDANCE THRESHOLDS (SOURCE: MIDDLE COLUMBIA RIVER STEELHEAD DPS RECOVERY PLAN SUMMARY 2009).....	48
TABLE 15. HABITAT LIMITING FACTORS WITHIN THE BEECH CREEK MAJOR SPAWNING AREA IDENTIFIED IN NMFS (2009).....	49
TABLE 16. STATUS OF ENVIRONMENTAL BASELINE FOR THE UPPER JOHN DAY SUB-BASIN. ¹	53
TABLE 17. RANGES OF VALUES BY RISK CATEGORY FOR ELEMENTS USED IN THE WATERSHED RISK ANALYSIS.....	57
TABLE 18. SUB-WATERSHED RISK RATINGS FOR SIXTH FIELD HYDROLOGIC UNITS IN THE ESA ACTION AREA FOR THE SENECA, DEADHORSE, HANSCOMB, AND FIELDS PEAK ALLOTMENTS CONSULTATION (FROM MNF 2004B).....	60

TABLE 19. PIBO MONITORING RESULTS – DEADHORSE ALLOTMENT.....	64
TABLE 20. PIBO MONITORING RESULTS – FIELDS PEAK ALLOTMENT	65
TABLE 21. PIBO MANAGED AND REFERENCE MEAN VALUES FOR SELECTED STREAM, CHANNEL MORPHOLOGY, AND HABITAT ATTRIBUTES.....	67
TABLE 22. PRIMARY CONSTITUENT ELEMENTS OF MCR STEELHEAD CRITICAL HABITAT APPLICABLE TO THE ACTION AREA.	71
TABLE 23. ANALYSIS OF EFFECTS TO MPI INDICATORS CORRESPONDING TO PCEs OF DESIGNATED CRITICAL HABITAT FOR MCR STEELHEAD WITHIN THE SENECA, DEADHORSE, HANSCOMB, AND FIELDS PEAK ALLOTMENTS.	73
TABLE 24. SUMMARY OF EFFECTS OF THE PROPOSED ACTION BY THE PROJECT ELEMENTS OF LIVESTOCK GRAZING IN THE SENECA, DEADHORSE, HANSCOMB, AND FIELDS PEAK ALLOTMENTS TO THE INDICATORS ASSOCIATED WITH HABITAT FEATURES OF EACH PRIMARY CONSTITUENT ELEMENT OF MCR STEELHEAD CRITICAL HABITAT.	95
TABLE 25. EFFECTIVE SHADE PROVIDED BY THREE HEIGHTS OF GREENLINE VEGETATION AT VARYING ACTIVE STREAM CHANNEL WIDTHS. ¹	103
TABLE 26. ROCKY MOUNTAIN ELK AND MULE DEER MANAGEMENT OBJECTIVES AND WINTER POPULATION ESTIMATES FROM 2004-2010 FOR THE MURDERERS CREEK WILDLIFE MANAGEMENT UNITS IN OREGON.	111

LIST OF FIGURES

FIGURE 1. SENECA ALLOTMENT AND PASTURE MAP	5
FIGURE 3. DEADHORSE ALLOTMENT AND PASTURE MAP	9
FIGURE 4. HANSCOMB ALLOTMENT AND PASTURES	12
FIGURE 5. FIELDS PEAK ALLOTMENT AND PASTURE MAP	15
FIGURE 6. ESA ACTION AREA MAP FOR THE SENECA, DEADHORSE, HANSCOMB, AND FIELDS PEAK ALLOTMENTS CONSULTATION	45
FIGURE 7. VIABILITY RATINGS FOR THE MCR STEELHEAD MAJOR POPULATION GROUP (NMFS 2009). SHADES OF GREEN INDICATE LOWER RISK OF EXTINCTION AND SHADES OF RED INDICATE HIGHER RISK.	48
FIGURE 8. HEAT EXCHANGE BETWEEN A STREAM AND ITS ENVIRONMENT.	102
FIGURE 9. SHADE PROVIDED BY 150-FOOT TALL CONIFERS (PLATTS ET AL 1987), (PARK, 1993).	103

APPENDICES

- APPENDIX A. MALHEUR NATIONAL FOREST LAND AND RESOURCE MANAGEMENT PLAN AMENDMENT 29 DESIRED FUTURE CONDITIONS
- APPENDIX B. PACFISH ENCLOSURE B
- APPENDIX C. FOREST PLAN STANDARDS AND OBJECTIVES RELATED TO RIPARIAN AREAS, WATER QUALITY AND FISH HABITAT
- APPENDIX D. PROTOCOL TO BE USED FOR EVALUATING COMPLIANCE WITH STREAMBANK ALTERATION THRESHOLDS
- APPENDIX E. PACFISH/INFISH IMPLEMENTATION MONITORING DOCUMENTS
- APPENDIX F. STRATEGY TO MINIMIZE REDD TRAMPLING “TAKE” OF STEELHEAD AND BULL TROUT
- APPENDIX G. METHODS FOR DETERMINING MOST SENSITIVE RIPARIAN AREAS
- APPENDIX H. ADAPTIVE MANAGEMENT STRATEGY
- APPENDIX I. TEMPERATURE MONITORING DATA AND EVALUATION OF RELEVANT CRITERIA FOR MCR STEELHEAD CH STREAMS WITHIN THE DEADHORSE AND FIELDS PEAK ALLOTMENTS
- APPENDIX J. PIBO AND STREAM SURVEY MONITORING DATA

1 INTRODUCTION

The Blue Mountain Ranger District of the Malheur National Forest (MNF) proposes to authorize livestock grazing for the next five seasons, 2012-16 on the Seneca, Deadhorse, Hanscomb, and Fields Peak Allotments. Consistent with the Endangered Species Act (ESA) and its implementing regulations, this Biological Assessment (BA) documents the analysis and conclusions of the Forest Service regarding the effects of implementing the livestock grazing it intends to authorize during this period. The analysis in the BA evaluates the effects on: (1) the Middle Columbia River Steelhead Distinct Population Segment (DPS) listed by the National Marine Fisheries Service (NMFS) as Threatened, and the Columbia River bull trout DPS listed by the US Fish and Wildlife Service (FWS) as Threatened; (2) designated critical habitat (CH) for both of these DPSs; and (3) Essential Fish Habitat (EFH) established for Chinook salmon, a species regulated under a Federal fisheries management plan. It is prepared in compliance with the requirements of Forest Service Manual (FSM) 2630.3, FSM 2672.4 ESA regulations, and regulations promulgated pursuant to the Magnuson-Stevens Fishery Conservation Act (MSA) as amended by the Sustainable Fisheries Act of 1996 (Public Law 104-267) (MSA §305(b)(2)).

1.1 CONSULTATION HISTORY

Past and ongoing informal and formal consultations that overlap the ESA action area and the 6th field HUC subwatersheds of Seneca, Deadhorse, Hanscomb, and Fields Peak Allotments are described in this section.

1.1.1 INFORMAL CONSULTATIONS (RECENT AND ONGOING)

1.1.1.1 LIVESTOCK GRAZING

Consultation with NMFS was concluded on the 2006 grazing season for all MNF allotments. A letter of concurrence (LOC) was completed for the 2006 grazing season for the Seneca, Deadhorse, Hanscomb, and Fields Peak Allotments since the effects were determined to not likely to adversely affect (NLAA) MCR steelhead or designated CH.

1.1.1.2 BLUE MOUNTAIN EXPEDITED SECTION 7 CONSULTATION PROCESS

The MNF received a concurrence letter in June 2007 (NMFS 2007a) from NMFS (2007/02970) for a consultation on the effects of the Blue Mountain Expedited Section 7 Consultation Process (Process). The LOC is currently active, and applies through June 2012. The Process evaluates consistency of an action with a set of project design criteria (PDC). Among the categories of actions specifically considered for application of the Process are: (1) vegetation management (mechanized and non-mechanized); (2) livestock grazing (range improvements); (3) wildlife, fish or watershed improvement projects; (4) road maintenance; (5) low impact permits; and, (6) recreation and administrative sites. Other types of projects can be covered under the programmatic as long as all of the PDC are met. The Level I team will determine if the use of the expedited process is appropriate for that project.

The action agency prepares documentation evaluating whether or not the action is consistent with the PDC covered by the concurrence letter. If an action is determined to be consistent with all of the PDC after evaluation by the Level 1 team, then an ESA effect determination of “May Affect, NLAA” applies. The Level 1 team then sends a letter to the respective action agency official, documenting its finding regarding consistency with the Process and stating that the letter tiers to the concurrence letter.

Actions occurring in the allotments that are covered by the Process:

MNF Road Maintenance

The MNF has consulted with the NMFS and USFWS on Forest wide road maintenance. On January 29, 2010 a letter was sent to the Malheur National Forest Supervisor from the interagency members of the Malheur Level 1 Team (FS, BLM, NMFS and FWS). The letter tiered to the 2007 NMFS concurrence letter for the Process and stated that the team had reviewed the PDC documentation package for the MNF Road Maintenance program for consistency with the Process. The team agreed with the MNF finding that the project “may affect, but is not likely to adversely affect (NLAA) the species and their designated CH for Mid-Columbia River Steelhead and Bull Trout based on the rationale that was presented for consistency with all PDC in the documentation package.” The Malheur Level 1 Team also concluded that the documentation package demonstrated that the action would adequately avoid, minimize or otherwise offset potential effects to designated EFH and fulfilled requirements under the MSA. The letter provided ESA and MSA coverage for the Forest’s Road Maintenance program from 2010 to 2015.

1.1.2 FORMAL CONSULTATION (RECENT AND ONGOING)

1.1.2.1 AQUATIC AND RIPARIAN RESTORATION PROGRAMMATIC CONSULTATION

The Forest Service and BLM concluded formal consultation on June 27, 2008 with NMFS (2008/03505) on Forest Service and BLM aquatic restoration activities for administrative units in Oregon and Washington including the MNF. The biological opinion (BO) applies through CY 2012, and provides coverage for 19 aquatic restoration program activity types:

1. Large Wood, Boulder, and Gravel Placement
2. Reconnection of Existing Side Channels and Alcoves
3. Head-cut Stabilization and Associated Fish Passage
4. Bank Restoration
5. Fish Passage Culvert and Bridge Projects
6. Irrigation Screen Installation and Replacement
7. In-channel Nutrient Enhancement
8. Floodplain Overburden Removal
9. Reduction of Recreation Impacts
10. Estuary Restoration
11. Riparian Vegetation Treatment (non-commercial, mechanical)
12. Riparian and Upland Juniper Treatment (non-commercial)
13. Riparian Vegetation Treatment (controlled burning)
14. Riparian Area Invasive Plant Treatment
15. Riparian Exclusion Fencing (with water gaps and stream crossings)
16. Riparian Vegetation Plantings
17. Road Treatments
18. Removal of Legacy Structures
19. Fisheries, Hydrology, Geomorphology Wildlife, Botany, and Cultural Surveys in Support of Aquatic Restoration

Actions occurring in the ESA action area that are covered by the Aquatic and Riparian Restoration Programmatic Consultation:

Fields Peak Allotment Improvement Project Murderers Creek Fence. This project included construction of fences to restrict livestock use and improve distribution to maintain/restore stream channel, riparian vegetation, and floodplain functions of Murderers Creek, Lemon Creek, Lime Creek, and Grapefruit

Creek. The project occurred in 2009 and included design criteria specified in aquatic restoration program activity type #15.

1.1.2.2 LIVESTOCK GRAZING

In 2007 the MNF formally consulted with NMFS on the 2007-2011 livestock grazing in the Seneca, Deadhorse, Hanscomb, and Fields Peak Allotments (NMFS reference number 2007/01290, NMFS 2007b). Although outside the ESA action area of these allotments but within two of the same subwatersheds, a formal consultation with NMFS (reference number 2007/01290) was also completed in 2007 for the Murderers Creek allotment, which is located in the Headwaters Murderers Creek (formerly Tex Creek) 6th field subwatershed.

2 DESCRIPTION OF PROJECT AREA

2.1 SENECA ALLOTMENT

The Seneca Allotment is located within the Upper John Day River (HUC # 17070201) and Silvies (HUC # 17120002) subbasin (Table 1). The pastures comprising the Seneca Allotment lie within the Canyon Creek (HUC # 1707020107), Headwaters Silvies River (HUC # 1712000201, outside of the ESA action area), and Laycock Creek-John Day River (HUC # 1707020109) watersheds, mostly within T 15 S, and R 30 and 31 E. The Allotment includes approximately 10,166 acres of National Forest System (NFS) Lands. Approximately 100 acres of private land are intermingled with NFS lands. These lands are unfenced and management of these lands has not been waived to the Forest Service. Elevations within the allotment range from approximately 4,000 feet in Vance Creek to 5,800 feet along Starr Ridge.

Dominant grass species throughout the allotment are bluebunch wheatgrass, Idaho fescue, elk sedge, and pine grass. Riparian overstory vegetation generally consists of a mix of hardwood and conifer species along the stream with alder being the dominant species.

Throughout this allotment, livestock have varying levels of access to streams and the associated riparian communities. Parameters such as gradient, valley form, geologic substrate, vegetative structure, and forage availability can greatly influence livestock movement, use patterns, and distribution relative to streams. Other factors, such as the presence of “windthrown” or “jack-strawed” timber, may also influence livestock accessibility to streams and riparian communities.

Shade is provided by grass and grass-like species, riparian hardwood species and conifer species along the stream, as well as topography due in part to significantly incised stream channels on several streams within the allotment. Historically, riparian areas were logged by conventional tractor yarding. Railroad logging also occurred in and along many of the streams within the Seneca Allotment. The combination of logging, insect epidemic, and valley bottom roads has reduced shading from conifer species.

The watersheds encompassing the Seneca Allotment support a mix of National Forest System and State and private lands. Activities that have occurred or continue to occur within these watersheds include historic mining, timber harvest, grazing, roads, trails, prescribed and natural fire, noxious weed treatment, and recreation.

The Seneca Allotment is divided into 4 pastures: Vance Creek, Camp Creek, Camp Management, and Koehler (Figure 1). The allotment contains 1.02 miles of steelhead CH (Table 3).

Descriptions of individual pastures are presented below.

TABLE 3. STREAM MILES OF MIDDLE COLUMBIA RIVER STEELHEAD CRITICAL HABITAT IN THE SENECA ALLOTMENT

Stream Name	Miles of Steelhead Critical Habitat
Vance Creek	0.92
Hanscombe Creek	0.10
Total	1.02

Seneca Allotment




FIGURE 1. SENECA ALLOTMENT AND PASTURE MAP

Vance Creek Pasture

Elevations in the Vance Creek pasture vary from approximately 5,800 feet to approximately 4,000 feet. Overstory vegetation in the pasture contains primarily Ponderosa pine and Douglas fir. Understory vegetation consists primarily of mixed wheat grasses and bluebunch wheatgrass/Idaho fescue. Elk sedge and pine grass can be found under dense overstory stands on north slopes. This pasture has not been grazed since 2007.

The Vance Creek pasture contains 1.02 miles of designated steelhead critical habitat in the Vance and Hanscombe Creeks. Only 0.1 miles of Hanscombe Creek lie within the pasture in the extreme northwest corner of the pasture although there is no steelhead access to this stream section due to a large falls downstream on private land (R. Holliday pers. comm. permittee comments letter). Vance Creek has dense stands of alder and dogwood throughout much of the stream. Access by livestock is limited to a few existing trail crossings. No PIBO data are available for Vance Creek, but a level 2 survey was conducted in 1993 that found relatively little cattle use along the stream as much of it was enclosed by dense stands of alder, willow, and dogwood (MNF 1993) South Fork Vance Creek does not contain critical habitat, but is within the Vance Creek pasture within several hundred feet of its confluence with Vance Creek and had heavy grazing utilization during its level 2 stream survey conducted in the early 1990s (MNF 1993). Hanscombe Creek was surveyed in 1995, was found to be non-fish-bearing and had heavy grazing utilization in its lower reach, which included private land (MNF 1995a).

Camp Creek, Camp Management and Koehler Pastures

The Camp Creek, Camp Management and Koehler pastures are located entirely within the Silvies subbasin which does not support anadromous fishes or their habitat. These pastures will not be discussed further.

2.2 DEADHORSE ALLOTMENT

The Deadhorse Allotment is located within the Upper John Day River (HUC # 17070201) and Silvies (HUC # 17120002) subbasin (Table 1). The pastures comprising the Deadhorse Allotment lie within the Headwaters Silvies River (HUC # 1712000201) and Laycock Creek-John Day River (HUC # 1707020109) watersheds, mostly within T 14 and 15 S, and R 29 and 30 E. The Allotment includes approximately 15,527 acres of National Forest System (NFS) Lands. Elevations within the allotment range from approximately 3,600 feet on Riley Creek at the Forest boundary to 7,000 feet on Packsaddle Ridge.

Dominant grass species throughout the allotment are bluebunch wheatgrass, Idaho fescue, elk sedge, and pine grass. Riparian overstory vegetation generally consists of a mix of hardwood and conifer species along the stream with alder being the dominant species.

Throughout this allotment, livestock have varying levels of access to streams and the associated riparian communities. Parameters such as gradient, valley form, geologic substrate, vegetative structure, and forage availability can greatly influence livestock movement, use patterns, and distribution relative to streams. Other factors, such as the presence of “windthrown” or “jack-strawed” timber, may also influence livestock accessibility to streams and riparian communities.

Shade is provided by grass and grass-like species, riparian hardwood species and conifer species along the stream, as well as topography due in part to significantly incised stream channels on several streams within the allotment. Historically, riparian areas were logged by conventional tractor yarding. Railroad logging also occurred in and along many of the streams within the Seneca Allotment. The combination of logging, insect epidemic, and valley bottom roads has reduced shading from conifer species.

The watersheds encompassing the Deadhorse Allotment support a mix of National Forest System and State and private lands. Activities that have occurred or continue to occur within these watersheds include historic mining, timber harvest, grazing, roads, trails, prescribed and natural fire, noxious weed treatment, and recreation.

There are two major drainages within the Deadhorse Allotment; Riley Creek and Ingle Creek. Riley Creek bisects the Aldrich Mountains and flows mostly through a rocky narrow canyon that is inaccessible to cattle. There is a 15-foot-high waterfall approximately 1 mile upstream of the Malheur National Forest (MNF) boundary on Riley Creek that blocks access to steelhead (MNF 1995b). On MNF land Riley Creek contains steelhead/redband and westslope cutthroat trout, however livestock access to the stream is limited by steep canyon walls, abundant downed wood and thick riparian vegetation, and little use was reported in the 1995 stream survey report. This lower mile of Riley Creek is contained within a steep, confined canyon with large cobble and small boulder substrate, mixed in with spawning gravels.

The Deadhorse Allotment is divided into 4 pastures: North, Riley, Riley Meadows, and Percival (Figure 3). The allotment contains 4.28 miles of steelhead.

Descriptions of individual pastures are presented below.

TABLE 4. STREAM MILES OF MIDDLE COLUMBIA RIVER STEELHEAD CRITICAL HABITAT IN THE DEADHORSE ALLOTMENT

Stream Name	Miles of Steelhead Critical Habitat
Ingle Creek	2.85
Riley Creek	1.43
Total	4.28




FIGURE 3. DEADHORSE ALLOTMENT AND PASTURE MAP

North Pasture

The North pasture is a very large (10,236 acres) pasture consisting of generally steep (very steep in some areas – with cliffs in the north end of the unit, adjacent to Riley Creek), open hills bisected by narrow drainages. This pasture has no roads and is located in a non-motorized area. Grazing has not been authorized in this pasture since 2007.

Riley Creek and Ingle Creek are the only streams with potential steelhead habitat in the North Unit. A waterfall approximately one mile upstream from the MNF boundary on Riley Creek creates a fish barrier. Due to the steep and dry terrain, cattle from this allotment do not normally get into the lower (north) portion of Riley Creek, but limited use has been previously reported (MNF 1995b). Ingle Creek contained steelhead/redband trout 0.8 miles above the forest boundary during a 1995 survey (MNF 1995c). This survey also reported that anadromous fish do not reach the forest boundary (meaning the trout would all be redband) but no rationale was given why these were all resident fish. Heavy cattle (and elk) use was documented in spring and wetland areas adjacent and hydrologically connected to Ingle Creek in 1995, but the use appeared to be unauthorized as the pasture was deferred that season (MNF 1995c). The report also mentioned that livestock and big game grazing and trampling of the wetlands is a problem in this drainage and that reducing grazing activities [presumably referring to unauthorized use] in reach 2 (critical habitat reach) to increase riparian ground cover, shrub cover, and allow wetlands to recover would help reduce potential risk of sedimentation, and effect [decrease] water temperature (MNF 1995c).

Riley and Riley Meadows Pastures

The Riley and Riley Meadows pastures are located within the Upper John Day River subbasin, however they do not contain steelhead critical habitat and will not be discussed further.

Percival Pasture

The Percival pasture is located entirely within the Silvies subbasin which does not support anadromous fish or their habitat. This pasture will not be discussed further.

2.3 HANSCOMB ALLOTMENT

The Hanscomb Allotment is located within the Upper John Day River (HUC # 17070201) and Silvies (HUC # 17120002) subbasin (Table 1). The pastures comprising the Hanscomb Allotment lie within the Headwaters Silvies River (HUC # 1712000201) and Laycock Creek-John Day River (HUC # 1707020109) watersheds, mostly within T 14 and 15 S and R 30 E. The Allotment includes approximately 9,102 acres of National Forest System (NFS) Lands. Elevations within the allotment range from approximately 4,500 feet in Laycock Creek to 6,700 feet on Coal-Pit Mountain.

Dominant grass species throughout the allotment are bluebunch wheatgrass, Idaho fescue, elk sedge, and pine grass. Riparian overstory vegetation generally consists of a mix of hardwood and conifer species along the stream with alder being the dominant species.

Throughout this allotment, livestock have varying levels of access to streams and the associated riparian communities. Parameters such as gradient, valley form, geologic substrate, vegetative structure, and forage availability can greatly influence livestock movement, use patterns, and

distribution relative to streams. Other factors, such as the presence of “windthrown” or “jack-strawed” timber, may also influence livestock accessibility to streams and riparian communities.

Shade is provided by grass and grass-like species, riparian hardwood species and conifer species along the stream, as well as topography due in part to significantly incised stream channels on several streams within the allotment. Historically, riparian areas were logged by conventional tractor yarding. Railroad logging also occurred in and along many of the streams within the Seneca Allotment. The combination of logging, insect epidemic, and valley bottom roads has reduced shading from conifer species.

The watersheds encompassing the Hanscomb Allotment support a mix of National Forest System and State and private lands. Activities that have occurred or continue to occur within these watersheds include historic mining, timber harvest, grazing, roads, trails, prescribed and natural fire, noxious weed treatment, and recreation.

The Hanscomb Allotment is divided into 4 pastures: Laycock, Upper Geary, Geary Meadows, and Allen/Morris (Figure 4). The allotment contains 4.28 miles of steelhead CH and 0.19 miles of “Most Sensitive Riparian Areas” (MSRA) along Laycock Creek (Table 5). The MSRA are areas that the MNF has identified that are the most accessible and sensitive to livestock impacts within streams containing steelhead CH.

Descriptions of individual pastures are presented below.

TABLE 5. STREAM MILES OF MIDDLE COLUMBIA RIVER STEELHEAD CRITICAL HABITAT IN THE HANSCOMB ALLOTMENT

Stream Name	Miles of Steelhead Critical Habitat	Miles of MSRA
Laycock Creek	1.48	0.19
Hanscombe Creek	0.39	
Total	1.87	0.19




FIGURE 4. HANSCOMB ALLOTMENT AND PASTURES

Laycock Pasture

Elevations in the Laycock pasture vary from approximately 4,500 feet to approximately 6,700 feet. The Laycock Creek pasture consists of steep slopes covered with relatively dense timber on the upper two thirds of the unit grading into pine/juniper and sagebrush on the lower slopes.

Streams containing steelhead critical habitat within the Laycock pasture are: Laycock and Hanscombe Creeks. The upper portions of Laycock Creek and Hanscombe Creek contain mainly unconsolidated ash material in the substrate. Laycock Creek contained small populations of cutthroat and redband/steelhead trout in a 1995 level II survey, but fish habitat was severely limited by sedimentation caused primarily by a landslide/debris torrent (MNF 1995d). Livestock grazing was reported to be a contributing factor to sedimentation, which was a concern in this stream (MNF 1995d). Reach 2 of Laycock Creek, which contains CH, reported that riparian ground cover was 29% below minimum forest standards and that streambank trampling needs to be minimized especially in the marsh areas where soils are easily disturbed (MNF 1995d). See description of Hanscombe Creek above in Seneca Allotment narrative.

Upper Geary, Geary Meadows, Allen/Morris Pastures

The Upper Geary, Geary Meadows, Allen/Morris pastures are located entirely within the Silvies subbasin which does not support anadromous fish or their habitat. These pastures will not be discussed further.

2.4 FIELDS PEAK ALLOTMENT

The Fields Peak Allotment is located within the Upper John Day River (HUC # 17070201) subbasin (Table 1). The pastures comprising the Fields Peak Allotment lie within the Murderers Creek (HUC # 1707020103), Fields Creek-John Day River (HUC # 1707020110), and Laycock Creek-John Day River (HUC # 1707020109) watersheds, mostly within T 14 and 15 S and R 28, 29, and 30 E. The Allotment includes approximately 30,718 acres. Approximately 272 acres of private land are intermingled with NFS lands. The private lands are unfenced and management of these lands has not been waived to the Forest Service. Elevations within the allotment range from approximately 3,200 feet in Fields Creek to 7,300 feet.

Dominant grass species throughout the allotment are bluebunch wheatgrass, Idaho fescue, elk sedge, and pine grass. Riparian overstory vegetation generally consists of a mix of hardwood and conifer species along the stream with alder being the dominant species.

Throughout this allotment, livestock have varying levels of access to streams and the associated riparian communities. Parameters such as gradient, valley form, geologic substrate, vegetative structure, and forage availability can greatly influence livestock movement, use patterns, and distribution relative to streams. Other factors, such as the presence of “windthrown” or “jack-strawed” timber, may also influence livestock accessibility to streams and riparian communities.

Shade is provided by grass and grass-like species, riparian hardwood species and conifer species along the stream, as well as topography due in part to significantly incised stream channels on several streams within the allotment. Historically, riparian areas were logged by conventional tractor yarding. Railroad logging also occurred in and along many of the streams within the Seneca Allotment. The combination of logging, insect epidemic, and valley bottom roads has reduced shading from conifer species.

The watersheds encompassing the Fields Peak Allotment support a mix of National Forest System and State and private lands. Activities that have occurred or continue to occur within these watersheds include historic mining, timber harvest, grazing, roads, trails, prescribed and natural fire, noxious weed treatment, and recreation.

The Fields Peak Allotment is divided into 5 pastures: Fields Peak, Tex Creek, Miners Creek, North Murderers Creek, and Murderers Creek (Figure 5). The allotment contains 22.36 miles of steelhead CH. Streams that contain MSRA within the Fields Peak Allotment are Fields, Wickiup, Tex, and Murderers creeks (Table 6). Cattle are proposed to actively graze, trail, and water in these MSRAs.

Descriptions of individual pastures are presented below.

TABLE 6. STREAM MILES OF MIDDLE COLUMBIA RIVER STEELHEAD CRITICAL HABITAT IN THE FIELDS PEAK ALLOTMENT

Stream Name	Miles of Steelhead Critical Habitat	Miles of MSRA
Fields Creek	5.10	0.10
Wickiup Creek	0.90	0.11
Buck Cabin Creek	2.30	
Tex Creek	4.68	3.22
Miner Creek	0.83	
Sugar Creek	0.67	
Basin Creek	0.48	
White Creek	0.67	
Charlie Mack Creek	0.51	
Lemon Creek	0.89	
Murderers Creek	5.33	4.51
Total	22.36	7.94

Fields Peak Allotment




FIGURE 5. FIELDS PEAK ALLOTMENT AND PASTURE MAP

Fields Peak Pasture

Elevations in the Fields Peak pasture vary from approximately 3,300 feet at the Forest boundary along the 21 Road to approximately 7,300 feet at Fields Peak.

Streams containing steelhead critical habitat within the Fields Peak allotment are: Fields Creek, Wickiup Creek, and Buck Cabin Creek.

Fields Creek is likely the only stream with steelhead spawning habitat within the Fields Creek pasture, although other tributaries to Fields Creek likely provide rearing habitat for MCR steelhead. A 1992 Level II survey found embeddedness to exceed 35% in most reaches. It also reported “grazing throughout the reaches has caused sediment to be added to the stream, along with eroding some streambanks. This grazing however is not causing major damage” (MNF 1992).

Miners Creek Pasture

Elevations in the Miners Creek pasture vary from approximately 4,700 feet on Tex Creek to approximately 7,000 feet near the pasture boundary with the Fields Creek pasture. The Miners Creek pasture is primarily south and west facing, with the Tex Creek drainage being the major feature of the pasture. Wild horses have been observed in this pasture. They are typically individual horses or small bands of 3-5 horses.

Streams containing steelhead critical habitat within the Miners Creek pasture are: Tex Creek, Miner Creek, and Sugar Creek.

Tex Creek Pasture

Elevations in the Tex Creek pasture vary from approximately 4,400 feet along Tex Creek to approximately 5,500 feet at the Fields Creek summit on the 21 road. Overstory vegetation is primarily mixed conifer of Ponderosa pine and Douglas fir with lesser components of Grand fir and western Larch. There are a couple of wild horse bands that reside within this pasture.

Tex Creek is the only stream containing steelhead critical habitat within the pasture and is a documented steelhead spawning stream (Powell and Delano 2004).

North Murderers Creek Pasture

Elevations in the North Murderers Creek pasture vary from approximately 4,500 feet along the 21 Road to approximately 5,800 feet in the northeast corner of the pasture. In 2009 this pasture was created when a new fence was constructed along Murderers Creek and FS Road 21. The fence was constructed to provide for additional protection to Murderers Creek and allow for greater management flexibility and grazing rotations.

This pasture is predominately south facing with Ponderosa pine being the dominate conifer. There are also several aspen stands within the pasture. Wild horses have been observed in this pasture.

Streams containing steelhead critical habitat within the North Murderers Creek pasture are: Charlie Mack Creek, Basin Creek, and White Creek. Although identified as containing steelhead critical habitat, MNF biologists have determined that the lower section of Charlie Mack Creek does not support fish during summer months. The lower section of the stream suitable for spawning loses surface flow by mid to late June. Above this section (at about rivermile 0.3) stream gradient increases and it becomes enclosed by dense alders, aspens, and other hardwoods.

Murderers Creek Pasture

Elevations in the Murderers Creek pasture vary from approximately 4,500 along Murderers Creek to 5,700 feet in the southwest corner of the pasture. This pasture is a north facing pasture dominated by Ponderosa pine and Grand fir. Wild horses have been observed in this pasture.

Streams containing steelhead critical habitat within the Murderers Creek pasture are: Murderers Creek and Lemon Creek. Murderers Creek is a documented steelhead spawning stream (Powell and Delano 2004).

3 FOREST DIRECTION AND POLICIES GUIDING DEVELOPMENT OF PROPOSED ACTION

Forest direction and policies provide a management framework that direct or guide development of grazing actions on the MNF. Components of the management framework include the MNF Land and Resource Management Plan (LRMP), pertinent LRMP amendments and Forest policies. The most pertinent amendments to the MNF LRMP are PACFISH and Amendment 29. The MNF Riparian Monitoring Strategy is a forest policy, MNF (2006).

3.1 MALHEUR NATIONAL FOREST LRMP

The MNF LRMP (MNF 1990) contains goals and objectives for the Range Program that provide direction with respect to range management and other resources. Goals 20 - 22 of the MNF LRMP for the Range program state:

“Provide a sustained production of palatable forage for grazing by livestock and dependent wildlife species.”

“Manage rangelands to meet the needs of other resources and uses at a level which is responsive to site-specific objectives.”

“Permit livestock use on suitable range when the permittee manage livestock using prescribed practices.”

A Range program MNF LRMP Objective also provides context:

“Analyze allotments to determine proper stocking levels. Use specific management area goals and standards to resolve conflicts between wild horses, cattle and, and big game.”

Rangeland will be managed to meet the needs of ESA-listed MCR Steelhead and big game as “other resources.” The MNF Range program LRMP Objective directs that when there are conflicts between wild horse, cattle and big game uses in determining stocking levels, management area goals and standards will be used to resolve the conflicts.

The LRMP direction described above provides conservation benefits to ESA-listed MCR Steelhead and its designated CH by directing that the needs of other resources will be met.

Other components of the Forest management framework (MNF LRMP) that guide the development of the proposed action are discussed under the Forest amendments sections of the BA. The most pertinent amendments to the MNF LRMP are PACFISH and Amendment 29.

3.1.1 LRMP AMENDMENT 29 DESIRED FUTURE CONDITIONS

The MNF Land and Resource Management Plan (MNF 1990) was amended in 1994 (Amendment 29) in response to the Columbia River Basin Anadromous Fish Habitat Management Policy and Implementation Guide (USDA FS 1991). The Forest modified Standard 5 of the Fish and Wildlife resource elements. The amended Standard 5 included specific numerical desired future conditions (DFCs) for Management Area 3A (non-anadromous riparian areas) and Management Area 3B (anadromous riparian areas). The DFCs addressed: 1) sediment/substrate, 2) water quality, 3) stream channel morphology, and 4) riparian vegetation. See Amendment 29 for the specific numeric values (Appendix A). The numerical DFCs were selected to protect water quality, features of riparian vegetation, and components of fish habitat.

Amendment 29 did not set specific quantifiable standards for livestock grazing activities. However, grazing activities can directly affect the attainment of Amendment 29 DFCs for: 1) sediment/substrate (cobble embeddedness), 2) channel morphology (bank stability, lower bank angle, width to depth ratios, 3) riparian vegetation (ground cover, percentage of stream bank vegetated), and 4) shade/canopy closure (hardwood/meadow complex). DFCs were developed to provide the criteria against which attainment or progress toward attainment of the riparian goals are measured. The MNF manages according to the more stringent standards applicable to habitat components of anadromous riparian areas as between Amendment 29 DFCs and the Riparian Management Objectives (RMOs) of the PACFISH amendment, although it should be added that the two are not always directly correlative or equally applicable; for example, with respect to bank stability, the Amendment 29 DFC applies to forested areas only. Nevertheless, this overarching directive provides conservation benefits to ESA-listed MCR Steelhead and its designated CH. Table 7 presents Amendment 29 DFCs and PACFISH RMOs by habitat indicator/criterion and displays which of the two is more stringent to the extent that both may be applicable in a given management situation.

TABLE 7. IDENTIFICATION OF MORE STRINGENT HABITAT INDICATOR NUMERIC VALUES OR CRITERIA BETWEEN AMENDMENT 29 DESIRED FUTURE CONDITIONS OR PACFISH RIPARIAN MANAGEMENT OBJECTIVES.

Habitat Indicator	Desired Future Condition or Riparian Management Objective		More Stringent Condition or Objective
	Amendment 29	PACFISH RMO	
Cobble embeddedness	<20%	NA	Amendment 29
Water temp.	No increase if <68°F, reduce to 68°F if >68°F (See Appendix A for DFC details)	No measurable increase. Max below 64°F for migration/rearing, max below 60°F for spawning	PACFISH RMO
Large Woody Debris	Varies by ponderosa (20-70/mi), Mixed conifer (80-	>20/mi >12" dia >35' length	Amendment 29 is more specific

	120/mi), lodgepole (100-350/mi). Sizes vary.		
Pool frequency	Range expected for Rosgen B&C streams, upper limits adjusted for streams >75 ft. to be consistent w/PACFISH. Provides table w/ranges by bankfull width	Table provided shows pools/mile by wetted width. All values fall within ranges by BFW of Amendment 29	Same
Bank stability (forested)	90% and no decrease if above 90%	>80%	Amendment 29
Lower bank angle (non-forested)	50-75% of banks w/90 degree angle or greater	>75% w/90 degree angle	PACFISH RMO
W/D ratio	<10	<10	Same
Potential LWD forest	To provide a rate of input to maintain LWD standard	NA	Amendment 29
Ground cover	90% of site potential	NA	Amendment 29
% streambank vegetated	90% of site potential	NA	Amendment 29
Shade/canopy closure	Varies by conifer species forest. Hardwood/meadow complex 80% shaded	NA	Amendment 29

3.1.2 PACFISH LRMP AMENDMENT

PACFISH applies specifically to the MNF lands within the range of anadromy including the Seneca, Deadhorse, Hanscomb, and Fields Peak Allotments. PACFISH amended Forest Service Land and Resource Management Plans (LRMPs). PACFISH contains the following components that provide the necessary direction and objectives, and regulatory certainty that FS management actions will be designed to maintain and restore ecological processes that support high quality habitat for salmon and steelhead over the long term:

- Riparian Goals;
- Riparian Management Objectives (RMOs);

- Delineation of streamside areas (Riparian Habitat Conservation Areas) that are important to maintenance of high quality aquatic habitat and where special management considerations are applied;
- Standards and/or guidelines to ensure projects do not prevent or retard attainment of riparian goals and management objectives;
- Designation of Key watersheds where additional management emphasis and/or watershed analysis is required to ensure that salmon and steelhead habitat is maintained or provided priority for restoration;
- Watershed analyses to provide sufficient context for designing actions that support maintenance or restoration of aquatic habitats needed for recovery of ESA-listed salmon and steelhead;
- Targeted watershed restoration identified through watershed analysis;
- Monitoring program to evaluate the implementation (compliance) and effectiveness of PACFISH in improving aquatic habitat on federal lands.

Riparian Goals provide management context for proposed activities. The goals of PACFISH establish an expectation of the characteristics of healthy, functioning watershed, riparian areas, and associated fish habitats. They are stated in relatively broad, generic terms such that they can be said to apply to most riparian areas regardless of stream type and other more site-specific conditions, but need to be evaluated in the context of the particular stream at issue. Since the quality of water and fish habitat in aquatic systems is inseparably related to the integrity of upland and riparian areas within watersheds, PACFISH articulates the following goals to maintain or restore:

- Water quality, to a degree that provides for a stable and productive riparian and aquatic ecosystem;
- Stream channel integrity, channel processes and sediment regime (including the elements of timing, volume, and character of sediment input and transport) under which riparian and aquatic ecosystems developed;
- Instream flows to support healthy riparian and aquatic habitats, stable and functioning channels, and the ability to route flood flows;
- Natural timing and variability of water tables in meadows and wetlands;
- Diversity and productivity of native and desirable non-native plant communities in riparian zones;
- Riparian vegetation to provide for 1) an amount and distribution of large woody debris characteristic of natural aquatic and riparian ecosystems, 2) adequate summer and winter thermal regulation within the riparian and aquatic zone, and 3) rates of surface erosion, bank erosion, and channel migration characteristics of those under which the communities developed;
- Riparian and aquatic habitats necessary to foster unique genetic fish stock that evolved within the specific geo-climatic region; and,
- Habitat to support populations of well-distributed native and non-native plant, vertebrate and invertebrate populations that contribute to the viability of riparian-dependent communities.

3.1.2.1 PACFISH RIPARIAN MANAGEMENT OBJECTIVES

Interim quantitative Riparian Management Objectives (RMOs) for stream channel, riparian and watershed conditions were also developed as part of PACFISH to provide criteria against which attainment or progress of the strategy's riparian goals may be measured. The objectives need to be evaluated and assessed temporally to reflect the ecological capabilities of specific ecosystems and the fact that attainment of or progress toward many of the objectives is only able to occur over extended periods of time. In general, and to the extent applicable and feasible, the MNF manages livestock grazing so as not to prevent or retard attainment of these RMOs unless Forest Plan amendment # 29 is more stringent, which will benefit habitat for MCR Steelhead.

Bank Stability: at least 80%

Water Temperature: No measureable increase in maximum temperature; Meet state water quality standards. The standard is defined as: All streams identified as having anadromous fish passage and salmonid rearing use for Designated Beneficial Use purposes. 7 Day Mean Max 64°F (17.8°C) (migration and rearing habitat); 7 Day Mean Max 60°F (15.6°C) (spawning habitat).

Width-to-Depth Ratio (W:D): W:D <10, mean wetted width divided by mean depth (NMFS PACFISH BO 2000); or **Bankfull Width-to-Depth Ratio** within 75th percentile of the range for minimally managed or reference watershed conditions (i.e. healthy streams) by stream type (analysis pending from PACFISH/INFISH biological opinions (PIBO) Effectiveness Monitoring Team).

3.1.2.2 PACFISH RIPARIAN HABITAT CONSERVATION AREAS AND STANDARDS

Project and site-specific standards apply to all Riparian Habitat Conservation Areas (RHCAs) and to projects and activities in areas outside RHCAs that would degrade them. Standards and/or guidelines were developed to ensure to the extent practicable given site conditions that projects do not prevent or retard attainment of or reasonable progress toward riparian goals and management objectives. PACFISH (USDA FS and USDI BLM 1994) standards for livestock management are presented below.

- GM-1 - Modify grazing practices (e.g., accessibility of riparian area to livestock, length of grazing season, stocking levels, timing of grazing, etc.) that retard or prevent attainment of Riparian Management Objectives or are likely to adversely affect listed anadromous fish. Suspend grazing if adjusting practices is not effective in meeting Riparian Management Objectives and avoiding adverse effects on listed anadromous fish (PACFISH).
- GM-2 – Locate new livestock handling and/or management facilities outside of Riparian Habitat Conservation Areas. For existing livestock handling facilities inside the Riparian Habitat Conservation Areas, assure that facilities do not prevent attainment of Riparian Management Objectives or adversely affect listed anadromous fish. Relocate or close facilities where these objectives cannot be met.
- GM-3 – Limit livestock trailing, bedding, watering, salting, loading, and other handling efforts to those areas and times that will not retard or prevent attainment of Riparian Management Objectives or adversely affect listed anadromous fish.

Implementing these standards clearly provide a conservation benefit to MCR Steelhead and designated CH.

3.1.2.3 PACFISH MONITORING

The PACFISH Monitoring Strategy was designed to feed information back to management for decision making. The implementation strategy uses “endpoints” and “triggers” to assess whether the authorized grazing is having the anticipated effects to resources and as an additional precautionary measure that helps the Forest Service to ensure that PACFISH direction is met as the season progresses. The endpoints measure annual conditions after grazing has been completed and the triggers measure conditions during grazing to determine if adjustments are necessary or appropriate to be made to the rotation and schedule the Forest Service has developed based on its best professional judgment and projections for a particular grazing season. The results of implementation monitoring are to be fed into effectiveness monitoring, which uses “trend” of long-term indicators of habitat condition to assess the need for revising management. This monitoring has a direct application to habitat features essential to long-term conservation of salmon and steelhead.

The PIBO Effectiveness Monitoring (EM) Program (under the Interagency Deputy Team) was initiated to evaluate the effects of land management activities on aquatic and riparian communities at multiple scales and to determine whether PACFISH management practices are effective in maintaining or improving the structure and function of riparian and aquatic conditions. A pilot study was begun in 1998 on Forest Service lands within the Salmon River basin of central Idaho. In 2000, the pilot study was expanded to include additional Federal Lands in the interior Columbia River basin. The study area includes 20 National Forests and nine BLM field units within the interior Columbia River basin. Results from sample size analysis suggested that the monitoring program will be able to detect changes in resource condition at the scale of individual Forests and BLM field offices (35 to 90 sites) for many of the attributes measured. The PACFISH Effectiveness Monitoring Program sampling design anticipates collecting information at least through 2015.

The PIBO EM and Implementation Monitoring (IM) programs are coordinated such that data collected to assess trend is linked to management actions taken under the PACFISH strategy. Thus, monitoring sites selected for evaluating the effectiveness of PACFISH are also monitored for compliance with the standards and guidelines. Preliminary results from broad-scale aquatic habitat status and trend monitoring of FS and BLM lands within the interior Columbia River basin since 2001 indicates conditions have improved over the past 5 years, continuing the habitat recovery the agencies intended to commence upon their adoption in 1995 of the protections in PACFISH instituted (NMFS 2009).

3.1.2.4 PACFISH ENCLOSURE B: LIVESTOCK GRAZING GUIDELINES

A revision of PACFISH Enclosure B, the “Recommended Livestock Grazing Guidelines,” was sent to the PACFISH Forest Supervisors on August 14, 1995 (Appendix B). The guidelines were recommended for use in modifying applicable allotment management plans, annual operating plans, project decision documents and instructions to permittees to provide a high degree of assurance that objectives for conservation and restoration of anadromous fish habitat would be met.

The revision identified a set of key assumptions. One of the assumptions is that the goals or desired outcomes of management efforts provide the foundation for the recommended programmatic livestock grazing guidelines. The PACFISH EA was described as providing suitable riparian goals. All management activities should be structured so as not to prevent or meaningfully hinder accomplishment of the goals.

A summary of key priorities identified in the Enclosure B revision are:

- Maintain or allow for improvement of conditions where criteria for late seral ecological status are met or exceeded.
- Adjust management practices where the criteria for mid-seral ecological status are met but the trend is static or downward. This is especially important where vegetative factors are primarily responsible for the mid-seral rating.
- Adjust management practices where the criteria for early seral ecological status are met, with the understanding that deteriorated stream bank and channel conditions may not be recovered in the near term.

The Enclosure B revision stated that Al Winward, in Clary and Webster (1989) defined ecological status as a measure of the degree of similarity between current vegetation and potential vegetation for a given riparian area. Refined definitions for the three ecological classes were presented:

- Early seral. Percent similarity of riparian vegetation to the potential natural community/composition less than or equal to 25%; or, stream bank/channel condition rating “poor”.
- Mid-seral. Percent similarity of riparian vegetation to the potential natural community/composition 26-50% or better; and, stream bank/channel condition of at least “fair”.
- Late seral. Percent similarity of riparian vegetation to the potential natural community/composition greater than or equal to 50% or better; and, stream bank/channel condition rating “good” or better.

The MNF is utilizing Winward (2000) to evaluate ecological status of riparian vegetation, in place of the process described in Enclosure B. If similarity of riparian information is lacking, the Enclosure B revision suggested using PFC condition classes as a substitute.

3.2 MALHEUR NATIONAL FOREST RIPARIAN MONITORING STRATEGY

Many accepted methodologies and analytical tools are available to monitor short-term and long-term rangeland and forest health. The methods and tools chosen are dependent on the specific monitoring objectives as well as constraints such as timing, available funding and personnel, other priorities, and the geographical area to be monitored. Described below are the overall monitoring strategy, methods and analytical tools that the Malheur National Forest is currently using for determining condition and trend of riparian ecosystems as they relate to grazing activities. **The assessments and monitoring methods used are intended to be an important part of the adaptive management process and are subject to changes or modifications based on new scientific findings and improvements in methodologies as well as changes in definitions and policy. In particular, see Appendix L for a discussion of the monitoring protocol the Forest Service intends to use to evaluate compliance with bank alteration**

thresholds. Moreover, risk analyses and prioritization should generally be completed in all areas prior to initiating monitoring in order to determine the level and intensity of quantitative data collection. PFC assessments serve as the risk analyses/prioritization step.

Below are the key components of the MNF Riparian Monitoring Strategy:

1. Information Gathering and Interpretation
 - Proper Functioning Condition (PFC) Assessment –qualitative condition assessment over a stream reach (geomorphic or unit specific)
 - Multiple Indicator Monitoring (MIM) – quantitative monitoring protocol at designated DMAs
 - Analysis – interpretation and evaluation of assessment and monitoring information to determine current riparian condition and, to the extent feasible, trend)
 - Channel cross-section, streambed particle size distribution, and reach description measurements (i.e. Rosgen Channel Type)
 - Forest Service Region 6 Level II Stream Inventory Surveys – extensive quantitative assessment of stream channel, riparian vegetation, aquatic habitat condition, and biota to determine condition of selected stream systems
 - Spawning Surveys – quantitative assessment of redd vulnerability to disturbance
2. Support determinations of plan compliance -Provide information on which MNF can assess compliance with Forest Plan, including PACFISH & INFISH amendments. See Appendix C. for further discussion of Forest Plan standards and objectives related to riparian areas, water quality and fish habitat.
 - Standards are GM 1-4 in PACFISH & INFISH; standards 15-21 in Forest Plan (see Chapter IV).
 - Management Objectives for stream and riparian areas are described in PACFISH & INFISH amendments (RMO's) and in Amendment 29 of Forest Plan for MA3A/B (DFC's).
3. Recommendations
 - Shows linkage between condition, trend, and past/current management activities
 - A process that provides support for grazing management decisions or any necessary or appropriate adaptive management adjustments
 - Allows annual adjustment of management strategies, as needed, to achieve compliance with plan direction

Proper Functioning Condition Assessments

Proper functioning condition (PFC) assessments are a qualitative method for determining the condition of riparian areas. The term PFC is used to describe both the assessment process, and a defined, on-the-ground condition of a riparian area. PFC assessments can be an appropriate starting point for determining and prioritizing the type and location of quantitative inventory or monitoring necessities, and has been proven to be an excellent communication tool for bringing a wide diversity of publics to agreement. All PFC assessments are to be conducted with a journey level interdisciplinary team. One purpose of these assessments is to help correlate the findings with the trend towards attainment of the Malheur Forest Plan Riparian Management Objectives (RMOs), more specifically, to determine whether grazing practices are retarding attainment of

Near Natural Rates of Recovery of RMOs. No PFC assessments have been conducted in these 5 allotments.

Multiple Indicator Monitoring

The July 1, 2003 PACFISH/INFISH Implementation Monitoring Program Manual provides the background and direction for monitoring. The Multiple Indicator Monitoring (MIM) supplement by Cowley/Burton, dated May 2005 with addendums, provides the procedures in use by the MNF to monitor stream banks and riparian vegetation. The Interagency Implementation Team created the above documents; see Appendix E for these documents. The authors of MIM recently issued a 2011 technical guide as well (Burton et. al. 2011). MIM for grazing activities is designed to determine whether or not livestock grazing management is resulting in “Near Natural Rates of Recovery” as defined by PACFISH/INFISH. Below are the three components, which comprise MIM. Monitoring is to be conducted by an interdisciplinary professional team trained in riparian plant identification and channel classification. Multiple indicator monitoring consists of implementation (endpoint indicator) monitoring and effectiveness (riparian objective) monitoring at designated locations (i.e. designated monitoring areas).

Designated Monitoring Areas

Designated Monitoring Areas (DMAs) are the locations in riparian areas and along streambanks where quantitative monitoring takes place. They are monitored to provide information concerning the management of critical areas. Essentially DMA selection relies on the theory that if proper management occurs in that location, proper management will be occurring throughout the rest of the management unit. See Appendix D for the procedures used to collaboratively establish DMAs. The goal is to establish more DMAs each grazing season in order to establish a 5-year re-monitoring schedule and have coverage across the Forest’s allotments.

Implementation Monitoring - Endpoint Indicators

Implementation (endpoint indicator) monitoring measures indicators to determine if the authorized livestock grazing strategy for a particular season has had the projected effects to resources that the MNF has anticipated in developing the strategy and to determine if adaptive management adjustments need to be made for the following season(s). It provides information to assist with making decisions under adaptive management. Presently, implementation monitoring includes: modified extensive browse utilization (Interagency Technical References, 1996), modified stubble height (Interagency Technical Reference, 1996 and Challis Resource Area, 1999), and streambank alteration (Cowley 2002). These procedures provide information to refine and make annual adjustments to livestock grazing management practices necessary to meet long-term management objectives (adaptive management). They can be used as early warning indicators that current grazing impacts may prevent the achievement of management objectives and can also be used to help explain changes in riparian vegetation and channel conditions over time. See web-site (<http://www.rmsmim.com/>) for sampling procedures used.

Effectiveness Monitoring - Riparian Objectives

Effectiveness (riparian objective) monitoring is designed to address the question of whether or not management practices currently applied to the area are achieving the desired results. These procedures are designed to assess the current condition and measure changes in streambanks, channels, and streamside vegetation over time, i.e., trend. They help determine if local livestock grazing management strategies and other land management actions are making progress toward

achieving the long-term goals and objectives for streamside riparian vegetation and aquatic resources. The goal is to conduct effectiveness monitoring every three to five years on riparian areas and streambanks. This period of time is considered to be the minimum necessary to detect changes, although unusually wet years and/or flood events may result in short-term changes that validate the need to monitor more frequently, or at least at the time of the event. Budget and personnel constraints may limit the extent in which monitoring of this type will be conducted. Presently, effectiveness monitoring includes: modified greenline composition (adapted from Winward 2000 and USDI BLM 1996a), woody species height class (Kershner et al. 2004), streambank stability and cover (adapted from Kershner et al. 2004), woody species age class (adapted from Winward 2000), greenline-to-greenline width (Burton et al. 2008), substrate (Bunte and Apt 2001), and residual pool depth and pool frequency (Lisle 1987). These provide data and information concerning the present conditions and trend of riparian vegetation, channels, and streambanks, and to help determine if aquatic systems are being degraded, maintained, or restored across the Malheur National Forest. See web-site (<http://www.fs.fed.us/biology/fishecology/emp/>) for PIBO data and sampling procedures used.

Forest Service Region 6 Level II Stream Inventory Surveys

Forest Service Region 6 Level II Stream Inventory Surveys generate comparable baseline information on conditions of fish-bearing streams to support a variety of management activities. As inventories are completed and repeated over time, the information generated by them can be useful in measuring changes in stream channel conditions and determining attainment of habitat management objectives. The Level II inventory generates quantitative measurements and estimates of channel conditions and habitat attributes, including core attributes of streamflow, temperature, substrate composition, width/depth ratio, channel length and sinuosity, gradient, pool frequency, large wood, bank stability, and special habitats. Numerous non-core optional attributes may also be evaluated based on Forest needs, such as stream shading and overstory/understory vegetation. The Forest goal is to inventory 10 percent of fish-bearing streams per year, inferring a 10-year re-inventory recurrence interval. The 2010 Region 6 Stream Inventory Handbook can be found at:

(<http://www.fs.fed.us/r6/water/fhr/sida/handbook/Stream-Inv-2010.pdf>)

Spawning Surveys

Spawning surveys are a quantitative method to assess steelhead redd presence and vulnerability to livestock disturbance and may also be used to assess compliance with the level of “take” authorized within a Biological Opinion. The Forest has developed a strategy to avoid redd trampling “take” of steelhead and bull trout (see Appendix F).

Uplands Monitoring

Beginning in the 1930s, permanent camera points were established on the Malheur National Forest. Their purpose was to monitor the effects of management on the resources of the Forest (Fifty Years of Change on the Range, R6-Mal-035-89). Many camera points have been re-photographed a number of times. This monitoring will continue.

In the 1950s and early 1960s Parker Three-Step C&T (Condition & Trend) Transects were installed throughout the Forest. The majority of these were established in the uplands. Over the last five years some of these transects have been re-examined. The procedure has been to read the transect using the original Three-Step method and then reread the transect using a modified

Daubenmire cover/frequency method (see Technical Reference 1734-4). This allows comparisons between old and new information to determine ecological condition and trend and establishes a baseline using the more accurate cover/frequency method for gathering future data. The re-examining of these established transects will continue. If new trend transects are established the modified Daubenmire cover/frequency will be used.

There are a variety of additional or other monitoring methods available for use. The method or methods to be used will depend on the questions needing to be answered and considering other priorities. In some cases ocular observation (qualitative) will be sufficient to measure utilization, but when specific concerns are identified the forest may need quantitative methods such as Paired Clipped plots or development and use of height/weight curves may be necessary. Some of the more commonly used methods can be found in “*Utilization Studies and Residual Measurements*” (Interagency Technical Reference 1996, BLM/RS/ST-96/004+1730). (Please refer to ***Malheur National Forest Range Monitoring Guidelines***, October 16, 2006 for additional accepted methodology)

All of the monitoring methods used by the Forest are also intended to facilitate communications between forest range and resource personnel, grazing permittees and consulting agency personnel. This will largely be accomplished through participation and one-on-one interaction during the interdisciplinary, on the ground implementation.

4 PROPOSED ACTION AND ESA ACTION AREA

The MNF used the LRMP direction and policies presented in Section 3 to design the 2012-2016 proposed grazing action for Seneca, Deadhorse, Hanscomb, and Fields Peak Allotments. Public laws such as Clean Water Act were considered. The development of the project design criteria, grazing end-points and grazing strategies for these five allotments considered the PACFISH RMOs and Grazing Management Standards, desired conditions and standards from amendment 29 to the MNF LRMP, PACFISH livestock grazing guidelines (Enclosure B) and MSRAs (appendix G). Forest policies on Riparian Monitoring and MSRAs were also informed by the LRMP direction and Clean Water Act. Examples of resource objectives and their sources include:

- Greenline successional status value of at least 61, indicating late seral or the current value, whichever is greatest (Winward 2000, Burton et al. 2008) was developed in response to the PACFISH Enclosure B revision;
- Woody species regeneration sufficient to develop and maintain healthy woody plant communities (diversity of age and structure classes) was developed in response to MNF LRMP Amendment 29;
- Bank stability criteria (80% or current value, whichever is greatest for non-priority watersheds; at least 90% or current value, whichever is greatest, for priority watersheds) were from PACFISH;
- Water temperature criteria are from the MNF LRMP; and
- Width-depth ratio criteria are from PACFISH

4.1 PROPOSED ACTION

Seneca Allotment

The Seneca allotment is located south of the town of John Day on the north edge of Bear Valley mostly within Township 15, Ranges 30 and 31E. The Allotment encompasses approximately 10,166 acres and is divided into 4 pastures: Vance Creek, Camp Creek, Camp Creek Management, and Koehler (Figure 1).

Deadhorse On/Off Allotment

The Deadhorse allotment is located southwest of the town of John Day mostly within T 14 and 15 S, and R 29 and 30 E. The Allotment includes approximately 15,527 acres and is divided into 4 pastures: North, Riley, Riley Meadows, and Percival (Figure 3).

Hanscomb Allotment

The Hanscomb allotment is located southwest of the town of John Day, northeast of Bear Valley mostly within T 14 and 15 S and R 30 E. The Allotment includes approximately 9,102 acres and is divided into 4 pastures: Laycock, Upper Geary, Geary, and Allen/Morris (Figure 4).

Fields Peak Allotment

The Fields Peak allotment is located southeast of the town of John Day mostly within T 14 and 15 S and R 28, 29, and 30 E. The Allotment includes approximately 30,718 acres and is divided into 4 pastures: Fields Peak, Tex Creek, Miners Creek, North Murderers Creek, and Murderers Creek (Figure 5).

4.1.1 PERMIT INFORMATION AND GRAZING SYSTEMS

The Seneca, Deadhorse, Hanscomb, and Fields Peak Allotments are managed under one grazing permit. Permit information is displayed in Table 8 and Table 9.

TABLE 8. PERMIT INFORMATION FOR THE SENECA, DEADHORSE, HANSCOMB, FIELDS PEAK ALLOTMENT

Permit Number	Permit Issuance Date	Permit Expiration Date
01845	6/01/2007	12/31/2016

TABLE 9. PERMIT INFORMATION - ALLOTMENT SPECIFIC

Allotment	Permitted Livestock (Cow/Calf Pairs) / AUMs	On Date	Off Date
Seneca	170 / 1018	6/15	10/30
Deadhorse	175 / 1040	6/1	10/15
Hanscomb	121 / 719	6/1	10/15

Fields Peak	240 / 1437	6/15	10/15
Fields Peak	2 (horses) / 11	6/1	10/15

Grazing System:

- The Seneca Allotment will be managed using a deferred rest-rotation grazing strategy.
- The Deadhorse, Hanscomb, Fields Peak Allotments will be managed as one allotment utilizing a deferred rest-rotation grazing strategy.
- Range Readiness and utilization levels may result in varying on/off dates and pasture rotations within the parameters of authorized use.

4.1.2 PROJECT DESIGN CRITERIA

The following project design criteria (PDC) will be used to minimize or eliminate adverse effects of the PEs on MCR Steelhead and designated CH. The MNF regards these PDC as integral components of the proposed action and expects that all proposed project activities will be completed consistent with those measures.

1. Management will be framed in a manner that will allow managers to manipulate grazing strategies (dates, stocking levels, rotational patterns) depending on annual environmental factors and permittee success at meeting standards during the previous year.
2. Permittees must maintain perimeter and interior fences prior to turn-out.
3. Standards that are required of the permittee (e.g., turn on dates, move triggers, end point standards) will be outlined in an addendum to Part III of the grazing permit.
4. MSRA will be located and used to identify stream sections that are most vulnerable to livestock impacts. Identifying MSRA locations will guide application of bank alteration values.
5. Spawning surveys will occur within all pastures containing MSRA's where turn out is expected to occur prior to June 30. Of the remaining CH reaches 20% will be randomly surveyed for redds where turn out is prior to June 30 (See Appendix F. Strategy to minimize Redd Trampling "Take" of Steelhead and Bull trout).
6. Where there is significant risk for redd trampling, the Forest and permittees will utilize a number of tools to protect redds, which include but are not limited to: alternative rotation, rest, exclusion with water gaps, temporary electric fences, additional riding. Specific methods selected will be determined collaboratively with the permittees on a pasture-specific basis.
7. Complete all required monitoring at PIBO Effectiveness Monitoring DMAs. DMA's to be monitored are provided to the Forest yearly by the EM Team via the Regional Office. This will effectively satisfy Interagency Implementation Team (IIT) monitoring requirements.
8. The Forest Service will visually inspect riparian livestock use in each pasture containing steelhead CH near the mid-point of the grazing rotation for that pasture, and will conduct applicable Multiple Indicator Monitoring ("MIM") on any such pasture where it appears that riparian conditions are approaching one or more move triggers or end-point indicators. This will help meet our long term riparian resource objectives.

9. Annual use indicators will dictate when livestock are moved between units or off the allotment, within the terms of the term grazing permit, including moves in response to fish spawning. This will help the Forest Service meet its long term riparian resource objectives.
10. The Forest Service will provide the Services with an End of Year Grazing Report by March 1 of each year.
11. Use of roads and off-road travel by permittees and staff will follow these PDC:
 - a. Vehicles are not authorized to travel through seeps, springs or streams except for use of existing fords on road crossings;
 - b. All refueling activities and fuel storage will occur at least 150 feet away from live streams;
 - c. OHV routes within 100 feet of streams will be camouflaged so that access routes do not become new trails and minimize disturbance to riparian vegetation;
 - d. OHV travel off established roads within 100 feet of streams would occur only during periods when soil is dry.

4.1.3 GRAZING USE INDICATORS AND SUPPORTING RATIONALE

The Forest Service's Regions 1, 4, 6 and Bureau of Land Management's Idaho, Montana, Nevada, Oregon and Washington have made commitments through the PACFISH and INFISH Management Strategies to protect and improve aquatic resources found in the interior Columbia River basin. Since the Forest Service Pacific Northwest Region (Oregon and Washington) which includes the Malheur National Forest began implementing these strategies, there has been marked improvement in management of aquatic resources. To strengthen the implementation of the aquatic strategies, an interagency group consisting of deputies from the various action and regulatory agencies (Deputy Team) was formed to provide oversight of the strategies and subsequent biological opinions commonly known as PIBO. Under the Deputy Team oversight, an implementation monitoring module was developed for livestock grazing, and its application is required where listed fish species occur in the interior Columbia River basin. Compliance with these requirements is monitored and presented to the Deputy Team during their annual reviews.

The PNW region requires application of the PIBO implementation and effectiveness monitoring program for National Forest LRMPs amended by PACFISH and INFISH, and the regional office annually coordinates the PIBO monitoring programs with the National Forests with listed fish species. FS line officers continue to work with their staffs and grazing permittees to ensure that implementation monitoring requirements are met. As described in the PIBO monitoring strategy and the annual regional coordination letter, the Forest established Designated Monitoring Areas (DMAs) and annually monitors the grazing use indicators at these PIBO DMAs as well as DMAs established by the MNF.

Data collected at the PIBO effectiveness monitoring DMAs reflect the grazing use indicators applicable to stream banks and stream channels. Accordingly, the regional monitoring coordination letter identifies the following requirements will apply to these DMAs:

- A. Measurements will be on the greenline (first perennial vegetation above the channel). Measurements must include, at minimum: 1) bank alteration and 2) stubble height if any herbaceous vegetation is present.

- B. Where woody riparian vegetation dominates the DMA with little or no herbaceous vegetation along the greenline, woody use (browse) should be measured and may be sampled in lieu of stubble height.
- C. These measures will be made using the current MIM protocol.

Therefore, the grazing use indicators required by the PIBO and used by the Forest riparian monitoring programs at all DMAs are: browse of woody vegetation, stubble height of greenline vegetation and streambank alteration. Woody vegetation browse is used to regulate impacts on woody recruitment to streams, greenline stubble height is used to regulate grazing impacts on greenline ecological status and streambank alteration is used to regulate grazing impacts on streambank stability and channel width. For consistency with the PIBO monitoring program and regional direction regarding coordination with it, the Forest elected to use the current MIM for their monitoring protocol.

The MNF utilizes move trigger and endpoint (annual) indicators to manage livestock. The underlying concept behind the use of end-point indicators for livestock grazing management is that the selected end-points, if not exceeded, will allow for the attainment of or reasonable progress to be made toward desired conditions for riparian areas and fish habitat as described in Section 3 – Malheur NF LRMP and Section 4 – Proposed Action.

The MNF developed values for livestock move trigger and annual endpoint indicators (Table 6). The ranges of values are starting points based on research and the MNF’s best collective professional judgment for establishing desired riparian conditions. The end-point indicator values (allowable use in riparian areas) are, to the extent feasible and appropriate data are available, be site-specifically designed to prevent any meaningful carry-over effects. They also provide for the evaluation of management practices to determine if they are effective in maintaining the desired and/or proper functioning condition, or improving the structure and function of riparian and aquatic conditions. These values could be adjusted as more site-specific information is gathered. End-point indicators (allowable use in riparian areas) should be adjusted for timing, intensity, frequency, and duration. The rationale for the development of the move trigger and end-point grazing use indicators is discussed below.

Livestock grazing along the greenline of stream channels will be limited to attain the numeric move trigger and end-point indicator values in Table 10.

TABLE 10. SENECA ALLOTMENT LIVESTOCK MOVE TRIGGER AND END-POINT INDICATORS BY PASTURE.

Pasture Name/ DMA or Key area Name / Creek Name	Monitoring Attribute	Key Species	Move Trigger ²	Endpoint Indicator
Vance Creek, DMA site yet to be determined	Browse use		40% 30%	50% (early season use) 40% (late season use)
MSRA-None	Greenline stubble ¹	Deep-rooted hydric spp.	4 in.	4 in. (early season use) 6 in. (late season use)

Pasture Name/ DMA or Key area Name / Creek Name	Monitoring Attribute	Key Species	Move Trigger ²	Endpoint Indicator
	Streambank Alteration		15%	20%
Upland Sites (All Pastures)	% Utilization	Upland grass species	35%	45%
Riparian Areas (All Pastures)	% Utilization	Riparian grass species	35%	45%
<p>¹Browse use and greenline stubble will be used until next trend reading is completed to determine which attribute will be best suited to attain long term objectives.</p> <p>²The move triggers identified in this document are not intended to be used as a “standard.” They are designed to function as a tool to help permittees successfully meet allowable use standards. The move trigger values are set at lower levels than the endpoint indicators to serve as a trigger point for permittees to begin gathering and moving livestock to the next scheduled pasture. Meeting move triggers is not a requirement of the term grazing permit as are the endpoint indicators.</p>				

TABLE 11. DEADHORSE ALLOTMENT LIVESTOCK MOVE TRIGGER AND END-POINT INDICATORS BY PASTURE.

Pasture Name/ DMA or Key area Name / Creek Name	Monitoring Attribute	Key Species	Move Trigger ²	Endpoint Indicator
North, Riley Creek DMA #1, Riley Creek MSRA-None	Browse use		40% 30%	50% (early season use) 40% (late season use)
	Greenline stubble ¹	Deep-rooted hydric spp.	4 in.	4 in. (early season use) 6 in. (late season use)
	Streambank Alteration		15%	20%
Upland Sites (All Pastures)	% Utilization	Upland grass species	35%	45%
Riparian Areas (All Pastures)	% Utilization	Riparian grass	35%	45%

Pasture Name/ DMA or Key area Name / Creek Name	Monitoring Attribute	Key Species	Move Trigger ²	Endpoint Indicator
		species		
<p>¹Browse use and greenline stubble will be used until next trend reading is completed to determine which attribute will be best suited to attain long term objectives.</p> <p>²The move triggers identified in this document are not intended to be used as a “standard.” They are designed to function as a tool to help permittees successfully meet allowable use standards. The move trigger values are set at lower levels than the endpoint indicators to serve as a trigger point for permittees to begin gathering and moving livestock to the next scheduled pasture. Meeting move triggers is not a requirement of the term grazing permit as are the endpoint indicators.</p>				

TABLE 12. HANSCOMB ALLOTMENT LIVESTOCK MOVE TRIGGER AND END-POINT INDICATORS BY PASTURE.

Pasture Name/ DMA or Key area Name / Creek Name	Monitoring Attribute	Key Species	Move Trigger ²	Endpoint Indicator
Laycock, Laycock Creek DMA, Laycock Creek MSRA-Present	Browse use		40% 30%	50% (early season use) 40% (late season use)
	Greenline stubble ¹	Deep-rooted hydric spp.	4 in.	4 in. (early season use) 6 in. (late season use)
	Streambank Alteration		10%	15%
Upland Sites (All Pastures)	% Utilization	Upland grass species	35%	45%
Riparian Areas (All Pastures)	% Utilization	Riparian grass species	35%	45%
<p>¹Browse use and greenline stubble will be used until next trend reading is completed to determine which attribute will be best suited to attain long term objectives.</p> <p>²The move triggers identified in this document are not intended to be used as a “standard.” They are designed to function as a tool to help permittees successfully meet allowable use standards. The move trigger values are set at lower levels than the endpoint indicators to serve as a trigger point for permittees to begin gathering and moving livestock to the next scheduled pasture. Meeting move triggers is not a requirement of the term grazing permit as are the endpoint indicators.</p>				

TABLE 13. FIELDS PEAK ALLOTMENT LIVESTOCK MOVE TRIGGER AND END-POINT INDICATORS BY PASTURE.

Pasture Name/ DMA or Key area Name / Creek Name	Monitoring Attribute	Key Species	Move Trigger ²	Endpoint Indicator
Fields Peak, Fields Creek PIBO DMA, Fields Creek MSRA-Present	Browse use		40% 30%	50% (early season use) 40% (late season use)
	Greenline stubble ¹	Deep-rooted hydric spp.	4 in.	4 in. (early season use) 6 in. (late season use)
	Streambank Alteration		10%	15%
Tex Creek, DMA site yet to be determined, Tex Creek MSRA-Present	Browse use		40% 30%	50% (early season use) 40% (late season use)
	Greenline stubble ¹	Deep-rooted hydric spp.	4 in.	4 in. (early season use) 6 in. (late season use)
	Streambank Alteration		10%	15%
Miners Creek, DMA site yet to be determined, Miners Creek MSRA-Present	Browse use		40% 30%	50% (early season use) 40% (late season use)
	Greenline stubble ¹	Deep-rooted hydric spp.	4 in.	4 in. (early season use) 6 in. (late season use)
	Streambank Alteration		10%	15%
North Murderers Creek, DMA site yet to be determined MSRA-None	Browse use		40% 30%	50% (early season use) 40% (late season use)
	Greenline stubble ¹	Deep-rooted hydric spp.	4 in.	4 in. (early season use) 6 in. (late season use)
	Streambank		15%	20%

Pasture Name/ DMA or Key area Name / Creek Name	Monitoring Attribute	Key Species	Move Trigger ²	Endpoint Indicator
	Alteration			
Murderers Creek, Murderers Creek DMA #2, Murderers Creek MSRA-Present	Browse use		40% 30%	50% (early season use) 40% (late season use)
	Greenline stubble ¹	Deep-rooted hydric spp.	4 in.	4 in. (early season use) 6 in. (late season use)
	Streambank Alteration		10%	15%
Upland Sites (All Pastures)	% Utilization	Upland grass species	35%	45%
Riparian Areas (All Pastures)	% Utilization	Riparian grass species	35%	45%
<p>¹Browse use and greenline stubble will be used until next trend reading is completed to determine which attribute will be best suited to attain long term objectives.</p> <p>²The move triggers identified in this document are not intended to be used as a “standard.” They are designed to function as a tool to help permittees successfully meet allowable use standards. The move trigger values are set at lower levels than the endpoint indicators to serve as a trigger point for permittees to begin gathering and moving livestock to the next scheduled pasture. Meeting move triggers is not a requirement of the term grazing permit as are the endpoint indicators.</p>				

Rationale to Support the Range of Initial Values for Selected End-Point Indicators/Condition Thresholds/Allowable Use Criteria

Stubble Height: 4-6 inches. Stubble height has been identified as being related to the physiological health and vigor of individual plants/communities as well as the ability of vegetation to protect streambanks and filter during overbank flows, although by itself it is generally not sufficient to establish a relationship between grazing and riparian vegetative conditions. Research is limited, but the literature generally suggests 4-6 inches of residual stubble height allows for improved riparian grazing management and provides for adequate riparian protection. Clary and Leininger (2000) conducted studies on stubble height and its ability to improve riparian habitats and to capture and stabilize sediment. They concluded that stubble heights of 4-6 inches appear to stabilize the greatest amount of sediment. Clary (1999) states that by maintaining stubble heights of 4-5.5 inches allowed for streambank recovery. End-point indicator values are intended to vary by site depending on similarity to desired conditions and the resiliency of the site being monitored (University of Idaho Stubble Height Study Team

2004, Clary and Leininger 2000, Clary et al. 1996, Hall and Bryant 1995, Appendix B - PACFISH Enclosure B, Clary and Webster 1989). Stubble height is an annual use indicator that should be used in combination with long-term monitoring of vegetation and stream channel attributes.

Bank Alteration: 10-30%. In general, the most widespread impact livestock have on riparian areas is trampling stream banks (Bengeyfield, 2006). Like stubble height, streambank alteration is another annual or short-term indicator used to evaluate the potential effects of livestock grazing in riparian areas, primarily evaluating potential effects to long-term streambank stability and channel shape. It is used as a tool to assess the intensity of grazing along streambanks and to determine when such intensity may be appropriate or deemed excessive. It can also prove useful in determining the cause-and-effect relationships between livestock grazing and stream channel conditions and whether management changes are needed for the following year. Streams are naturally dynamic and have the ability to repair a certain amount of annual disturbance each year (the amount is variable based upon stream gradient, substrate composition, streambank materials, vegetation type and abundance, channel geometry, flow regime, etc.). Again, although the literature is not extensive, it generally suggests that 10-30% of annual streambank alteration is consistent with providing adequate riparian protection, and is intended to vary by site depending on similarity to desired conditions and the resiliency of the site being monitored (Burton et al. 2011, Heitke et al. 2008, Bengeyfield 2006, Cowley 2002, Bengeyfield and Svoboda 1998). Bengeyfield (2006) found that when streambank alteration measured 15-20%, width to depth ratios showed an improving trend. He also noted that the vegetation improvements kept pace with the physical changes.

The streambank alteration procedure described here is an intercept procedure recording presence/absence of current year's disturbance along the greenline. It is not a measure of the percent of the area of streambank altered, but rather an estimate of the percent of the length of bank altered along the greenline based on the presence or absence of a hoofprint(s) intercepting one (or more) of the five lines within a plot. This procedure samples only that part of the streambank associated with the greenline, often at the top of the streambank, and only within a 42-by-50cm plot. The streambank may be wider or narrower than the width of the plot.

Streambank alteration is an annual use indicator that should be used in combination with long-term monitoring of streambank stability and channel geometry. In addition, it is worth noting that research is continuing to be conducted on the various ways that can be used to monitor for and measure actual streambank alteration (including MIM, which the District is presently using) to account for accuracy of results, reduction of variability among observers, and the resources necessary to carry out such measures.

Mean incidence of use on woody species: 30-50%. Woody vegetation is an important component of many stream/riparian ecosystems as it can provide a strong root system, filter sediment, and provide stream shade and habitat diversity. Woody species browse is a short-term indicator of grazing utilization of woody species. Overall, there is generally a reduction in seed production of woody plants that receive more than 55 percent utilization, and when heavy and severe utilization levels are sustained over time overall plant health, including size and root strength, is reduced. Although the literature is not extensive, it generally suggests light to moderate allowable use on woody species (~30-50%) can be sustained and not meaningfully impede the potential for improved conditions of affected woody plant communities; and is intended to vary by site depending on similarity to desired conditions and the resiliency of the

site being monitored (Winward 2000, USDI BLM 1996, Appendix B-PACFISH Enclosure B). Woody species browse is an annual use indicator that should be used in combination with the long-term monitoring indicators of woody species age class and greenline composition to help determine the health of woody plant communities.

Livestock grazing along the greenline of stream channels will be limited to attain the numeric move trigger and end-point indicator values in Table 6. The numeric values in Table 6 are considered starting points for allowable use since values could be adjusted as more site-specific information is gathered.

Initial Values for Grazing Use Indicators

Based on the best available science, applied science publications, and professional judgment, the Forest interdisciplinary team selected initial values for each indicator. The season of use determined the initial values of endpoint indicators for woody shrub use and stubble height of greenline vegetation. The early season initial values for shrub use and stubble height are 50% and 4 inches, and the late season initial values for shrub use and stubble height are 40% and 6 inches, respectively. Grazing use in the early season allows time for vegetation growth after livestock use. The exact dates and times of early and late season can vary across the Forest and between given years, and therefore are not specified. However, to provide some typical guidelines, early season is usually defined as the beginning of the growing season to mid-July and late season from mid-August to the end of the growing season.

To determine an initial value for the streambank alteration grazing use indicator, the Forest also looked at a Regional Technical Team (RTT) report prepared under the Streamlining Consultation Procedures resolution process. The NMFS had stated that “The best available science indicates that the 10% and 20% bank alteration levels are appropriate in preventing bank destabilization and protecting habitats critical to listed fish.” Their position paper cited numerous references to support these values. The RTT reviewed documents cited by NMFS, and they concluded that NMFS had reasonably established a causal link between streambank alteration-related habitat effects caused by livestock grazing activity and the taking of the species (i.e., grazing will affect stream channel conditions that will affect fish habitat conditions such as water quality, food, cover, etc.). The literature generally supports the concept that increased streambank alteration will, at some point, adversely affect stream channel conditions, and therefore fish habitat conditions. However, there was uncertainty relative to the percentage of streambank alteration at which habitat conditions were significantly altered and take of the species is likely to occur.

There is little field research supporting any specific percent streambank alteration standard using a defined and repeatable measurement protocol. The above RTT report and citations provide recommendations and professional judgments that range from 10% to 30% streambank alteration, but do not present empirical evidence from grazing monitoring data to support the percentages. The Forest interdisciplinary team recognizes a connection between the streambank alteration grazing use indicator and long term fish habitat conservation objectives in the LRMP, but couldn't determine a consensus value. Therefore, the Forest interdisciplinary team selected an initial value of 20% streambank alteration for endpoint grazing use indicator which is the statistical median of the range.

Adjustments to Values of Grazing Use Indicators and/or Grazing Strategy

The interdisciplinary team considers available information on riparian condition (eg. succession status of greenline vegetation and woody species regeneration) and presence of MSRAs to adjust values of grazing use indicators and/or the grazing strategy. Information wasn't available for the successional status of the greenline and woody species regeneration for the pastures in this allotment. Therefore, it didn't affect the values of the grazing use indicators in Table 6. However, the presence of MSRAs resulted in reductions of the streambank alteration values for several pastures.

Criteria used to evaluate the riparian condition (eg. succession status of greenline vegetation and woody species regeneration):

When these conditions apply:

- Greenline plant communities show moderate to high similarity to desired condition class/seral stage -- greenline successional status value is 41% or greater (mid-late seral) as defined by Winward (2000); *and/or*
- Stream/riparian systems have been assessed as being in Properly Functioning Condition or Functioning-at-Risk (high to moderate) category (Prichard et al. 1998); *and/or*
- Riparian/channel attributes are near desired conditions in a Unit,

Then allowable use within riparian areas will be:

1. <40-50% browse on both clumped, multi-stemmed species (i.e. most willows (*Salix spp.*) and single-stemmed species (i.e. coyote willow (*Salix exigua*), birch (*Betula spp.*), alder (*Alnus spp.*), cottonwood or quaking aspen (*Populus spp.*)) i.e. <50% browse if early season use, and <40% browse for mid and late season use⁴; and
2. >4-6 inch residual stubble height (will vary based on greenline successional status/seral stage, and season of use) i.e. >4 inches if early season use, and >6 inches for mid and late season use; and
3. Allowable bank alteration will be limited to 20% streambank alteration⁵ by large herbivores (% of linear length of greenline altered) (Multiple Indicator Monitoring (MIM) of Stream Channels and Streamside Vegetation (Interagency Technical Reference 1737-23 2010)⁶. The estimated combined variability, observer error and sampling error or sample size, results in a 95% confidence interval of 6% for this bank alteration monitoring method. Thus, by setting a trigger for moving livestock at

⁴ The exact dates and times of “early”, “mid” and “late” can vary across the Forest and between given years, and therefore are not specified. However, to provide some typical guidelines, “early” is usually defined as the beginning of the growing season to mid-July, “mid” season from mid-July to mid-August, and “late” season from mid-August to the end of the growing season.

⁵ The allowable level of bank alteration for a specific site should allow for no more than 5% of the lineal bank distance (includes both sides of the stream) displaying evidence of new bank instability that has become perceptible after livestock grazing is initiated in a pasture. Note: hoof prints by themselves are not a sign of instability unless they move the bank by > 10 cm (direct shearing or sloughing of the bank).

⁶ Research is presently ongoing, which may result in a new and or modified method of measuring Bank Alteration. If in fact this occurs the PIBO EM Team and or other researchers would present findings and provide a cross walk and rational to the existing monitoring method and endpoint indicators.

~14%, we can be reasonably confident that livestock would be off the pasture before an additional 12% alteration was reached. The upper level for reasonable confidence would be 26% -- which represent an upper limit for the associated conservation measure.

However, when these conditions apply:

- When greenline plant communities show low similarity to desired condition class/seral stage -- greenline successional status value is less than 41 (early seral) as defined by Winward (2000); and/or
- Stream/riparian systems have been assessed as being in a *Functional-at-Risk (low) to Non-Functional category* (Prichard et al. 1998) {a Non-Functional system is one that clearly does not provide adequate vegetation, landform, or large woody debris to dissipate stream energy associated with high flows and thus are not reducing erosion, improving water quality, etc.; there is an absence of certain physical attributes such as a floodplain where one should be} in a Unit,

Then,

1. Consider resting the area/s for one or more years until condition reaches moderate similarity for those riparian areas with moderate and low gradient channels, such as Rosgen "B" and "C" channel types, with substrates composed of medium to fine easily eroded materials; or
2. If grazing is allowed, use should be for only short duration (i.e. to facilitate moves, etc.) and during a period of least environmental impacts

Most Sensitive Riparian Areas:

The MNF has identified stream reaches of high quality steelhead spawning and rearing critical habitat called Most Sensitive Riparian Areas (MSRAs). The process and criteria for identifying MSRAs is described in detail in Appendix G- Methods for determining Most Sensitive Riparian Areas in relation to Mid Columbia River Steelhead. The miles of MSRA by stream and location in the allotment are displayed in Table 3, 4, 5, and 6 and Figures 1, 2,3, 4, and 5 respectively.

MSRAs are typically steelhead critical habitat that is most accessible and sensitive to livestock use. MSRA and the grazing strategies described below are part of the proposed action. Certain grazing strategies can be used to minimize livestock and stream interactions and promote maintenance of, or recovery towards, desired conditions. Pastures containing MSRA that include one or more of the following grazing strategies would result in allowable use levels of 20% bank alteration versus the more restrictive standard of 15%:

- Rest Rotation – 1 year of complete rest during a grazing cycle (grazing cycle is typically 3-4 pastures)
- Double Rest Rotation – pasture is rested for two consecutive years, then grazed either early or late to following year depending on recovery needs (i.e. herbaceous or shrubs)
- Corridor Fencing – complete rest from grazing for a specified period of time or until specified objectives are met.

If at any point during this consultation a permittee adopts one or more of the strategies listed above the endpoint indicator would be adjusted to reflect such management changes in the annual instructions.

Other useful tools to minimize riparian use include – using a full-time rider (7 days/week), using electric fence, using low-stress stockmanship, placing low-moisture nutrient supplement blocks (as well as using other supplementations) in uplands, less than 21 days grazing duration in any pasture during the hot season (typically mid and late seasons, or mid-July to end of growing season). The use of these tools will be evaluated by the IDT on an annual basis to determine if the level of allowable use would be raised to 20% bank alteration. If none of these tools are in place, allowable use will remain at the 15% bank alteration.

4.1.4 MONITORING

The Malheur National Forest monitoring strategy for determining condition and trend of riparian ecosystems as they relate to grazing activities was described in detail in Section 3.2. The goal is to determine site-specific desired riparian/stream channel conditions and the levels of allowable use (annual indicators also known as end-points) that will improve conditions that are not at the desired and/or proper functioning condition. The assessments and monitoring protocols used, as well as the values for desired conditions and allowable use, are intended to be an important part of the adaptive management process and are subject to changes or modifications based on new scientific findings and improvements in methodologies as well as changes in definitions and policy.

The annual indicators are used in implementation monitoring to ensure that grazing does not prevent the attainment of the desired conditions. Riparian annual use indicators used on the Malheur National Forest include greenline stubble height, bank alteration, and woody browse. Greenline stubble height is used to regulate grazing impacts on greenline ecological status, bank alteration is used to regulate grazing impacts on bank stability, and woody browse is used to regulate impacts on woody recruitment. The specific indicators selected for a specific unit should be those that correspond with the riparian resources that are most sensitive to the impacts of livestock grazing. For example, if bank stability was the riparian feature most likely to be impacted by livestock grazing in a unit, then bank alteration would be selected as the annual use indicator for that unit.

Annual use indicators will be measured at key areas by key species (on uplands) and at DMA greenlines annually. Key areas are monitoring sites chosen to reflect the effects of grazing over a larger area (Burton et al. 2008). Key species are preferred by livestock and an important component of a plant community, serving as an indicator of change (Coulloudon et al 1999). The Interagency Technical Reference or other best available science would be used to monitor grazing use. The MIM Interagency Technical Bulletin (Burton et al 2008) or other best available science would be used to monitor grazing use at DMAs. The Forest Service will monitor annual use indicators. Triggers will be used by permittees as a tool to help ensure annual use indicators are met. Endpoint indicators will be monitored by MNF personnel at designated monitoring areas (DMAs) at the end of the grazing period, following the MIM protocol (Burton et al. 2011). Move trigger evaluations will be conducted by the permittee.

Annual use indicators will be adjusted if desired conditions are not being met. Effectiveness (riparian objective) monitoring is designed to address the question of whether or not

management practices currently applied to the area are achieving the desired results. These procedures are designed to assess the current condition and measure changes in streambanks, channels, and streamside vegetation over time, i.e., trend. They help determine if local livestock grazing management strategies and other land management actions are making progress toward achieving the long-term goals and objectives for streamside riparian vegetation and aquatic resources. The goal is to conduct effectiveness monitoring every three to five years on riparian areas and streambanks. This period of time is considered to be the minimum necessary to detect changes, although unusually wet years and/or flood events may result in short-term changes that validate the need to monitor more frequently, or at least at the time of the event. Budget and personnel constraints may limit the extent in which monitoring of this type will be conducted.

Presently, effectiveness monitoring includes: modified greenline composition (adapted from Winward 2000 and USDI BLM 1996a), woody species height class (Kershner et al. 2004), streambank stability and cover (adapted from Kershner et al. 2004), woody species age class (adapted from Winward 2000), greenline-to-greenline width (Burton et al. 2008), substrate (Bunte and Apt 2001), and residual pool depth and pool frequency (Lisle 1987). These provide data and information concerning the present conditions and trend of riparian vegetation, channels, and streambanks, and to help determine if aquatic systems are being degraded, maintained, or restored across the Malheur National Forest.

4.1.5 ADAPTIVE MANAGEMENT

The adaptive management strategy described below and depicted in Appendix H diagrams 1.0 (Long-term) and 2.0 (Annual) is intended for allotments requiring consultation. It is designed to provide the MNF the ability to make management decisions based on new information, changing conditions, or the results of implementation/effectiveness monitoring. Adaptive management will be used to ensure: 1) sites at desired condition remain in desired condition; 2) sites not in desired condition have an upward trend; and 3) direction from consultation with the Services is met.

The overall strategy consists of a long-term adaptive management strategy and an annual adaptive management strategy. The long-term strategy describes how adaptive management will be used to ensure the three objectives previously stated are achieved and to maintain consistency with Forest Plan level direction. The annual adaptive management strategy describes how adjustments will be made within the grazing season to ensure annual use indicators and other direction from consultation is met, it also describes when and how regulatory agencies will be contacted in the event direction from consultation is not going to be met.

Ideally, the value associated with the annual use indicator is customized to the specific circumstances in each unit. However, customizing this value generally requires a significant amount of data and/or experience with a particular unit. As data is gathered and analyzed the annual use indicators may be adjusted to reflect the new information.

The annual use indicators within the Multiple Indicator Monitoring (MIM) method will be used to detect the annual use of wild horses, wildlife, and livestock at the end of a grazing period or growing season, whichever occurs first. Although the Proposed Action includes a suite of measures designed to avoid such an outcome, the MNF acknowledges that it is nevertheless possible that annual use indicators could be exceeded in a particular year. If this occurs, the MNF proposes the adaptive management process to be initiated immediately and will make any

necessary adjustments to the current or future grazing strategy to ensure that the exceedances do not recur.

When the annual utilization data is collected at the end of the growing season, the MNF will consider adjustments of livestock numbers, timing of grazing, duration of grazing, and endpoint indicators. If the big game populations exceed ODFW Management Objectives, appropriate coordination will occur among the agencies.

If there are recurring exceedances of annual indicators, or if there is a failure to comply with the terms and conditions of the grazing permit, the issuance of a Notice of Non-Compliance may be warranted. This notice, issued to the permittee(s), is likely to be in addition to the outcomes that result from following the adaptive management process described above. The issuance of a Notice of Non-Compliance and resulting action taken by the MNF will be consistent with FSH 2209.13 Section 16 and 36 CFR 222.4. All exceedances of annual indicators and subsequent grazing strategy adjustment recommendations will be documented by the MNF in the annual EOY Report and presented to the Level I consultation team. A specific strategy for when the endpoint indicator for streambank alteration is exceeded is discussed in Section 4.1.5.1 below.

4.1.5.1 COMPLIANCE STRATEGY FOR THE STREAMBANK ALTERATION ENDPOINT INDICATOR

The MNF acknowledges that there is a $\pm 6\%$ margin of error associated with the MIM protocol (see section 4.1.3). Action would only be taken for permit violations and not as a result of wild ungulate or unauthorized use. The MNF will follow the strategy outlined below for exceedance of the bank alteration endpoint indicator. For each level of exceedance, the BMRD will incorporate adaptive management strategies into the following season's grazing strategy which may include: adjustments of livestock numbers, timing of grazing, or duration of grazing.

- **Measured bank alteration $\leq 6\%$ over the endpoint indicator:** will be evaluated by the District IDT. The IDT will examine the level of measured use on stubble height and woody browse to determine if an exceedance of the endpoint indicator occurred. If the IDT concludes that the endpoint indicator has been exceeded the permittee will be contacted via phone or in person to notify them of the IDT findings. The permittee would be given 1 year to remedy. A follow-up letter will be sent to the permittee to document the verbal discussion and include what action is expected of the permittee to remedy the situation, to what standard, and by when (FSH 2209.13, 10, 16.2e).
- **Measured bank alteration 7-13% over the endpoint indicator:** the BMRD will, at a minimum, issue a notice of Non-Compliance for violation of terms and conditions of the term grazing permit and be given 1 year to remedy the non-compliance. Adjustments to the grazing strategy may be made following the adaptive management process. Failure to remedy the non-compliance during the following grazing season may result in a reduction of 25% of permitted AUMs for the following grazing season, **or** rest the pasture the following grazing season (FSH 2209.13, 10, 16.2e).
- **Measured bank alteration 14-20% over the endpoint indicator:** the BMRD will, at a minimum, issue a notice of Non-Compliance for violation of terms and conditions of the term grazing permit and will give the permittee 1 year to remedy the non-compliance. Adjustments to the grazing strategy may be made following the adaptive management process. When documented inspection indicates that the initial non-compliance has not

been remedied as specified, or if a second situation of non-compliance has occurred, the permittee will be contacted by phone or in person describing the specific non-compliance. The BMRD will either reduce the authorized use by 25% of permitted AUMs for the following grazing season, **or** rest the pasture the following grazing season. A follow-up letter of a notice of permit action for non-compliance will be sent to the permittee indicating that a specified part of the permitted numbers or seasons is being suspended for a period of at least two years (FSH 2209.13, 10, 16.2e).

- **Measured bank alteration >21% over the endpoint indicator:** the BMRD will, at a minimum, issue a notice of Non-Compliance for violation of terms and conditions of the term grazing permit and will give the permittee 1 year to remedy the non-compliance. Adjustments to the grazing strategy may be made following the adaptive management process. When documented inspection indicates that the initial non-compliance has not been remedied as specified, or if a second situation of non-compliance has occurred, the permittee will be contacted by phone or in person describing the specific non-compliance. The BMRD will reduce the authorized use by 25% of permitted AUMs for the following grazing season, **and** rest the pasture the following grazing season using the adaptive management process. A follow-up letter of a notice of permit action for non-compliance will be sent to the permittee indicating that a specified part of the permitted numbers or seasons is being suspended for a period of at least two years (FSH 2209.13, 10, 16.2e).

Recurring non-compliance may lead to suspension of AUMs and/or the cancellation in part or whole of the Term Grazing Permit. Permit action involving the suspension or cancelation of grazing permits as per direction outlined in FSH 2209.13, 10, 16.2 and 36 CFR 222.4.

4.1.6 COORDINATION AND REPORTING

Reporting

Annual end-of-year (EOY) grazing reports are prepared by BMRD staff for all livestock grazing allotments. The reports include monitoring results, descriptions of any exceedance of grazing end-points and recommendations for management changes for the next grazing season. See monitoring section for a description of the grazing use and stream channel condition indicators for which information is collected, evaluated and reported. The report is sent to the NMFS and/or FWS by March 1 of each year and to permittees as well.

Coordination

EOY report. Both internal and external coordination takes place regarding information and recommendations for changes in management found within the EOY report. The recommendations for changes in management in the EOY report are developed in an interdisciplinary manner. Typically, range conservationists, fish biologists, hydrologists, and the line officer will be involved. On occasion, wildlife biologists and botanists will participate.

Level 1 Team Meeting: A Level 1 team meeting is scheduled after a draft EOY report is sent to NMFS and/or FWS. The Level 1 Team discusses the draft EOY monitoring results, proposed remedies, and application of the compliance strategy (Section 4.1.5.1).

4.2 INTERRELATED ACTIONS

The regulations require the MNF to impose penalties for violation of prohibited acts on public lands. Unauthorized use is a prohibited act, and therefore is not a federal action. If unauthorized use occurs, the MNF's response could constitute a separate, interrelated federal action.

Unauthorized livestock grazing can occur in the allotment, and is reasonably certain to occur in the future. It is not a large problem in the allotments analyzed in this BA. Typically, it involves a few cow/calf pairs every few years.

Forest Service grazing regulations define unauthorized use, also known as "trespass," as occurring when livestock not under permit enter National Forest System (NFS) lands. It is a violation of 36 CFR 261.7. When unauthorized use occurs, the MNF attempts to identify and contact the owner of the livestock with instructions to remove the unauthorized livestock from NFS. The MNF can then bill the owner for the unauthorized use at the appropriate rate as identified in 36 CFR 222.50(h). If the ownership of the livestock is unknown, or the owner fails to comply with instructions to remove the livestock, the impoundment of said livestock by the MNF can occur as per 36 CFR 262.10.

4.3 PROJECT ELEMENTS

Project elements are the component parts of the action. Project elements will be assessed in the effects analysis section of the BA. Several of the project elements involve the use of vehicles on and off roads to access sites, such as four wheel drive trucks and OHVs.

1. Livestock use of allotment/pastures. Livestock will utilize the allotment/pastures consistent with the permitted numbers, season of use and grazing system described above and in the term grazing permit.
2. Permittee management of livestock and infrastructure maintenance. This includes move-in and move-out of cattle, herding, placement of nutrient (salt blocks) in the uplands, and maintenance of troughs, springs, ponds, fences and gates. Use of highway and off-road vehicles is included in this PE.
3. Range improvements. This includes the construction of fences for riparian pastures, and the construction/development of off-stream water sources.
4. Exclusionary fencing. Fences are constructed or placed to exclude areas from grazing. This is done to prevent livestock damage of riparian areas and in the case of electric fencing, to eliminate the potential for cattle stepping on redds.
5. Monitoring. A variety of implementation and effectiveness monitoring techniques are employed to determine if desired conditions are being met. The MNF Riparian Monitoring Strategy is discussed in detail in Section 4.1.4 below. Workers use manual and electronic equipment to measure vegetation, water quality and stream channel/streambed characteristics.
6. Adaptive management. An adaptive management strategy is designed to provide the MNF the ability to make management decisions based on new information, changing conditions, or the results of implementation/effectiveness monitoring. It will be used to ensure: (1) Sites at desired condition remain in desired condition; (2) sites not in desired condition have an upward trend; and (3) direction from ESA consultation with NMFS is met. The adaptive management strategy describes how adjustments will be made to ensure annual endpoint indicators as well as other direction from this consultation are met, and describes when and how regulatory agencies will be contacted in the event

direction from this ESA consultation is not going to be met. The MNF Adaptive Management Strategy is described in Section 4.1.5.

4.4 ESA ACTION AREA

The ESA action area for this consultation is defined by the entire Fields Peak and northern portions of the Seneca, Deadhorse, and Hanscomb allotment boundaries that flow into the John Day River basin (Figure 6). The ESA action area is displayed by cross-hatched area within the allotment boundary of the 6th field HUCs listed in Table 1.




FIGURE 6. ESA ACTION AREA MAP FOR THE SENECA, DEADHORSE, HANSCOMB, AND FIELDS PEAK ALLOTMENTS CONSULTATION

5 STATUS OF THE SPECIES AND DESIGNATED CRITICAL HABITAT

The status of the Middle Columbia River Steelhead distinct population segment (DPS) and its designated critical habitat (CH) is presented in this section. No bull trout or bull trout CH are found in the action area. Therefore, no description of the status of bull trout and its designated CH is presented.

5.1 MIDDLE COLUMBIA RIVER STEELHEAD DISTINCT POPULATION SEGMENT

5.1.1 LISTING HISTORY AND LOCATION

The Middle Columbia River Steelhead DPS was listed by NMFS as Threatened under the Federal ESA on March 25, 1999 (64 FR 15417). NMFS reaffirmed its threatened status on January 5, 2006 (71 FR 834). Protective regulations for MCR Steelhead were issued under section 4(d) of the ESA on July 10, 2000 (65 FR 42423). The NMFS revised the 4(d) protective regulations on June 28, 2005 (70 FR 37160).

The MCR Steelhead DPS includes all naturally-spawned populations of steelhead in streams within the Columbia River basin from above the Wind River in Washington and the Hood River in Oregon (exclusive), upstream to, and including, the Yakima River in Washington, excluding steelhead from the Snake River basin (64 FR 14517; March 25, 1999). The major tributaries occupied by this DPS are the Deschutes, John Day, Klickitat, Umatilla, Walla Walla, and Yakima River systems. The John Day River (JDR) probably represents the largest naturally spawning and most viable native stock of steelhead in the region (NMFS 2009). The MCR Steelhead DPS does not include co-occurring resident forms of *O. mykiss* (rainbow trout).

5.1.2 LIFE HISTORY AND HABITAT REQUIREMENTS

Steelhead trout are the anadromous form of *O. mykiss*. Adult summer steelhead typically return to freshwater from June through September. Adults overwinter in large rivers while sexually maturing. Adults resume migration to spawning streams in early spring.

The JDR adult summer steelhead enter the lower river as early as September and as late as March, depending on water temperatures. Adult migration in the JDR generally peaks in October. The JDR below the North Fork JDR is used only for migration due to high summer water temperatures. Spawning takes place from March through May. Eggs incubate during the spring and emergence occurs from April through July depending on water temperatures. Juveniles typically rear for 2 to 3 years in freshwater before smolting and migrating to the ocean.

Juvenile steelhead generally utilize habitats with higher water velocities than juvenile Chinook salmon. In winter, juveniles utilize deep pools with abundant cover. Juveniles may reside in their natal stream for their entire freshwater rearing phase or may migrate to other streams within a watershed. Smoltification occurs during late winter and emigration to the ocean occurs during spring. Smolts outmigrate rapidly, taking 45 days or less to reach the ocean from upstream rearing areas. In the JDR below the North Fork, smolts generally stay within the thalweg, taking advantage of cover provided by depth and turbidity. Approximately 80% of the steelhead rear in the ocean for 2 years before returning to the JDR system as adults to spawn (PD BLM 2006).

5.1.3 MCR STEELHEAD POPULATIONS

The Interior Columbia Basin Technical Recovery Team (ICTRT) (2003) identified 15 populations in four major population groups (MPG) (Cascades Eastern Slopes Tributaries, John Day River (JDR), the Walla Walla and Umatilla Rivers, and the Yakima River) and one unaffiliated independent population (Rock Creek) in this steelhead DPS. There are two extinct populations in the Cascades Eastern Slopes Tributaries MPG, the White Salmon River and Deschutes River above Pelton Dam.

The JDR Subbasin contains the MCR Steelhead JDR MPG that consists of the Lower Mainstem John Day (LMJD), North Fork John Day (NFJD), Middle Fork John Day (MFJD), South Fork John Day (SFJD), and Upper Mainstem John Day (UMJD) populations (ICTRT 2003). The action area is associated with the NFJD, MFJD and UMJD populations.

5.1.4 MCR STEELHEAD DPS VIABILITY STATUS

The status of a salmon or steelhead species is expressed in terms of likelihood of persistence over 100 years, or in terms of risk of extinction within 100 years. The ICTRT defined viability at two levels: less than 5 percent risk of extinction within 100 years (viable) and less than 1 percent risk of extinction within 100 years (highly viable). A third category, “maintained,” represents a less than 25 percent risk. The risk level of the steelhead DPS as a whole is built up from the aggregate risk levels of the populations and MPGs. The viable salmonid population (VSP) parameters (abundance, productivity, spatial structure, and diversity of the component populations) must be taken into account to determine the risk level.

The MCR Steelhead DPS does not currently meet viability criteria because its four component MPGs are not at low risk. However, for this DPS the outlook is relatively optimistic. One population, North Fork John Day, is currently at very low risk or “highly viable.” Two populations are currently viable (Deschutes Eastside, Fifteenmile); eleven are at moderate risk, with good prospects for improving. However, three large populations at high risk (Deschutes Westside, Naches, and Upper Yakima) are important to DPS viability; these present significant challenges.

Significant programs are underway for natural recolonization (White Salmon) or reintroduction (Deschutes Crooked River above Pelton Dam) of two of the extirpated populations to historically accessible habitat. Success of these programs should help improve overall DPS viability.

The MCR Steelhead Recovery Plan (NMFS 2009) presented viability ratings for the MCR Steelhead MPG. The risk of extinction is displayed as a combination of ratings for Abundance/Productivity Risk and Spatial Structure/Diversity Risk (Figure 7). The North Fork John Day MPG rates low/very low by the two criteria. The Middle Fork and South Fork MPGs rate low/moderate and the Lower Mainstem and Upper Mainstem MPGs have the highest extinction risk at moderate/moderate.

5.1.5 JOHN DAY RIVER MPG POPULATION STATUS

The current status of the MCR Steelhead John Day River MPG populations, showing 10-year geometric mean abundance by population, estimated productivity, and the minimum abundance threshold needed for long-term viability is summarized in Table 14. The table also includes the 10-year geometric mean proportion of hatchery spawners for the populations where data are available, and the risk ratings of high, moderate, low, and very low, for abundance and productivity combined, and spatial structure and diversity combined.




FIGURE 7. VIABILITY RATINGS FOR THE MCR STEELHEAD MAJOR POPULATION GROUP (NMFS 2009). SHADES OF GREEN INDICATE LOWER RISK OF EXTINCTION AND SHADES OF RED INDICATE HIGHER RISK.

TABLE 14. MCR STEELHEAD JOHN DAY RIVER MPG SUMMARY OF ABUNDANCE, PRODUCTIVITY, RISK RATINGS, AND MINIMUM ABUNDANCE THRESHOLDS (SOURCE: MIDDLE COLUMBIA RIVER STEELHEAD DPS RECOVERY PLAN SUMMARY 2009).

Population	Abundance Threshold ¹	Size Category	Run Timing	10-year Geomean abundance	Abundance Range	10-yr Hatchery Fraction ²	Productivity ³	Productivity Standard Error	A&P Risk Rating ⁴	SSD Risk Rating
Lower Mainstem John Day	2250	Very Large	Summer	1800	563-6257	0.1	2.99	0.24	M	M
North Fork John Day	1500	Large	Summer	1740	369-10,235	0.08	2.41	0.22	VL	L
Upper Mainstem John Day	1000	Intermed.	Summer	524	185-5169	0.08	2.14	0.33	M	M
Middle Fork John Day	1000	Intermed.	Summer	756	195-3538	0.08	2.45	0.16	M	M
South Fork John Day	500	Basic	Summer	259	76-2729	0.08	2.06	0.27	M	M

¹ Abundance threshold for viability based on habitat intrinsic potential

² Average proportion of hatchery spawners over most recent 10 years in the data series.

³ Geomean return per spawner calculated over most recent 20 years in data series.

⁴ Abundance & Productivity Risk Ratings: H = high risk, M= moderate risk, L = low risk, VL = very low risk

5.1.6 POPULATION LIMITING FACTORS

The Middle Columbia River Steelhead ESA Recovery Plan (NMFS 2009) identified population limiting factors. For the NFJD population the primary tributary habitat limiting factors identified by the recovery planning team are degraded floodplain connectivity and function, degraded channel structure and complexity (habitat quantity, habitat diversity, channel stability), altered sediment routing, water quality (temperature), and altered hydrology. For the MFJD population they are degraded floodplain and channel structure (habitat quantity/diversity), altered sediment routing, altered hydrology and water temperature. The primary tributary limiting factors for the SFJD population include altered sediment routing, degraded floodplain and channel structure (habitat quantity and habitat diversity), altered hydrology, water quality (temperature) and blocked or impaired fish passage.

Tributary limiting factors for the UMJD population include degraded channel structure and complexity (habitat quantity and diversity), degraded riparian areas and LWD recruitment, altered sediment routing, water temperatures and altered hydrology. Impaired fish passage is also a priority limiting factor in Beech and Laycock creeks. Habitat limiting factors identified in ODFW (2010) for the Beech Creek Major Spawning Area within the ESA action area are displayed in Table 15.

TABLE 15. HABITAT LIMITING FACTORS WITHIN THE BEECH CREEK MAJOR SPAWNING AREA IDENTIFIED IN NMFS (2009).

Limiting Factor	Beech Creek Major Spawning Area
Degraded floodplain connectivity and function	X
Degraded channel structure and complexity	Lower reaches next to Highway 395
Altered hydrology	Lower reach of McClellan Creek
Altered sediment routing	X
Water Temperature	X
Degraded riparian communities	X
Man-made block to migration	
Impaired fish passage	X

5.2 CRITICAL HABITAT FOR MIDDLE COLUMBIA RIVER STEELHEAD DPS

5.2.1 DESIGNATION HISTORY

Critical habitat (CH) was designated for MCR Steelhead on February 16, 2000 (65 FR 7764) that encompassed the major Columbia River tributaries known to support the DPS, including the

Deschutes, John Day, Klickitat, Umatilla, Walla Walla, and Yakima Rivers, as well as the Columbia River and estuary.

In late 2000, a lawsuit was filed challenging the NMFS February 2000 final designation of CH for ESUs/DPSs of Pacific salmon and steelhead listed under the ESA. A federal court ruled that the agency did not adequately consider the economic impacts of the CH designations. In April 2002, NMFS withdrew its 2000 CH designations.

Critical habitat for MCR Steelhead was designated again on September 2, 2005 (70 FR 52630). Designated CH includes the stream channels within the designated stream reaches, and includes a lateral extent as defined by the ordinary high-water line (33 CFR 319.11). In areas where ordinary high-water line has not been defined, the lateral extent is defined by the bankfull elevation. Bankfull elevation is the level at which water begins to leave the channel and move into the floodplain and is reached at a discharge which generally has a flood recurrence interval of 1 to 2 years on the annual flood series.

5.2.2 PRIMARY CONSTITUENT ELEMENTS

The physical or biological features of CH essential to the conservation of the species are known as primary constituent elements (PCEs). The PCEs of MCR Steelhead CH are those sites and habitat components that support one or more life stages, including:

- (1) Freshwater spawning sites with water quantity and quality conditions and substrate supporting spawning, incubation and larval development;
- (2) Freshwater rearing sites with:
 - (i) Water quantity and floodplain connectivity to form and maintain physical habitat conditions and support juvenile growth and mobility;
 - (ii) Water quality and forage supporting juvenile development; and
 - (iii) Natural cover such as shade, submerged and overhanging large wood, log jams and beaver dams, aquatic vegetation, large rocks and boulders, side channels, and undercut banks.
- (3) Freshwater migration corridors free of obstruction and excessive predation with water quantity and quality conditions and natural cover such as submerged and overhanging large wood, aquatic vegetation, large rocks and boulders, side channels, and undercut banks supporting juvenile and adult mobility and survival.

5.2.3 STATUS OF MIDDLE COLUMBIA RIVER STEELHEAD CRITICAL HABITAT

Migratory habitat quality for MCR steelhead has been severely degraded by the development of the Federal Columbia River Power System. Depending on their natal watershed, adults and out-migrating juvenile steelhead encounter between one and three mainstem Columbia River dams migrating to and from the ocean. Hydroelectric development has modified natural flow regimes resulting in higher water temperatures, changes in fish community structure, and increased travel time for migrating adults and juvenile salmonids. Physical features of dams such as turbines also kill migrating fish. The only substantial habitat blockages at present for this species are Pelton Dam on the Deschutes River and Condit Dam on the White Salmon River. However, minor blockages from smaller dams, impassable culverts, and irrigation dams occur

throughout the region. Several dams in the John Day River basin previously blocked habitat, but they have since been modified with ladders; however, there is a possibility that local native stocks were extirpated before these ladders were built (NMFS 2004).

Water quality impairment that affects spawning, migration, and rearing is a problem in many areas of designated CH for the MCR Steelhead. Summer stream temperature is the primary water quality problem for this species, and many of the stream reaches proposed as CH are listed on the Clean Water Act (CWA) 303(d) list for water temperature. Many areas that were historically suitable rearing and spawning habitat are now unsuitable due to high summer stream temperatures. Elevated stream temperatures may form thermal barriers to juvenile migration within tributaries. Removal of riparian vegetation, alteration of natural stream morphology, and withdrawal of water for agricultural or municipal use all contribute to elevated stream temperatures. Contaminants such as insecticides and herbicides from agricultural run-off and heavy metals from mine waste are common in some areas of designated critical habitat for this species.

Low summer stream flow is also a common characteristic affecting spawning, rearing, and migration PCEs for this DPS. There is little or no late summer flow in sections of the lower Umatilla and Walla Walla rivers. Withdrawal and storage of natural stream flow in spawning and rearing areas have altered hydrological cycles, causing a variety of adverse impacts to MCR Steelhead habitat. Increased summer stream temperatures, migration blockages, stranding of fish, and alteration of sediment transport processes can result from water withdrawal for irrigation or municipal use (NMFS 1996; Spence *et al.* 1996). In many river basins, the amount and quality of available rearing habitat has been reduced by water withdrawals. Many stream reaches are over-appropriated under state water law, with more allocated water rights than existing stream flow conditions can support.

Spawning and rearing salmonids, such as steelhead, require physically complex lotic habitats with pools, large woody debris, undercut banks, and substrates with low levels of fine sediments (Spence *et al.* 1996; Bjornn and Reiser 1991). Although these habitat conditions are still present in many wilderness, roadless, and undeveloped areas, recent subbasin assessments and plans (NPCC 2005) indicate that habitat complexity has been greatly reduced in many areas of designated critical habitat. Channel and riparian alterations for agricultural purposes, transportation, mining, forestry and other development activities have affected spawning, rearing and migration PCEs by reducing overall habitat complexity, cover, food availability, and spawning and rearing quality and quantity.

Under section 303(d) of the Clean Water Act, the Oregon Department of Environmental Quality (ODEQ) identified many streams within the LJD, UJD, MFJD, and NFJD watersheds that are water quality limited for high temperatures, dissolved oxygen, or biological criteria. Additionally, the ODEQ identified total phosphates and fecal coliform as water quality limitations for many streams within the Lower Mainstem John Day River, and sediment for many NFJD streams (NMFS 2004).

Critical Habitat Analytical Review Teams (CHARTs) were convened by NMFS for each recovery domain (NMFS 2005). CHARTs were charged with analyzing the best available data for each listed species, to make findings regarding the presence of essential habitat features in each watershed, identify potential management actions that may affect those features, and determine the conservation value of each watershed within each species' range. The action area

occurs within four 5th-field HUCs in the range of MCR steelhead: Canyon Creek, Laycock Creek-John Day River, Fields Creek-John Day River, and Murderers Creek. These HUCs have medium to high conservation value. Mid-Columbia CHART members noted that PCEs in these HUCs support unique genetic resources since there is minimal hatchery influence on these populations.

The John Day Subbasin Plan (NPCC 2005) included an Ecosystem and Diagnostic Treatment (EDT) analysis of habitat conditions for the 5th field HUCs located in the action area. The approach was to display the top quartile protection and/or restoration 5th field HUCs and their important restoration attributes.

Fifteen 5th field HUCs identified as important to Upper John Day summer steelhead were evaluated. Laycock and Fields creeks were in the top quartile for restoration benefit. The attributes for restoration in common for these watersheds were habitat diversity, sediment load and key habitat quantity. In the South Fork John Day, Murderers Creek was a high priority HUC5 for both protection and restoration, signifying that it should be protected from further degradation and that restoration on any of the limiting factors listed (temperature, sediment load and key habitat quantity) would have the potential to increase productivity and abundance for the population.

The Subbasin Plan also evaluated 5th field HUCs by restoration priority by population. The Upper Mainstem and South Fork John Day River populations were evaluated together. Murderers, Fields, and Laycock Creek HUC6 watersheds were listed as high priority for restoration to benefit MCR steelhead.

6 ENVIRONMENTAL BASELINE

As mentioned in earlier sections, the predominant land use activity in the action area is livestock grazing for which there have been MNF formal and informal ESA consultations. The past, present and anticipated impacts of future Federal livestock grazing which have undergone formal consultation have been taken into account in the following description of the environmental baseline.

6.1 NMFS MATRIX OF PATHWAYS AND INDICATORS

A NMFS process paper titled “Making Endangered Species Act Determinations of Effect for Individual or Grouped Actions at the Watershed Scale” (NMFS 1996) is used to describe the environmental baseline. It is commonly known as the NMFS Matrix of Pathways and Indicators, hereafter referenced as the “NMFS MPI.” The NMFS MPI identifies indicators to analyze for the following pathways: 1) Water quality; 2) Habitat access; 3) Habitat elements; 4) Channel condition and dynamics; 5) Flow/hydrology; and, 6) Watershed condition. The condition of each indicator is described as either “Properly Functioning” (PF), “At Risk (AR),” or “Not Properly Functioning (NPF)” based upon specific numeric or qualitative criteria. Table 16 presents the current status of the environmental baseline using the NMFS MPI for the Upper John Day River. Although the action area is located in two sub-basins (Upper John Day and Silvies), only Upper John Day contains MCR steelhead and its critical habitat. Therefore, Silvies sub-basin will not be discussed or analyzed further. Table cells in bold print indicate the current status of each indicator. The habitat indicators in the NMFS matrix also correspond to the PCEs of designated CH. The relationship between NMFS MPI habitat indicators and PCEs of CH is discussed in

Section 7.2 (Analysis of Effects to Designated Critical Habitat).

TABLE 16. STATUS OF ENVIRONMENTAL BASELINE FOR THE UPPER JOHN DAY SUB-BASIN.¹

Pathway	Indicators	Properly Functioning	At Risk	Not Properly Functioning
Water Quality	Temperature	50 – 57° F (max 7-day average)	57 – 61° F (spawning, max 7-day average)	> 61° F (spawning, max 7-day average)
			57 – 64° F (migration and rearing, max 7-day average)	> 64° F (migration and rearing, max 7-day average)
	Sediment	< 12% fines (<0.85mm) in gravel	12 – 20% fines	> 20% fines
	Chemical Contaminants or Nutrients	Low levels of chemical contamination from agricultural, industrial, and other sources; no excess nutrients; no CWA 303d designated reaches	Moderate levels of chemical contamination from agricultural, industrial, and other sources; some excess nutrients; one CWA 303d designated reach	High levels of chemical contamination from agricultural, industrial, and other sources; high levels of excess nutrients; more than one CWA 303d designated reach
Habitat Access	Physical Barriers	Any man-made barriers present in watershed allow upstream and downstream fish passage at all flows	Any man-made barriers present in watershed do not allow upstream and/or downstream fish passage at base/low flows	Any man-made barriers present in watershed do not allow upstream and/or downstream fish passage at a range of flows
Habitat Elements	Substrate	Dominant substrate is gravel or cobble (interstitial spaces clear), or embeddedness <20%	Gravel and cobble is subdominant, or if dominant, embeddedness 20 – 30%	Bedrock, sand, silt, or small gravel dominant, or if gravel and cobble dominant, embeddedness >30%
	Large Woody Debris	> 20 pieces/mile (> 12 inch diameter and > 35 ft. length), and adequate sources of woody debris recruitment in riparian areas	Currently meets standards for Properly Functioning, but lacks potential sources from riparian areas of woody debris recruitment to maintain that standard	Does not meet standards for Properly Functioning and lacks potential large woody debris recruitment

Pathway	Indicators	Properly Functioning	At Risk	Not Properly Functioning
	Pool Frequency	Meets pool frequency standards and meets large woody debris recruitment standards for Properly Functioning habitat	Meets pool frequency standards but large woody debris recruitment inadequate to maintain pools over time	Does not meet pool frequency standards
	Pool Quality	Pools > 1 meter deep (holding pools) with good cover and cool water; minor reduction of pool volume by fine sediment	Few deeper pools (> 1 meter) present or inadequate cover/temperature; moderate reduction of pool volume by fine sediment	No deep pools (> 1 meter) and inadequate cover/temperature; major reduction of pool volume by fine sediment
	Off Channel Habitat	Backwaters with cover, and low energy off-channel areas (ponds, oxbows, etc.)	Some backwaters and high energy side channels	Few or no backwaters; no off-channel ponds
	Refugia	Habitat refugia exist and are adequately buffered (e.g., by intact riparian reserves); existing refugia are sufficient in size, number, and connectivity to maintain viable populations or subpopulations (all life stages and forms)	Habitat refugia exist but are not adequately buffered (e.g., by intact riparian reserves); existing refugia are insufficient in size, number, and connectivity to maintain viable populations or subpopulations (all life stages and forms)	Adequate habitat refugia do not exist
Channel Condition & Dynamics	Width/Depth Ratio	< 10	10 – 12	> 12
	Stream Bank Condition	> 80% of any stream reach has > 90% stability	50 – 80% of any stream reach has > 90% stability	< 50% of any stream reach has > 90% stability
	Floodplain Connectivity	Off-channel areas are frequently hydrologically linked to main channel; overbank flows occur and maintain wetland functions, riparian vegetation, and succession	Reduced linkage of wetland, floodplains, and river areas to main channel; overbank flows are reduced relative to historic frequency, as evidenced by moderate degradation of wetland function and riparian vegetation/succession	Severe reduction in hydrologic connectivity between off-channel, wetland, floodplain, and riparian areas; wetland extent drastically reduced, and riparian vegetation/success altered significantly

Pathway	Indicators	Properly Functioning	At Risk	Not Properly Functioning
Flow/Hydrology	Change in Peak/Base Flows	Watershed hydrograph indicates peak flow, base flow, and flow timing characteristics comparable to an undisturbed watershed of similar size, geology, and geography	Some evidence of altered peak flow, base flow, and/or flow timing relative to an undisturbed watershed of similar size, geology, and geography	Pronounced changes in peak flow, base flow, and/or timing relative to an undisturbed watershed of similar size, geology, and geography
	Increase in Drainage Network	Zero or minimum increases in drainage network density due to roads	Moderate increases in drainage network density due to roads (e.g., 5%)	Significant increases in drainage network density due to roads (e.g., 20 – 25%)
Watershed Condition	Road Density & Location	< 2 mi/mi ² ; no valley bottom roads	2 – 3 mi/mi ² ; some valley bottom roads	> 3 mi/mi²; many valley bottom roads
	Disturbance History	< 15% ECA (entire watershed) with no concentration of disturbance in unstable or potentially unstable areas, and/or refugia, and/or riparian areas	< 15% ECA (entire watershed) but disturbance concentrated in unstable or potentially unstable areas, and/or refugia, and/or riparian areas	> 15% ECA (entire watershed) and disturbance concentrated in unstable or potentially unstable areas, and/or refugia, and/or riparian areas
	Riparian Management Areas	The riparian reserve system provides adequate shade, large woody debris recruitment, and habitat protection and connectivity in all subwatersheds, and buffers or includes known refugia for sensitive aquatic species (>80% intact), and/or for grazing impacts; percent similarity of riparian vegetation to the potential natural community/ composition > 50%	Moderate loss of connectivity or function (shade, LWD recruitment, etc.) of riparian reserve system, or incomplete protection of habitats and refugia for sensitive aquatic species (~ 70 – 80% intact), and/or for grazing impacts; percent similarity of riparian vegetation to the potential natural community/ composition 25 – 50% or better	Riparian reserve system is fragmented, poorly connected, or provides inadequate protection of habitats and refugia for sensitive aquatic species (< 70% intact), and/or for grazing impacts; percent similarity of riparian vegetation to the potential natural community/ composition < 25%

¹ Bold text in table cells indicates current status of the indicator

The environmental baseline using the NMFS MPI ratings in Table 14 is based on scientific literature review, management documents and the professional judgment of MNF Forest and District fishery biologists, hydrologists, soil scientists and range conservationists. The MCR Steelhead Recovery Plan, the Malheur National Forest Roads Analysis Report, and the Forest

water temperature monitoring program support the environmental baseline ratings. The rationale from the supporting documents and programs for these ratings are summarized in the following three sections: 6.1.1; 6.1.2; and 6.1.3.

Historic and current cattle grazing in the ESA action area likely play varying roles in the current environmental baseline ratings for these affected subbasins. In some situations the actual small streams and corresponding 6th field subwatersheds draining the ESA action area may be properly functioning or functioning at risk while the larger subbasins are not properly functioning. Grazing is one of multiple natural and human-caused watershed disturbances influencing environmental baseline ratings. In some circumstances in these subbasins the proposed action has causal mechanisms that affect fish habitat indicators analyzed in the environmental baseline ratings.

6.1.1 MIDDLE COLUMBIA RIVER STEELHEAD RECOVERY PLAN

The Middle Columbia River Steelhead Recovery Plan (NMFS 2009) describes habitat conditions for the UMJD and SFJD River populations. The recovery plan states that there has been a loss of off-channel and side-channel habitats. Consequently there is reduced spawning and rearing habitat and a loss of refugia from high flows. Floodplain connectivity has been degraded. Removal of beavers has reduced habitat complexity, floodplain function and the amount of stored water. Livestock grazing has increased channelization and negatively impacted large wood debris (LWD), cover and bank stability.

More than 50 percent of measured streams had greater than 35 percent substrate embeddedness (NMFS 2009). Embeddedness is also a problem in 5 of 8 Canyon Creek reaches and 10 of 32 mainstem John Day River reaches (MNF 2004). Livestock grazing, logging, road construction, beaver removal and fire suppression have contributed to altered flow regimes. Water withdrawals dewater the lower reach of McClellan Creek. Elevated stream temperatures are common to almost all streams in the Upper John Day. Historical mining has added to water temperature problems by removing riparian vegetation, simplifying stream channels and changing substrate composition (NMFS 2009). High water temperatures in tributaries and portions of the mainstem alter or block juvenile steelhead movements in the summer months.

Riparian conditions are degraded. Roads have altered riparian functions. The MCR steelhead Recovery Plan states that the MNF identified about 124 miles of roads within Riparian Habitat Conservation Areas (MNF 2004). Forest practices and grazing on private and public land have altered riparian vegetation and LWD potential has also been reduced.

There are additional sources of information to describe the environmental baseline at finer scales than the UMJD River population. They include a MNF roads analysis report, water temperature monitoring information, PIBO EM results, Forest Service level II stream inventories, annual end-of-year grazing reports for the years 2008-2010 (MNF 2009a, MNF 2009b, MNF 2011), Proper Functioning Condition (PFC) assessments (Prichard et al. 1994) and Multiple Indicator Monitoring (MIM) (Burton et al. 2011) monitoring results at Designated Monitoring Areas (DMA) for specific pastures in the allotments for years ranging from 2008 to 2010. The information provided by each of these sources is presented and interpreted below.

6.1.2 MALHEUR NATIONAL FOREST ROADS ANALYSIS REPORT

The MNF prepared an analysis of its road system in a document titled “Malheur National Forest Roads Analysis Report” (MNF 2004). Among the issues analyzed was the risk of the existing road network to general watershed health at the scale of 6th level hydrologic unit codes (HUC), commonly known as sub-watersheds. Many scientific studies have documented the impacts of roads on to fish, fish habitat, and watershed function. Effects include habitat fragmentation from stream crossing structures that block migration, increases in peak flows from high road density, increased sedimentation and isolating streams from their floodplains (USDA FS 2001). The MNF used a Geographic Information System assessment to determine watershed risk. The following watershed risk rating elements were used:

- Total road density (roads in management levels 1-5)
- Road density (roads in management levels 1 and 2)
- Total road density within 200 feet of perennial and intermittent streams
- Density within 200 feet of perennial and intermittent streams (roads in management levels 1 and 2)
- Total road-stream crossing density (crossings/square mile)
- Geologic Sensitivity
- Soil Sensitivity

Ranges of values for each element were assigned a risk rating of low, moderate, high or extreme (Table 17). For example, for total road density, an “extreme” risk rating was for densities greater than five miles per square mile and a “low” rating was for densities less than one mile per square mile. To determine the overall subwatershed risk rating, the risk rating for each element was assigned a numeric value. They ranged from 1 for a rating of “low” to 4 for a rating of “extreme.” The individual element numeric scores were then added for a total score. Total scores exceeding 23 were given an overall watershed risk rating of “extreme,” scores in the 17-23 range were given a “high” rating, scores from 11-17 were given a “moderate” rating and scores less than 11 were given a rating of “low.”

TABLE 17. RANGES OF VALUES BY RISK CATEGORY FOR ELEMENTS USED IN THE WATERSHED RISK ANALYSIS.

Risk Element	Risk Category			
	Low	Moderate	High	Extreme
Total Road Density (miles/mile ²)	0 - 1	1 - 3	3 - 5	>5
Level 1-2 Road				

Risk Element	Risk Category			
	Low	Moderate	High	Extreme
Density (miles/mile ²)	0 - 1	1 – 2.5	2.5 – 4	>4
Road Density w/in 200 feet of streams (miles/mile ²)	0 – 0.2	0.2 – 0.6	0.6 – 0.9	>0.9
Level 1-2 Road Density w/in 200 feet of streams (miles/mile ²)	0 – 0.2	0.2 – 0.5	0.5 – 0.8	>0.8
Road Stream Crossing Density (#crossings/mile)	0 – 1.5	1.5 - 3	3 – 4.5	>4.5
Percent of Subwatershed with Sensitive Geology	1 - 20	20 – 50	50 – 100	Not applicable
Percent of Subwatershed with Sensitive Soils	1 - 20	20 – 50	50 – 100	Not applicable

The results for the eight 6th field HUCs represented in the action area for this consultation are shown in Table 18. Fields Creek was given a “high” watershed risk rating, the Middle Canyon Creek, Clark Creek-John Day River, and Laycock Creek 6th field subwatersheds had a “moderate” watershed risk rating, and the Riley Creek, Dry Creek-John Day, and Headwaters Murderers Creek 6th fields had a “low” watershed risk rating. The individual risk ratings for the road density, road proximity within 200 feet, and road crossings elements were “high” or “extreme” for the Headwaters Murderers Creek, Middle Canyon Creek, Laycock Creek, and Fields Creek 6th fields. This suggests that the legacy road system has negatively impacted riparian and aquatic environmental baseline conditions in these five subwatersheds.

The NMFS MPI values for the Road Density and Location (RDL) indicator are <2, 2-3 miles/mile² and >3 miles/mile² for the PF, AR and NPF categories, respectively. The “high” and

“extreme” risk ratings for both road density risk elements are equivalent to the NMFS MPI NPF category. Therefore, the Headwaters Murderers Creek, Middle Canyon Creek, Laycock Creek, and Fields Creek 6th fields are NPF for RDL. The Riley Creek 6th field HUC road density risk rating of “moderate” would be considered either PF or AR, since the road density elements’ ranges for “moderate” (1-3, 1-2.5) encompass the NMFS MPI numeric ranges for the PF and AR categories.

While there are no other roads risk analysis elements that are directly comparable to NMFS MPI indicators, it is logical that the Road Stream Crossing Density (RSCD) watershed risk element would inform an analysis of the NMFS MPI Increase in Drainage Network (IDN) indicator. The “high” or “extreme” risk score for the RSCD risk element for the Headwaters Murderers Creek, Middle Canyon Creek, and Fields Creek 6th field HUCs supports a NMFS MPI classification of NPF for the IDN indicator.

Road crossings at streams are the primary mechanism for rainfall runoff intercepted by roads to enter stream channels. Roads tend to concentrate runoff, resulting in higher peak flows than would occur without roads. Fine sediments from road surfaces also enter stream channels at road crossings, increasing turbidity, substrate embeddedness and substrate composition. The “extreme” or “high” risk ratings for the RSCD risk element for the Headwaters Murderers Creek, Middle Canyon Creek, and Fields Creek 6th field HUCs would logically support classification of NPF for the Change in Peak/Base Flows, Sediment and Substrate NMFS MPI indicators.

The vast majority of road crossings at streams are culverts. Poorly designed culverts can be barriers to juvenile or adult fish passage. The RSCD risk scores do not incorporate fish passage barrier information, but high or extreme risk ratings imply a large number of culverts with potential fish passage problems. The MNF has identified road crossing structures that may have fish passage concerns from Level II stream inventory reports. Seventeen culverts with potential fish passage problems were identified in streams within the ESA action area. The culverts that may have fish passage concerns are located in the following streams by allotment and pasture: Murderers Creek (8), Basin Creek (2), and White Creek (1) in the Fields Peak Allotment, Murderers Creek Pasture; Fields Creek (5), in the Fields Peak Allotment, Fields Creek Pasture; Tex Creek (1), in the Fields Peak Allotment, Tex Creek Pasture.

The two risk elements for road density within 200 feet of streams do not have a comparable NMFS MPI indicator. However, roads within floodplains have the potential to negatively affect the Off-channel Habitat and Floodplain Connectivity NMFS MPI indicators. Many FS roads are in the valley bottoms, in or adjacent to riparian areas, and affect the ability of a stream to meander laterally through its floodplain. There is no information to determine to what degree a distance of 200 feet includes the floodplains for the various streams associated with the road system in the eight 6th field HUCs represented in the Seneca, Deadhorse, Hanscomb, and Fields Peak allotments. However, the “extreme” or “high” risk ratings for the two risk elements for the Headwaters Murderers Creek, Middle Canyon Creek, Fields Creek, Laycock Creek, and Riley Creek 6th field HUCs would tend to support a NMFS MPI classification of NPF for the two NMFS MPI indicators.

TABLE 18. SUB-WATERSHED RISK RATINGS FOR SIXTH FIELD HYDROLOGIC UNITS IN THE ESA ACTION AREA FOR THE SENECA, DEADHORSE, HANSCOMB, AND FIELDS PEAK ALLOTMENTS CONSULTATION (FROM MNF 2004B).

Watershed Risk Element	Hydrologic Unit Code Name and Number						
	Middle Canyon Creek ² 170702010703	Laycock Creek 170702010901	Clark Creek-John Day River 170702010904	Riley Creek 170702010903	Dry Creek-John Day River 170702010906	Headwaters Murderers Creek ³ 170702010301	Fields Creek 170702011002
Road Density Risk (ML ¹ 1-5)	High	High	Low	Moderate	Low	High	High
Road Density Risk (ML 1-2)	High	High	Low	Moderate	Low	High	High
Road 200' Proximity Risk (ML 1-5)	Extreme	Moderate	Low	Moderate	Moderate	Extreme	Extreme
Road 200' Proximity Risk (ML 1-2)	Extreme	High	Low	High	Moderate	Extreme	Extreme
Road Crossings Risk	High	Low	Low	Moderate	Low	Extreme	Extreme
Geologic Sensitivity	Low	Low	Low	Low	Low	Low	Moderate
Soil Erosion Sensitivity	High	Moderate	High	High	High	Moderate	High

Watershed Risk Element	Hydrologic Unit Code Name and Number						
	Middle Canyon Creek ² 170702010703	Laycock Creek 170702010901	Clark Creek- John Day River 170702010904	Riley Creek 170702010903	Dry Creek- John Day River 170702010906	Headwaters Murderers Creek ³ 170702010301	Fields Creek 170702011002
Overall Watershed Risk	High	Moderate	Low	Moderate	Moderate	High	Extreme

Road maintenance level (ML) designations are defined as:

Level 1. These are intermittent service roads during the time they are closed to motorized traffic. The closure period must exceed one year. Basic custodial maintenance is performed to keep damage to adjacent resources to an acceptable level. Emphasis is normally given to maintaining drainage facilities and runoff patterns. Planned road deterioration may occur at this level. Roads receiving level 1 maintenance may be of any type, class, or construction standard.

Level 2. Roads open for use by high clearance vehicles. Passenger car traffic is not a consideration. Traffic is normally minor.

Level 3. Roads open and maintained for travel by a prudent driver in a standard passenger car. User comfort and convenience are not considered priorities. Roads in this maintenance level are typically low speed, single lane with turnouts and spot surfacing. Some roads may be fully surfaced with either native or processed material.

Level 4. Roads that provide a moderate degree of user comfort and convenience at moderate travel speeds. Most roads are double lane and aggregate surfaced. However, some roads may be single lane. Some roads may be paved and/or dust abated.

Level 5. Roads that provide a high degree of user comfort and convenience. These roads are normally double lane, paved facilities. Some may be aggregate surfaced and dust abated.

6.1.3 MALHEUR NATIONAL FOREST WATER TEMPERATURE MONITORING

Appendix I presents water temperature monitoring information for Fields, Wickiup, Buck Cabin, and Tex creeks in the Fields Peak Allotment for 1995-2005 in Table I-1. The former three sites were in the Fields Peak pasture while the Tex Creek site was in the Tex Creek pasture. One site (Fields Creek) had 6 years of data, but most had 1-2. All four monitoring sites were located in MCR Steelhead CH. Fields and Tex creeks are on the Oregon Department of Environmental Quality 303(d) list for water temperature (ODEQ 2010).

Appendix I also presents water temperature monitoring information for Riley Creek in the Riley Creek Meadow pasture of the Deadhorse Allotment for 2005 in Table I-2. This site was located about 4 miles upstream of MCR Steelhead CH in Riley Creek at Riley Creek Meadow. This stream is not on the Oregon Department of Environmental Quality 303(d) list for water temperature.

For the Fields Peak Allotment, the mean yearly maximum of seven day rolling means of the daily maximum in degrees Fahrenheit (7 day mean max) was Fields Creek 65.5 °F, Buck Cabin Creek 61.9 °F, Wickiup Creek 62.5 °F, Tex Creek 71.3 °F. The mean number of days per year over 64 degrees F was Fields Creek 14.5, Buck Cabin Creek 0, Wickiup Creek 1, Tex Creek 50.

For the Deadhorse Allotment the 2005 maximum of seven day rolling means of the daily maximum in degrees Fahrenheit (7 day mean max) was Riley Creek 75.5 °F. The number of days per year over 64 degrees F at the site was 70. In the tables, data are evaluated using the following criteria: 1) State water quality standards; 2) Amendment 29 DFC; 3) PACFISH RMO; and, 4) NMFS MPI.

The state water quality standard of the seven-day mean maximum temperature of 64 °F for streams with anadromous fish passage and salmonid rearing use *was met* for Buck Cabin and Wickiup creeks and *not met* for Fields, Tex, and Riley creeks. The Amendment 29 DFC for Chinook salmon and/or westslope cutthroat trout spawning and rearing habitat of seven day mean maximum of 55 °F was *not met* for any site. Cutthroat trout occur in streams draining north out of the Aldrich Mountains such as Fields, Laycock, and Ingle creeks (MNF corporate GIS files).

The PACFISH RMO has three criteria. Criterion 1 is no measurable increase in summer maximum temperature. Fields Creek was the only site with adequate sample size to determine if a trend existed. Inspection of the six year dataset showed no linear trend in annual temperature maxima, and only a weak correlation (14%) to annual air temperature recorded at John Day (MNF unpub. data). Criterion 2, seven-day mean maximum below 64 °F for migration and rearing habitat, *was met* for Buck Cabin and Wickiup creeks and *not met* for Fields, Tex, and Riley creeks. Criterion 3, seven-day mean maximum below 60 °F for spawning habitat, *was not met* for any stream. The data supported a NMFS MPI rating of NPF for Fields, Tex, and Riley creeks (seven day mean maximum >61 °F for spawning habitat; >64 °F for migration and rearing habitat and a rating of FAR for Buck Cabin and Wickiup creeks.

6.2 PIBO MONITORING

The PACFISH-INFISH Biological Opinion (PIBO) monitoring strategy is described in section 3.1.2.3. Monitoring consists of two components: effectiveness and implementation.

6.2.1 EFFECTIVENESS MONITORING

The following provides a summary of data collected by the PIBO Effectiveness Monitoring Program (EMP) for monitoring locations within the Deadhorse and Fields Peak Allotments; no

monitoring sites are found within the Seneca and Hanscomb Allotments. Sites included within the summary are both integrator⁷ reaches within randomly selected watersheds as well as Designated Monitoring Area (DMA)⁸ reaches that lie within pastures contained by the randomly selected watersheds. There have been 5 reach evaluations (sites evaluated two or more times) within the Deadhorse and Fields Peak Allotments, which occurred between 2003 and 2008. Within this monitored reach the data suggest that some stream attributes are improving, while others are remaining relatively static (see Tables 20 & 21 below).

⁷ PIBO monitoring sites chosen within randomly selected subwatersheds to show integrated effects of upstream management—most are located in the most downstream response reach (stream gradient less than 3%), while the remaining at the downstream most transport reach (stream gradient between 3 and 5%)

⁸ PIBO monitoring sites located within each randomly selected subwatershed where livestock grazing occurs within the riparian area; locations are selected by Field Units and used for annual implementation monitoring; objective is to develop link between implementation and effectiveness monitoring as part of adaptive management feedback process

TABLE 19. PIBO MONITORING RESULTS – DEADHORSE ALLOTMENT

Stream	Site Type	Year	Bf W/D	Bankfull Width (m)	Mean Part. Size (D50) (mm)	Pool % /Res. Pool depth (m)	Pool frequency (#pools/km)	%Fine s <2mm /<6mm	Bank Stab. (%)	Bank Angle (°)	Under-cut %	GL Wet Rat	GL Woody CV
Riley Cr.	I	2003	16.4	3.0	14	60.5/ 0.23	100.5	23.8/ 37.3	98	119	23.8	56	63
Riley Cr.	I	2008	15.1	4.0	5	29.6/ 0.29	59.1	9.3/ 18.5	100	113	33.3	44	38
Riley Cr.	K	2003	-	4.0	-	-	-	-	98	116	19.1	62	49
Riley Cr.	K	2008	10.0	4.0	-	33.5/ 0.29	75.6	-	100	126	14.7	-	-

TABLE 20. PIBO MONITORING RESULTS – FIELDS PEAK ALLOTMENT

Stream	Site Type	Year	Bf W/D	Bankfull Width (m)	Mean Part. Size (D50) (mm)	Pool % /Res. Pool depth (m)	Pool frequency (#pools/km)	%Fine s <2mm /<6mm	Bank Stab. (%)	Bank Angle (°)	Under-cut %	GL Wet Rat	GL Woody CV
Fields Cr.	I	2003	29.5	5.3	52	33.2/ 0.33	47.8	23.6/ 31.5	95	111	29.0	42	98
Fields Cr.	I	2008	23.1	5.9	32	32.1/ 0.29	60.8	6.2/ 13.2	95	118	26.3	30	74
Fields Cr.	K	2003	-	5.9	-	-	-	-	98	125	5.3	50	96
Fields Cr.	K	2008	21.3	5.7	-	20.2/ 0.30	56.6	-	97	130	8.3	62	80
Murderers Cr.	K	2003	-	4.0	-	-	-	-	95	112	32.5	81	9
Murderers Cr.	K	2008	18.1		-	49.4/ 0.49	55.6	-	85	103	34.2	71	8

In reviewing the data, variability exists between monitoring site parameters measured between 2003 and 2008. Overall, bankfull width-to-depth ratios, greenline wetland rating, and greenline woody cover and percent fines <6mm have decreased, undercut banks at Riley Creek have increased, while the remaining evaluated attributes, those considered to be potentially affected by livestock grazing – bank stability, bank angle, residual pool depth– remained relatively unchanged (a few of the attributes lacked data to make an evaluation). It should be noted that besides riparian vegetation, the stream attributes most directly affected by grazing activities are bank stability, bank angle, width to depth ratio, and percent undercut banks. There are too few reference sites within the John Day Basin to determine whether similar changes are occurring in unmanaged watersheds. Although only bankfull width-to-depth ratios, pool %, and bank stability showed improvement across the John Day Basin, this is actually a promising indicator given that both Bengeyfield (2006) and Rosgen (1996) indicated that the relationship between a stream’s width and depth is perhaps the most revealing of all stream channel indicators as to whether the stream is in a condition to perform the various tasks that lead to a healthy riparian area.

While some attributes, such as channel shape and the frequent floodplain, are generally formed in 1.5-2 year events, others, such as habitat complexity, are formed during moderately high events of 10-25 year return intervals. If the stream cannot maintain its dimension, pattern, and profile during these moderately high events, then habitat or other desired values will probably not be created or sustained over time. Thus, given the short 5-year time frame between site visits for monitoring and developing trends, and the analysis of data, it is our reasoned opinion that the evaluated attributes are being maintained or showing a slight overall improvement. There has been little some change in overall stream habitat conditions including improvements in bankfull width-to-depth ratios and percent fines <6mm on federal lands of the Malheur National Forest (MNF) within the Dead Horse and Fields Peak Allotments.

These results align with those found for integrator and DMA reaches as summarized in Dr. Roper’s 2009 declaration (ONDA v. Tidwell, Civ. No. 07-1871-HA, Docket #235). Based on his analysis he also stated that “it is my reasoned conclusion that the trend in the conditions of integrator reaches evaluated on the Malheur National Forest is similar or slightly better than those of other integrator reaches within the John Day Basin ... and it is my reasoned conclusion the trend in the DMA reaches within the Malheur National Forest are similar (or slightly better) than those of other DMA reaches within the John Day River Basin.”

6.2.1.1 EVALUATION OF EXISTING CONDITIONS TO PIBO MANAGED AND REFERENCE MEANS

Identifying the existing condition of streams within a particular watershed or management area is an important step in evaluating how land management may be affecting the quality of stream habitats. To help assess these conditions, we are utilizing recent information from the PIBO EMP to represent mean habitat conditions for both managed and reference conditions (see Table 19). As part of a recent publication, they developed an index of physical habitat conditions – 8 commonly collected metrics used in stream habitat monitoring – by evaluating the status of the condition of 217 reference and 934 managed streams in the Interior Columbia River and Upper Missouri River Basins (Al-Chokhachy et al. 2010). Comparing our existing condition information to these values helps provide for the evaluation of management practices to determine if they are effective in maintaining the desired and/or proper functioning condition, or improving the structure and function of riparian and aquatic conditions.

The values shown in Table 21 are not intended to replace current RMOs but to illustrate how specific streams on the MNF compare to other managed and reference stream sites within the Interior Columbia Basin. We believe this evaluation is merited as new information/articles are calling for the re-examination of certain RMOs/RMO values. Kershner and Roper (2010) found that many of the PIBO EMP reference reaches did not meet RMOs, such as wetted width-to-depth, percent undercut banks, number of pieces of large wood, and numbers of days exceeding 15°C. These authors also stated that the current RMOs were originally designed as an early warning of potential negative effects of land management on stream/riparian conditions, and values that did not meet RMOs were thought to potentially represent unsuitable habitat conditions for important salmonids. Their analysis of data from federally-managed sites in the interior Columbia River basin indicates that the usefulness of RMOs may be questionable. In summary, they found that none of the 726 reference and managed reaches surveyed met all RMOs, and in a previous analysis (Henderson et al. 2005) found that only 2% of the reference reaches met the RMO for wetted width-to-depth ratio and that 16% met the reference criteria for percent undercut banks. These may be examples of RMOs warranting exclusion or reconsideration.

Modifications in the selection of RMOs and their application are clearly needed if the results in this study are to represent conditions across the broader landscape. The authors also acknowledged that one of the drawbacks of the use of RMOs has been to disregard the role of disturbance in shaping stream habitats. Natural disturbances play an imperative role in shaping the setting of streams and the conditions that are found within them (Benda et al. 1998). They went on to say that it is apparent that all streams will most likely not meet all habitat objectives during some point in their history as the series of natural disturbances both influences and resets them. In fact, some of the PIBO reference sites come from wilderness areas that have experienced severe disturbance from wildfires and associated debris flows. These sites provide valuable information when describing the distribution of conditions that may be possible in a reference setting and provide important information on recovery trajectories in the absence of land management. Based on this information and findings, MNF staff feels it is important to evaluate conditions against selected managed and reference riparian/channel attribute values, shown in Table 21. Again, these values only represent interim aspirations and are intended as a provisional step in the measurement of movement towards existing LRMP RMOs and ESA related MPI habitat values.

TABLE 21. PIBO MANAGED AND REFERENCE MEAN VALUES FOR SELECTED STREAM, CHANNEL MORPHOLOGY, AND HABITAT ATTRIBUTES.

Stream/Channel/Habitat Attribute	PIBO Managed Mean	PIBO Reference Mean	RMSE¹	Source of Information
Bankfull W/D Ratio	23.9	22.6	4.0	Henderson et al. 2005
Percent fines sediment (<6mm)	26.7	18.0	4.9 (4.8)	Al-Chokhachy et al. 2010
Bank angle (°)	108.0	99.3	(6.5)	Al-Chokhachy et al. 2010
Percent undercut banks (%)	26.4	32.7		Al-Chokhachy et al. 2010

Residual pool depth (m):	0.26	0.31	.027 (.016)	Al-Chokhachy et al. 2010
Percent pool habitat (%)	40.9	43.3	12.9 (5.8)	Al-Chokhachy et al. 2010
Bank stability (%)	74.6	79.9		Henderson et al. 2005
D50 (mm)	43.0	58.0	13.8 (8.0)	Al-Chokhachy et al. 2010

¹ RMSE is an estimate of the potential sampling error for selected stream attributes (a measurement of the temporal variability – based on repeat measures analyses at sites over a 9-year period). Values are from Roper et al. 2010 and (Al-Chokhachy et al. 2010).

At the scale of the Deadhorse and Fields Peak Allotments, all five monitoring sites presently show bankfull width-depth ratios, bank stability, residual pool depth, and percent fines <6mm within PIBO managed mean values, while only two or fewer of the five monitoring sites exhibit managed mean values for the remaining four stream attributes: % undercut banks, bank angle, D50, and pool %. Thus, a cursory examination of the Deadhorse and Fields Peak Allotments reveals that four of the eight stream attributes considered to be potentially affected by livestock grazing – bankfull width-depth ratios, bank stability, residual pool depth, and percent fines <6mm – are found to be within PIBO managed mean values, whereas the other four, % undercut banks, bank angle, D50, and pool %, are less than managed mean values.

It should be noted that besides riparian vegetation, the stream attributes most directly affected by grazing activities are bank stability, bank angle, width to depth ratio, and percent undercut banks. As discussed earlier, a channel’s bankfull width-depth ratio is an important indicator of whether a stream is able to perform the various tasks that lead to a healthy riparian area. This indicator, along with appropriate riparian vegetation, is critically important for a stream to maintain its dimension, pattern, and profile even during moderate to high (10-25+ year return intervals) flow events, like those that occurred in 2011.

If continued monitoring shows that overall channel shape was maintained, the expected outcome will be improvement in the other stream attributes, thereby enhancing habitat complexity. Continuing to meet the allowable use standards of the BA/BiOp and LRMP, both in the form of “move triggers” and end of season minimum requirements, should avoid any negative effects to riparian or aquatic habitats that would carry over in any meaningful way to the following grazing season. In the absence of site-specific information to the contrary, it is fair to say that livestock grazing that complies with these applicable allowable use standards has a high likelihood of not meaningfully impeding the capacity for the structure and function of riparian and aquatic conditions to achieve recovery, consistent with PACFISH/INFISH S&Gs.

The results of this analysis can further be used to assist in the evaluation of other important parameters, such as water temperature in low-gradient alluvial systems, as they are directly tied to stream morphology (e.g., width/depth ratio) and vegetative cover. For example, as the streams narrow and deepen over time in an improving trajectory, water temperatures will be less susceptible to diurnal fluctuations, and as inflow from stored water in riparian banks increases, cooler water temperatures in summer and warmer temperatures in winter would be expected. Thus, the evaluation of trend in stream morphology attributes such as width/depth ratio can help inform the Line Officer or Manager about other important fish habitat elements.

6.2.2 IMPLEMENTATION MONITORING

An interdisciplinary group using the Multiple Indicator Monitoring (MIM) method developed a monitoring program for pastures on the Malheur National Forest. Because permittees generally rested pastures that overlapped the ESA action area in the Seneca, Deadhorse, Hanscomb, and Fields Peak Allotments over the last 3 years (2008-2010), no monitoring was conducted in the riparian sites during this time span.

7 EFFECTS OF THE PROPOSED ACTION

The direct and indirect effects of implementing the action, including interrelated and interdependent actions, on the listed species and designated CH are evaluated in this section. In addition, the probability of directly affecting juveniles, spawning adults, and incubating embryos in redds will be assessed. The environmental impacts of implementing the project elements (PE) will be evaluated by use of NMFS MPI indicators to determine effects to ESA-listed MCR Steelhead and designated CH.

As described in this document, the proposed action is expected to allow previously degraded riparian areas/habitat indicators to continue recovery. However, it is anticipated that the proposed grazing activities in all cases will maintain the current environmental baseline condition for each indicator. In some cases indicators are rated as Not Properly Functioning, which suggests that the proposed grazing activities will be maintaining this risk rating. However, because the environmental baseline rating is determined at the subbasin scale, the proposed grazing activities tend to influence only portions of subbasins, and watershed restoration activities needed to improve the baseline indicators at the subbasin scale will not likely occur over the life of this consultation, it is anticipated that the proposed grazing activities will maintain the current environmental baseline condition. Historic and current cattle grazing in the ESA action area likely plays varying roles in the current environmental baseline ratings for these affected subbasins.

7.1 PROJECT ELEMENT AND INTERRELATED ACTION EVALUATION

The component parts of the action are listed in Section 4.1 as six project elements and are also shown below.

1. Livestock use of allotment/pastures
2. Permittee management of livestock and infrastructure maintenance
3. Range improvements
4. Exclusionary fences
5. Monitoring
6. Adaptive management

We determined that unauthorized use (trespass) is not an action. However, the implementation of FS enforcement actions regarding unauthorized use is an interrelated action.

7.1.1 PROJECT ELEMENTS DROPPED FROM FURTHER ANALYSIS

An initial step in the analysis process is to determine if any of the project elements are already provided ESA coverage in a concluded programmatic consultation. The consultation history section (Section 1.1) described the Blue Mountain Expedited Section 7 Consultation (BMESSC) programmatic consultation. Range improvements are covered under that consultation. Range improvements in the BMESSC concurrence letter are described as: “e.g. fencing, off-site water developments.” The consultation history section also described the Aquatic and Riparian Restoration Programmatic Consultation (ARRPC). Riparian exclusion fencing with water gaps and stream crossings is a category covered under the ARRPC biological opinion. Consequently, PEs 3 and 4 below already have existing ESA coverage and will not be further evaluated in this BA.

7.1.2 PROJECT ELEMENTS AND INTERRELATED ACTIONS WITH ENTIRELY BENEFICIAL EFFECTS

PE 6, adaptive management, provides a mechanism to adjust management if end-point indicators and desired conditions are not being met. Examples of adaptive management measures include reducing livestock numbers, changing the timing and duration of grazing, adjusting the numeric end-point indicators and constructing more exclusion fences. Making adjustments to ensure that end-point indicators and desired conditions are met will result in positive effects to habitat indicators and therefore to CH. The results would also have beneficial effects to the species, as many adaptive management adjustments will reduce the time that livestock are in or adjacent to streams.

Law enforcement actions to remove cattle not under permit will result in entirely beneficial effects to the species and designated CH.

7.1.3 PROJECT ELEMENTS REMAINING FOR ANALYSIS

Of the six PEs initially developed for this livestock grazing consultation, PEs 3 and 4 have been addressed as already covered by existing programmatic consultations still in effect, and the effect of implementing PE 6 has been determined to be entirely beneficial to CH and to the species. The PEs remaining for analysis are:

1. Livestock use of allotment/pastures
2. Permittee management of livestock and infrastructure maintenance
5. Monitoring

7.1.3.1 PE1: LIVESTOCK USE OF ALLOTMENT/PASTURES

Livestock will graze the allotment and individual pastures in the numbers, time frames and locations described in the proposed action section and in the term grazing permit.

7.1.3.2 PE2: PERMITTEE MANAGEMENT OF LIVESTOCK AND INFRASTRUCTURE MAINTENANCE

This PE includes the move-in and move-out of livestock using highway and off-road vehicles, and herding by range riders. While vehicles are also used to access sites for monitoring purposes

(PE 5), the effects of vehicle use to CH and to the species will only be assessed for this PE to reduce redundancy in the analysis. Project Design Criteria for vehicle use are provided by the PDCs described earlier in the proposed action section.

Several hundred troughs, springs and ponds are maintained by grazing permittees to provide off-stream water for livestock across the MNF. In addition, there are miles of fence and dozens of gates that are maintained each year. Typical maintenance activities involve the use of hand tools or machines on a small footprint of land. Some work such as repairing troughs or replacing wire will not involve any soil or vegetation disturbance. Other maintenance activities may disturb small amounts of soil and vegetation, but rarely within riparian areas adjacent to MCR Steelhead CH. Workers performing maintenance activities rarely walk in riparian areas or in stream channels where listed fish are present or in designated CH.

7.1.3.3 PE5: MONITORING

A variety of implementation and effectiveness monitoring techniques are employed to determine if desired conditions are being met. The MNF Riparian Monitoring Strategy is discussed in detail in the Monitoring section (Section 4.1.4). Workers use manual and electronic equipment to measure vegetation, water quality and stream channel/streambed characteristics. Some monitoring actions include wading in stream channels.

7.2 ANALYSIS OF EFFECTS TO DESIGNATED CRITICAL HABITAT

The three PEs will be analyzed first for their effects to designated CH, then for effects to the species. The freshwater primary constituent elements (PCE) of MCR Steelhead CH applicable to the action area are presented in Table 22.

TABLE 22. PRIMARY CONSTITUENT ELEMENTS OF MCR STEELHEAD CRITICAL HABITAT APPLICABLE TO THE ACTION AREA.

PCE	Description
1	Freshwater spawning sites with water quantity and quality conditions and substrate supporting spawning, incubation and larval development.
2	Freshwater rearing sites with: (i) Water quantity and floodplain connectivity to form and maintain physical habitat conditions and support juvenile growth and mobility; (ii) Water quality and forage supporting juvenile development; and (iii) Natural cover such as shade, submerged and overhanging large wood, log jams and beaver dams, aquatic vegetation, large rocks and boulders, side channels, and undercut banks.
3	Freshwater migration corridors free of obstruction and excessive predation with water quantity and quality conditions and natural cover such as submerged and overhanging large wood, aquatic vegetation, large rocks and boulders, side channels, and undercut banks supporting juvenile and adult mobility and survival.

The effects to each PCE, and ultimately to designated CH as a whole, can be determined by evaluating the effects to indicators of the NMFS MPI that correspond to each PCE. The MNF uses a crosswalk table format for this purpose. Table 23 presents the analysis for effects of the action to the PCEs of MCR Steelhead designated CH. Table 24 presents a summary of effects to the indicators associated with each PCE of MCR Steelhead CH. Measurable effects to several habitat indicators of PCEs were concluded.

TABLE 23. ANALYSIS OF EFFECTS TO MPI INDICATORS CORRESPONDING TO PCEs OF DESIGNATED CRITICAL HABITAT FOR MCR STEELHEAD WITHIN THE SENECA, DEADHORSE, HANSCOMB, AND FIELDS PEAK ALLOTMENTS.

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
(1) Freshwater spawning sites with water quantity and quality conditions and substrate supporting spawning, incubation and larval development;	Water Quantity	Flow/ Hydrology	Changes in Peak/Base Flows	<p>Riparian vegetation has been linked to the water-holding capacity of streamside aquifers (Platts 1991). As riparian vegetation is removed by grazing and streamside soils are compacted by hooves, the ability of areas to retain water is decreased. Decreased evapotranspiration and infiltration increase and hasten surface runoff, resulting in a more rapid hydrologic response of streams to rainfall. When this occurs, high flows in the spring tend to increase in volume, leading to bank damage and erosion, and channel downcutting. Summer and fall base flows are decreased, often resulting in flows that are insufficient to provide suitable rearing habitat for juvenile salmonids. If aquifers lose their capacity to hold and slowly deliver water to the stream, differences between peak and base discharge rates increase dramatically (USEPA 1993). Some streams that typically flowed perennially may experience periods of no flow in the summer or fall. Li <i>et al.</i> (1994) found that flow in a heavily grazed eastern Oregon stream became intermittent during the summer, while a nearby, well-vegetated reference stream in a similar-sized watershed had permanent flows. They suggested that the difference in flow regimes was due to diminished interaction between the stream and floodplain with resultant lowering of the water table.</p> <p>Indirect effects of historic livestock grazing in the ESA action area (including trailing and watering), on channel and bank features such as bank stability, undercut banks and width to depth ratio, as well as impacts to shrub recruitment and green line plant vigor, have likely affected peak and base flows on some streams. It is anticipated that PE 1 (livestock use) will have negative effects to this indicator, but they will be too small to be meaningfully measured, particularly to flows at the time of year when spawning, incubation and larval development occur. The use of BMPs, end point indicators, and adaptive management should minimize effects. If hydrophytic vegetation, bank stability, width-depth ratio, and undercut banks show a static and/or downward trend and the Forest is not meeting RMOs, grazing practices will be modified (See Adaptive Management Section VI). PE 2 (permittee mgt. and mtce.) includes off-road vehicle use. This has the potential to increase soil compaction, but it will be minimized by use of PDCs. Little to no riparian vegetation is affected by vehicle use, range riding or maintenance activities. PE 2 overall will have slight negative effects to the indicator that are too small to be meaningfully measured. PE5 (monitoring) will not increase compaction or remove vegetation, and therefore does not have</p>

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
				a mechanism to affect peak/base flow. The effect to the indicator is neutral.
			Increase in Drainage Network	None of the PEs has road construction, so no change to the drainage network will occur. The proposed action would have a neutral effect on the indicator.
			Temperature	<p>The temperature monitoring data for the Buck Cabin and Wickiup Creek monitoring sites met state water quality standards but rated NPF using the NMFS MPI criteria (see Section 6.1.3). Temperature monitoring data for the Tex Creek and Fields Creek monitoring sites failed state water quality standards and were also rated NPF using the NMFS MPI criteria.</p> <p>Interpretation of utilization monitoring results from pastures that were rested from livestock grazing in or adjacent to the ESA action area suggest that shade-producing vegetation along stream channels is being grazed or browsed to some degree by wild ungulates. However, this use has been light.</p> <p>Many grass/grass-like species found on the MNF have an ungrazed potential height of 2 to 3 feet (MNF 2007). In meadow streams with narrow channels, they often are the plants that provide stream shade. PE 1 (livestock use) will potentially reduce vegetation heights to 4 or 6 inches. This will considerably reduce stream shade in those circumstances compared to the ungrazed potential vegetation heights (see discussion that follows in Effects to Listed Species section).</p> <p>Livestock use (PE 1) is likely to result in measurable water temperature increases for certain stream reaches. These impacts are expected to be generally confined to low gradient stream channels less than 10 feet wide with grass/grass-like vegetation providing shade. The Tex Creek and Murderers Creek MSRAs are areas where small temperature increases are probable. The effect to this indicator by livestock use is negative and meaningfully measured for the Fields Peak and Hanscomb Allotments. Critical Habitat streams draining the Seneca and Deadhorse allotments such as Riley, Ingle, and Vance creeks, are higher gradient, well-shaded with timbered riparian zones or have low likelihood of cattle interaction. Therefore effects to the temperature indicator in these allotments would be negative but insignificant.</p>

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
				<p>Water temperatures typically are below concern thresholds when spawning, incubation and larval development of MCR Steelhead occurs, as flows are greater than later in the year. PE 2 (permittee livestock management and infrastructure maintenance) and PE 5 (monitoring) activities will not remove vegetation that provides shade nor affect channel-forming processes that might widen stream channels. Consequently, there is no mechanism for PEs 2 and 3 to affect water temperature and the effect of the PE for the indicator is neutral.</p> <p>Livestock grazing on federal land in the ESA action area is managed to attain the endpoint indicators, which were developed to meet PACFISH grazing standards and guidelines as well as water quality BMPs. The assumption is that meeting these endpoint indicators would move key riparian and stream channel elements (bank stability, W/D ratio, woody species regeneration) towards their Desired Conditions and meet Riparian Objectives, and thereby maintain water temperatures. If monitoring fails to show this trend, adaptive management would be implemented and endpoint indicators would be modified to minimize adverse impacts to this element of the PCE.</p>
			Sediment/ Turbidity	<p>The monitoring results presented above indicate that livestock use (PE 1), as well as use by wild ungulates, results in trampled and grazed riparian vegetation, and altered stream banks to some degree. Livestock also use trails to access streams for water. Livestock occasionally will concentrate their use in certain areas, potentially creating patches of relatively bare soil. Some of these areas may be adjacent to stream sections used by MCR Steelhead for spawning, incubation and larval development, such as Tex and Murderers creeks. Bare soil is prone to erosion and can result in fine sediment entering stream channels and resultant increases in turbidity. Habitat impacts are likely to include areas of exposed streambank up to a few feet wide where livestock access streams to drink or cross, and areas of bank disturbance where livestock graze in riparian areas. Exposed areas and other bank disturbances that occur are likely to result in a slight increase in turbidity for a short distance downstream during rainstorms or runoff events. However, given background levels of turbidity during runoff events it will be difficult to distinguish between turbidity resulting from these grazing impacts and background turbidity. A slight increase in fine sediment deposition for a short distance downstream of exposed and disturbed areas is also likely to occur.</p>

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
				<p>Endpoint indicators were developed in order to meet PACFISH grazing standards and guidelines as well as water quality BMPs. The assumption is that meeting these endpoint indicators would move key riparian and stream channel elements (bank stability, w/d ratio, woody species regeneration) towards their Desired Conditions and meet our Riparian Objectives. If monitoring fails to show this trend, adaptive management would be implemented (Section 4.1.5) and endpoint indicators would be modified to minimize adverse effects to critical habitat.</p> <p>However, livestock grazing will increase the amount of sediment entering streams by the mechanisms described above, particularly in streams where MSRAs exist such as Tex, Murderers, and Laycock Creeks. Laycock Creek is already dominated by sandy substrate and has unstable banks. These impacts are expected to be localized and short-term. Consequently, the effect to this indicator by PE1 (livestock use) is negative and expected to be measurable for the Fields Peak Allotment using bank alteration methods described above as a proxy for turbidity and sedimentation. Critical Habitat streams draining the Seneca and Deadhorse allotments such as Riley, Ingle, and Vance creeks, are higher gradient, with cobble substrate and armored banks, have low likelihood of cattle interaction because of brush or abundant instream large wood. Therefore effects to the temperature indicator in these allotments would be negative but insignificant</p> <p>PE 2 involves use of vehicles on and off roads, as well as infrastructure maintenance. There is the potential for fine sediment to be transported from unpaved roads to stream channels, primarily at road crossings, during rainstorms or runoff events. However, it is impossible to determine the proportion of the suspended sediment attributable to road use by permittees, given the use of the roads for other purposes. In addition, background levels of suspended sediment in streams will be high during rainstorms and runoff events, and the contribution by permittee use of roads to increased turbidity cannot be meaningfully measured. Use of off-road vehicles should not result in measurable effects due to use of PDCs. Range riding with horses will not cause any meaningfully measured increases in streambed sediment or turbidity. Maintenance activities are typically distant from designated CH, disturb little to no soil, and are not hydrologically connected to stream channels. There is no mechanism for maintenance activities to affect the indicator. Overall, the effects of PE 2 to the indicator are</p>

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
				<p>negative and not meaningfully measured.</p> <p>Monitoring (PE 3) activities such as pebble counts and measuring cross-sections involve wading in stream channels. Other monitoring activities involve walking or riding horses in riparian areas. The timing of these activities is typically after spawning, incubation and larval development of MCR Steelhead, although there may be some overlap in timing. Spawning surveys also involve wading. Wading may result in very small increases in turbidity downstream for a short distance (a few feet) that will quickly dissipate. Walking and riding horses in riparian areas should not result in fine sediment delivery to stream channels. However, there may be very small and transient increases in turbidity when a stream is being crossed. The monitoring PE effect to the indicator is negative, but not meaningfully measured.</p>
			<p>Chemical Contamination/ Nutrients</p>	<p>Urine and feces from livestock use (PE 1) in riparian areas increases the likelihood that nitrogen and phosphorous will enter streams. Increased nutrients will likely increase stream productivity at the source of nutrients and for a short distance downstream. It is anticipated that livestock grazing will have slight negative impacts to the indicator, but they are not expected to be meaningfully measured.</p> <p>PE 2 (permittee management and infrastructure maintenance) includes vehicle use. The risk of chemical contamination to streams will be minimized by use of PDC. Maintenance activities are typically distant from designated CH, and at locations not hydrologically connected to stream channels. Therefore, there is no mechanism for petroleum products spilling from power tools to affect CH. Use of horses for range riding will have similar effects (but much smaller scale) than that of PE 1, above. Maintenance activities are typically distant from stream channels. The overall effect of PE 2 is for slight negative effects to the indicator that are not expected to be meaningfully measured.</p> <p>Monitoring (PE 5) does not involve the use of chemicals and does not have the potential to affect nutrients in streams. PE 5 will have a neutral effect to the indicator.</p>

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
	Suitable Substrate	Habitat Elements	Substrate Embeddedness	<p>The analysis of effects to the sediment/turbidity indicator, above, determined that use of riparian areas by livestock is expected to increase the amount of sediment entering streams. A slight increase in fine sediment deposition for a short distance downstream of exposed and disturbed areas is likely to occur. There is the potential for fine sediment to slightly increase embeddedness within gravels suitable for spawning when the gravel is located immediately downstream from exposed and disturbed streambank areas. The effect to this indicator by livestock use (PE 1) is negative and meaningfully measurable, using bank alteration survey protocol as a proxy for embeddedness. The analysis for sediment/turbidity determined that PE 2 would have a slightly negative, but not meaningfully measured effect to the indicator. Therefore, the same conclusion is made for the substrate embeddedness indicator. As described above, monitoring (PE 5) would not introduce fine sediment into stream channels. The monitoring PE will have a neutral effect to the indicator.</p>
(2) Freshwater rearing sites with: (i) Water quantity and floodplain connectivity to form and maintain physical habitat conditions and support	Water Quantity	Flow/ Hydrology	Changes in Peak/Base Flows	See discussion above.
			Increase in Drainage Network	See discussion above.
	Water Quality	Water Quality	Temperature	See discussion above. The rearing period includes the summer months when elevated water temperatures are most concerning for juvenile salmonids, and the sun's position in the sky results in the greatest potential for increased solar radiation to streams. It is this time period when the small, but measurable, increases to water temperature would take place.
			Sediment/ Turbidity	See discussion above.

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
juvenile growth and mobility; (ii) Water			Chemical Contamination/ Nutrients	See discussion above. The conclusion was for a slight negative effect to the indicator from livestock use (PE 1) and permittee management and infrastructure maintenance (PE 2) since there would be an increase in nutrients into streams. However, the introduction of nutrients may lead to small increases in contamination from fecal coliform.

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
<p>quality and forage supporting juvenile development; and</p> <p>(iii) Natural cover such as shade, submerged and overhanging large wood, log jams and beaver dams, aquatic vegetation, large rocks and boulders, side channels, and undercut banks</p>	<p>Floodplain Connectivity</p>	<p>Channel Condition and Dynamics</p>	<p>Floodplain Connectivity</p>	<p>Channel entrenchment is the main concern for loss of floodplain connectivity. Indirect effects of livestock use (PE 1), including trailing and watering, on things such as bank stability, undercut banks, width depth ratio, shrub recruitment, and green line plant vigor have limited some streams' ability to access their floodplains, thus concentrating energies within confined channels and causing additional erosion. Many of these streams are still experiencing this phenomenon.</p> <p>Channel entrenchment as a result of livestock use (PE 1) will be prevented by use of endpoint indicators to meet PACFISH grazing standards and guidelines as well as water quality BMPs. The conclusion is that the effect to the indicator by livestock use is negative but not meaningfully measured. The assumption is that meeting these endpoint indicators would move key riparian and stream channel elements (bank stability, w/d ratio, woody species regeneration) towards their Desired Conditions and meet Riparian Objectives. If monitoring fails to show this trend, adaptive management would be implemented and endpoint indicators would be modified to minimize negative effects to floodplain connectivity.</p> <p>PE 2 (permittee management and infrastructure maintenance) includes on and off road vehicle use. Road use has no mechanism to affect floodplain connectivity. PDC for off-road use will prevent channel downcutting. Range riding with horses will occasionally cross a stream but effects to streambanks and beds will be so minimal as to not affect the indicator. Infrastructure maintenance actions do not affect streambanks or riparian vegetation adjacent to CH, and will therefore not affect floodplain connectivity. The overall effect of PE 2 is a neutral affect to the indicator.</p> <p>Monitoring (PE 5) does not remove riparian vegetation or otherwise have mechanisms to destabilize stream channels. PE 5 will have a neutral effect to the indicator.</p>

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
			Width/Depth Ratio	<p>Eight level II stream surveys reported bankfull width-depth ratios ranging from 3.4 to 10.6, with 16 of 19 stream reaches meeting the NMFS MPI criterion for PF (<10) (see Appendix J for PIBO and stream survey monitoring data). Data from the two PIBO integrator sites shows a slight decrease in W/D ratio between 2003 and 2008 at the Riley Creek site (16 to 15) while it decreased substantially at the Fields Creek site (29 to 23). Livestock use (PE 1) is anticipated to have a negative, but not meaningfully measured effect to the indicator. The potential for increases in width-depth ratio is reduced because of implementation of endpoint indicators for livestock grazing (which includes use by wild ungulates) and adaptive management. McDowell and Mowry (2002) found little difference in width to depth ratios between grazed areas and exclosures along Tex Creek, but did find a lower w:d ratio in exclosed portions of Murderers Creek. PE 2 (permittee management and infrastructure maintenance) includes on and off road vehicle use. Road use has no mechanism to affect W/D ratio. PDC for off-road use will prevent bank damage and effects to W/D ratio. Range riding with horses will occasionally cross a stream but effects to streambanks and beds will be so minimal as to not affect the indicator. Infrastructure maintenance actions do not affect streambanks or riparian vegetation adjacent to CH, and will therefore not affect W/D ratio. The overall effect of PE 2 is a neutral effect to the indicator. PE 5 (monitoring) does not remove vegetation or destabilize stream banks. There is no potential for it to increase W/D ratio. The monitoring PE will have a neutral effect to the indicator.</p>
	Forage	Habitat Elements	Substrate Embeddedness	<p>See discussion above for this indicator for the suitable substrate PCE habitat feature. The conclusion for livestock use (PE 1) was that a slight increase in fine sediment deposition for a short distance downstream of exposed and disturbed areas is likely to occur. This would result in small areas of increased embeddedness. Increased embeddedness may result in a decrease in the potential for production of aquatic macroinvertebrates (a forage item for rearing salmonids) in small, isolated patches. The conclusion is that PE 1 will have a slight negative effect on substrate embeddedness with respect to the production of forage. Consistent with the analysis for the suitable substrate PCE habitat feature, the effect of PCE 2 to the indicator is negative but not meaningfully measured, and the effect of PCE 5 is neutral.</p>

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
			Large Woody Debris	<p>Livestock grazing does not affect this indicator in conifer-dominated riparian forests. Livestock use can negatively affect this indicator when grazing occurs within hardwood stands such as aspen, alder, birch, and cottonwoods that could contribute larger pieces of wood to small streams. In sites in the action area that would be naturally dominated by cottonwood gallery riparian forests, livestock use (PE 1) will likely result in altering the level of cottonwood stocking and future large tree (and subsequent large woody debris) recruitment (Kaufmann et al. 1983, Case and Kaufmann 1997, Beschta and Ripple 2005). Therefore it is anticipated that livestock will graze young cottonwoods at levels meaningfully measured with respect to the future production of large woody debris. The effect to this indicator by livestock use (PE 1) is negative and meaningfully measured in the Fields Peak allotment because of the presence of meadow systems that could support cottonwood growth. Critical Habitat streams draining the Seneca and Deadhorse allotments such as Riley, Ingle, and Vance creeks, are higher gradient, well-shaded with mixed conifer, hardwood (Vance) or ponderosa pine riparian zones or have low likelihood of cattle interaction. Therefore effects to the large woody debris indicator in these allotments would be negative but insignificant. Using BMPs, end point indicators, and adaptive management will result in discouraging browse on existing hardwoods and willows but may not promote regeneration of new cottonwoods. PE 2 and PE 5 do not affect trees and associated LWD in any way. Therefore there is no mechanism for an effect and the effect is neutral to the indicator for both PEs</p>

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
			Pool Frequency	<p>Pool frequency at the PIBO monitoring sites and stream survey reaches within the allotments are not currently meeting the DFC as described within Amendment 29 of the MNF LRMP and would be NPF using NMFS MPI criteria. See Appendix J for PIBO and stream survey results.</p> <p>Indirect effects of livestock grazing (including trailing and watering), on bank stability, undercut banks, width-depth ratio, shrub recruitment, green line plant composition and vigor have the potential to affect this indicator. The use of BMP's for livestock management, end point indicators and adaptive management, should result in an overall effect by PE 1 (livestock use) to pool frequency that is not meaningfully measured and unlikely to occur.</p> <p>PE 2 (permittee management and infrastructure maintenance) includes on and off road vehicle use. Road use has no mechanism to affect pool frequency. PDC for off-road use will prevent bank damage and effects to pool frequency. Range riding with horses will occasionally cross a stream but effects to streambanks will be so minimal as to not affect the indicator. Infrastructure maintenance actions do not affect streambanks or riparian vegetation adjacent to CH, and will therefore not affect pool frequency. The overall effect of PE 2 is a neutral effect to the indicator.</p> <p>PE 5 (monitoring) does not have any mechanisms to affect plants or bank and channel features that would impact pool frequency. The monitoring PE has a neutral effect to the indicator.</p>

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
			Pool Quality	<p>PIBO data within the allotments (n=6) indicate that undercut banks are at less than the desired condition. Residual pool depth is generally low with very few to no pools greater than 1 meter deep in surveyed stream reaches. See Appendix J for PIBO and stream survey results. Based upon the PIBO and stream survey data, pool quality would be considered to be NPF using NMFS MPI criteria.</p> <p>Indirect effects of livestock grazing (including trailing and watering), on bank stability, undercut banks, width-depth ratio, shrub recruitment, green line plant composition and vigor have the potential to affect this indicator. The use of BMPs for livestock management, end point indicators (which are inclusive of wild ungulate use), and adaptive management, should result in an overall effect by PE 1 (livestock use) to pool quality that is negative and not meaningfully measured.</p> <p>PE 2 (permittee management and infrastructure maintenance) includes on and off road vehicle use. Road use has no mechanism to affect pool quality. PDC for off-road use will prevent bank damage and effects to pool quality. Range riding with horses will occasionally cross a stream but effects to streambanks will be so minimal as to not affect the indicator. Infrastructure maintenance actions do not affect streambanks or riparian vegetation adjacent to CH, and will therefore not affect pool frequency. The overall effect of PE 2 is a neutral effect to the indicator.</p> <p>PE 5 (monitoring) does not have any mechanisms to affect plants or bank and channel features that would impact pool quality. The monitoring PE has a neutral effect to the indicator.</p>

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
			Off-Channel Habitat	<p>There is very little off-channel habitat within tributaries of the UMJD River. Of 19 level II stream surveys that reported side-channel information, all except one reported less than 5% of the stream reaches contained side channels or off-channel area. The use of BMP's for livestock management, end point indicators (which are inclusive of wild ungulate use), and adaptive management, should result in an overall effect by PE 1 (livestock use) to off-channel habitat that is negative and not meaningfully measured.</p> <p>PE 2 (permittee management and infrastructure maintenance) includes on and off road vehicle use. Road use has no mechanism to affect off-channel area. PDC for off-road use will prevent bank damage and effects to off-channel habitat. Range riding with horses will occasionally cross a stream but effects to streambanks will be so minimal as to not affect the indicator. Infrastructure maintenance actions do not affect streambanks or riparian vegetation adjacent to CH, and will therefore not affect off-channel areas. The overall effect of PE 2 is a neutral affect to the indicator.</p> <p>PE 5 (monitoring) does not have any mechanisms to affect off-channel habitat. The monitoring PE has a neutral effect to the indicator.</p>

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
			Refugia	<p>The availability of refugia is a limiting factor identified in the recovery plan for the Oregon steelhead population of the MCR Steelhead distinct population segment (NMFS 2009). The NMFS MPI (NMFS 1996) defines the refugia indicator as: “important remnant habitat for sensitive aquatic species.” All of the habitat indicators in this crosswalk table are potential components of Refugia. Analysis for previous indicators has determined that PE 1 (livestock use) will have negative and meaningfully measured effects to several of them. This may occur in areas that meet the definition of Refugia. Therefore, PE 1 (livestock use) will have negative and meaningfully measured or evaluated effects to the refugia indicator for the Fields Peak and Hanscomb allotments.</p> <p>The highest level of effect to previous indicators by PE 2 (permittee management and infrastructure maintenance) was “negative but not meaningfully measurable.” This level of effects will not impact the function of Refugia to provide important remnant habitat. Therefore, the effect conclusion is neutral for PE 2.</p> <p>The highest level of effect to previous indicators by PE 5 (monitoring) was “negative but not meaningfully measurable” for small and transient increases in turbidity by wading in stream channels or crossing streams on foot or by horse. This level of effects will not impact the function of Refugia to provide important remnant habitat. Therefore, the effect conclusion is neutral for the monitoring PE.</p>

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
		Watershed Condition	Riparian Reserves	<p>As described above, PE 1 (livestock use) will result in negative effects to indicators within riparian areas. A negative effect to Riparian Management Areas (RMA) (east-side analog of Riparian Reserves) is indicated. However, the negative effects should not rise to the level that impacts to the processes and functions of RMAs are meaningfully measurable. Endpoint indicators were developed with seral class in mind to meet PACFISH grazing standards and guidelines, enclosure B of the LMRP and water quality BMPs. The assumption is that meeting these endpoint indicators would move key riparian and stream channel elements (bank stability, w/d ratio, woody species regeneration) towards their Desired Conditions and meet Riparian Objectives. If monitoring fails to show this trend, adaptive management would be implemented and endpoint indicators would be modified to minimize adverse effects to Riparian Reserves.</p> <p>The highest level of effect to previous indicators by PE 2 (permittee management and infrastructure maintenance) was “negative but not meaningfully measurable.” This level of effects will not impact the processes and functions of RMAs. Therefore, the effect conclusion is neutral for PE 2.</p> <p>The monitoring PE does not have any mechanisms to affect the processes and functions of RMAs. The monitoring PE has a neutral effect to the indicator.</p>

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
	Natural Cover	Habitat Elements	Substrate	<p>This indicator focuses on the composition of streambed substrate rather than its embeddedness, which was evaluated earlier. Stream survey data indicated that 6 of 19 stream reaches within the allotments have substrate that is dominated by sand. See Substrate Embeddedness indicator above. Based on the stream survey data the Substrate indicator would be classified as NPF using NMFS MPI criteria.</p> <p>The analysis of effects to the sediment/turbidity indicator for PCE 1, above, determined that use of riparian areas by livestock is expected to increase the amount of sediment entering streams. A slight increase in fine sediment deposition for a short distance downstream of exposed and disturbed areas is likely to occur. However, this is not expected to measurably change the composition of existing substrate with regard to its function as cover for juvenile or adult MCR Steelhead. Therefore, the effect to this indicator by PE 1 (livestock use) is negative and not meaningfully measurable. The use of BMP's for livestock management, end point indicators (which are inclusive of wild ungulate use), and adaptive management, should further minimize the magnitude of potential negative effects by PE 1.</p> <p>The analysis of effects to the sediment/turbidity indicator for PCE 1, above, determined that the effect of PE 2 (permittee management and infrastructure maintenance) was "negative and not meaningfully measured." This level of effects is not expected to measurably change the composition of existing substrate with regard to its function as cover for juvenile or adult MCRS Steelhead. Therefore, the effect to this indicator by PE 1 (livestock use) is negative and not meaningfully measurable.</p> <p>As described above, PE 5 (monitoring) would not introduce fine sediment into stream channels. The monitoring PE will have a neutral effect to the indicator.</p>
			Large Woody Debris	See Above.
			Pool Frequency	See Above.

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
			Pool Quality	See Above
			Off-Channel Habitat	See Above
			Refugia	See Above
		Watershed Condition	Riparian Reserves	See Above

PCE	PCE Habitat Feature	Matrix Pathway	Matrix Indicator	Rationale
(3) Freshwater migration corridors free of obstruction and excessive predation with water quantity and quality conditions and natural cover such as submerged and overhanging large wood, aquatic vegetation, large rocks and boulders, side channels, and undercut banks supporting juvenile and adult mobility and survival;	Migration Corridors Free of Obstruction	Habitat Access	Physical Barriers	No barriers will be created or removed by the actions of any PE. All PEs have a neutral effect on the physical barriers indicator.

TABLE 24. SUMMARY OF EFFECTS OF THE PROPOSED ACTION BY THE PROJECT ELEMENTS OF LIVESTOCK GRAZING IN THE SENECA, DEADHORSE, HANSCOMB, AND FIELDS PEAK ALLOTMENTS TO THE INDICATORS ASSOCIATED WITH HABITAT FEATURES OF EACH PRIMARY CONSTITUENT ELEMENT OF MCR STEELHEAD CRITICAL HABITAT.

Primary Constituent Element	PCE Habitat Feature	Indicator	Effect Conclusion by Project Element		
			PE1: Livestock Use	PE2: Permittee Management of Livestock and Infrastructure Maintenance	PE 5: Monitoring
1. Freshwater spawning sites with water quantity and quality conditions and substrate supporting spawning, incubation and larval development	Water quantity	Changes in Peak/Base Flows	NNMM ¹	NNMM	Neutral
		Increase in Drainage Network	Neutral	Neutral	Neutral
	Water quality	Temperature	<i>Fields Peak/Hanscomb: NMM²</i> Seneca/Deadhorse: NNMM	Neutral	Neutral
		Sediment/Turbidity	<i>Fields Peak/Hanscomb: NMM</i> Seneca/Deadhorse: NNMM	NNMM	NNMM
		Chemical Contamination/Nutrients	NNMM	NNMM	Neutral
	Suitable substrate	Substrate Embeddedness	<i>Fields Peak/Hanscomb: NMM</i> Seneca/Deadhorse: NNMM	NNMM	Neutral
2. Freshwater rearing sites with:	Flow/hydrology	Changes in Peak/Base Flows	NMM	NNMM	Neutral

Primary Constituent Element	PCE Habitat Feature	Indicator	Effect Conclusion by Project Element		
			PE1: Livestock Use	PE2: Permittee Management of Livestock and Infrastructure Maintenance	PE 5: Monitoring
<p>(i) Water quantity and floodplain connectivity to form and maintain physical habitat conditions and support juvenile growth and mobility;</p> <p>(ii) Water quality and forage supporting juvenile development; and</p> <p>(iii) Natural cover such as shade, submerged and overhanging large wood, log jams and beaver dams, aquatic vegetation, large rocks and boulders, side channels, and undercut banks</p>		Increase in Drainage Network	Neutral	Neutral	Neutral
	Water quality	Temperature	<i>Fields Peak/Hanscomb: NMM</i> Seneca/Deadhorse: NNMM	Neutral	Neutral
		Sediment/Turbidity	<i>Fields Peak/Hanscomb: NMM</i> Seneca/Deadhorse: NNMM	NNMM	NNMM
		Chemical Contamination/Nutrients	NNMM	NNMM	Neutral
	Floodplain connectivity	Floodplain Connectivity	NNMM	Neutral	Neutral
		Width/Depth Ratio	NNMM	Neutral	Neutral
	Forage	Substrate Embeddedness	<i>Fields Peak/Hanscomb: NMM</i> Seneca/Deadhorse: NNMM	NNMM	Neutral
		Large Woody Debris	<i>Fields Peak/Hanscomb: NMM</i> Seneca/Deadhorse:	Neutral	Neutral

Primary Constituent Element	PCE Habitat Feature	Indicator	Effect Conclusion by Project Element		
			PE1: Livestock Use	PE2: Permittee Management of Livestock and Infrastructure Maintenance	PE 5: Monitoring
			NNMM		
		Pool Frequency	NNMM	Neutral	Neutral
		Pool Quality	NNMM	Neutral	Neutral
		Off-Channel Habitat	NNMM	Neutral	Neutral
		Refugia	<i>Fields Peak/Hanscomb: NMM</i> Seneca/Deadhorse: NNMM	Neutral	Neutral
		Riparian Reserves	NNMM	Neutral	Neutral
	Natural cover	Substrate	NNMM	NNMM	Neutral
		Large Woody Debris	<i>Fields Peak/Hanscomb: NMM</i> Seneca/Deadhorse: NNMM	Neutral	Neutral
		Pool Frequency	NNMM	Neutral	Neutral
		Pool Quality	NNMM	Neutral	Neutral
		Off-Channel Habitat	NNMM	Neutral	Neutral
		Refugia	<i>Fields Peak/Hanscomb:</i>	Neutral	Neutral

Primary Constituent Element	PCE Habitat Feature	Indicator	Effect Conclusion by Project Element		
			PE1: Livestock Use	PE2: Permittee Management of Livestock and Infrastructure Maintenance	PE 5: Monitoring
			<i>NMM</i> Seneca/Deadhorse: NNMM		
		Riparian Reserves	NNMM	Neutral	Neutral
3. Freshwater migration corridors free of obstruction and excessive predation with water quantity and quality conditions and natural cover such as submerged and overhanging large wood, aquatic vegetation, large rocks and boulders, side channels, and undercut banks supporting juvenile and adult mobility and survival	Migration corridors free of obstruction	Physical Barriers	Neutral	Neutral	Neutral

¹NNMM = Negative, not meaningfully measured

²*NMM* = Negative, meaningfully measured

7.3 ANALYSIS OF EFFECTS TO LISTED SPECIES

7.3.1 GENERAL EFFECTS

Effects to MCR steelhead from livestock grazing can be in the form of direct impacts to individual fish or indirectly through habitat disturbance. Direct disturbance includes trampling on MCR steelhead redds, resulting in injury or death to incubating embryos or alevin; disturbing holding or spawning adults, forcing them to alter their behavior and seek cover; or disturbing rearing juveniles, forcing them to alter their behavior and seek cover.

Grazing can have a number of detrimental effects on riparian and aquatic habitat (Platts 1991). When riparian habitat is adversely affected, the survival and growth of listed fish species may also be negatively affected. For example, if temperatures increase to critical levels due to reduced shade, salmonid survival can decrease and some habitat may be abandoned as fish migrate to seek cooler temperatures. Loss of overhead cover in the form of overhanging vegetation or undercut banks is likely to result in increased predation of juvenile salmonids. Increases in fine sediment are likely to increase turbidity that can alter salmonid behavior, and is also likely to increase fine sediment in spawning gravels that decreases egg-to-fry survival.

However, the livestock grazing end-point indicators were developed to meet PACFISH grazing standards and guidelines, enclosure B of the LMRP and water quality BMPs. The assumption is that meeting the endpoint indicators would move key riparian and stream channel elements (bank stability, width to depth ratio, woody species regeneration) towards their Desired Conditions and meet Riparian Objectives. This will allow recovery of degraded riparian habitat to occur. Recovery of riparian vegetation results in the development of more complex habitat. Riparian recovery allows roots to stabilize streambanks, and stems and foliage to slow water velocities, trap fine sediment, provide overhead cover for fish, provide shade that may aid in keeping stream temperatures cool, and provide surfaces for macroinvertebrates to inhabit. Stable stream banks and fine sediment trapping result in less fine sediment in spawning substrate that would improve egg-to-fry survival (Tappel and Bjornn 1983, Bjornn and Reiser 1991). Reduced water velocities along stream edges increase the amount of available habitat for young salmonids (Bjornn and Reiser 1991). Spawning salmonids appear to prefer spawning in areas in close proximity of overhead cover (Bjornn and Reiser 1991), and overhead cover protects juvenile salmonids from predation. Shade provided by vegetation can be important in keeping stream temperatures cool for salmonids. Li et al. (1994) found that trout abundance decreased as solar input and water temperature increased. Macroinvertebrates inhabiting overhanging vegetation provide forage for juvenile MCR steelhead when they fall into the stream. Each of these benefits contributes to increasing the amount and quality of habitat available for all freshwater life stages of MCR Steelhead.

7.3.2 DIRECT EFFECTS TO SPECIES

The three allotments contain MCR Steelhead spawning and rearing habitat. At certain times and under various conditions it is possible for livestock use (PE 1) to directly impact listed MCR steelhead. These effects could manifest themselves as direct impacts to individual fish, fry, or incubating embryos.

Direct impacts are likely to occur if livestock waded into a stream and disturb rearing juveniles or spawning adults, and/or step on redds. Juveniles in close proximity to stream crossings or watering sites are likely to move out of an area when livestock enter or approach the stream. Juveniles are likely to be at increased risk of predation. Livestock will have access to spawning habitat in the Fields Peak allotment during the spawning period. Spawning steelhead have been observed at low densities (< 1 redd per mile) in the action area in Tex and Murderers creeks (Powell and Delano 2004). This interaction, which is not discountable or insignificant (e. g. Gregory and Gamett 2009), results in an LAA determination for direct effects to the species for the Fields Peak Allotment. For the Seneca and Hanscomb allotments, potential for cattle-fish interactions is discountable due to the difficult access in Vance Creek (Seneca Allotment, MNF 1993), lack of fish in Hanscombe Creek (Seneca and Hanscomb allotments, MNF1995a), and lack of suitable spawning habitat in Laycock Creek (Hanscomb allotment, MNF 1995d). Thus direct effects to the species are discountable (but see indirect effects to habitat below in 7.3.3 for Laycock Creek). For the Deadhorse Allotment steelhead-cattle interactions are also discountable in Ingle Creek (assumes anadromous fish do not reach the action area, MNF 1995b) and Riley Creek where steelhead are present (low likelihood of cattle accessing lower reach).

It is likely that spawning behavior could be interrupted, forcing adults to retreat to nearby cover, and that redds will be at risk of being stepped on by cattle, potentially causing direct mortality (Gregory and Gamett 2009). However, these risks will be minimized with the implementation of the *Malheur National Forest Strategy to Minimize Redd Trampling “Take” of Steelhead and Bull Trout* (Appendix G). Additionally, MNF staff (range and aquatic specialists) will take extra effort to monitor these sites when they are in the field. If active redds are located, mitigation actions will be taken to eliminate or significantly minimize the potential for redd trampling (PDC 5 and 6).

The potential for direct impacts from PE 2 (permittee management and infrastructure maintenance) is much smaller. Road use has no potential for direct impacts to the species. The PDCs do not allow off-road vehicles to cross streams except for use of existing fords on road crossings. Range riders on horses will occasionally cross streams, but redds will be identified and avoided, and any disturbance to adults or juveniles should be sufficiently brief to not result in significant disruption of normal behavioral patterns. Infrastructure maintenance actions are not located in stream channels, so there is no mechanism for direct impacts to the species.

Some monitoring activities (PE 5) involve walking in stream channels. Actions such as pebble counts and redd surveys will result in individuals walking across stream channels for time periods that may result in MCR steelhead being disturbed and moving out of the area, resulting in direct impacts to the species. Surveyors will search for steelhead redds when walking in streams and avoid if present.

7.3.3 DIRECT AND INDIRECT EFFECTS TO AQUATIC AND RIPARIAN HABITAT

Use of the NMFS MPI to determine effects to listed fish species is based upon using the effects of the action on habitat indicators as a surrogate for effects to the species. The premise is that the indicators and the range of environmental baseline conditions provided by the three classifications (PF, AR, NPF) depict the biological requirements of the listed fish species. Since there is a direct relationship between habitat condition and the growth and survival of individual fish at various life stages, the effects of the action on habitat variables can be linked to effects to individuals of the species, and ultimately to an ESA effect determination.

The analysis in the “Effects to Critical Habitat” section (Section 6.3) evaluated specific NMFS MPI indicators that correspond to the PCEs of CH. The PCEs are used to describe “those physical or biological features that are essential to the conservation of the listed species.” The same sub-set of NMFS MPI indicators evaluated for effects to PCEs also apply to the analysis of effects to the species. To eliminate redundancy, only those indicator/PE combinations for which a conclusion of effect to a component of a PCE was “negative and meaningfully measured” will be brought forward for further evaluation in this section, as they have the potential to adversely affect listed MCR Steelhead. This conclusion was only found for PE 1 (livestock use) and not for PE 2 (permittee management and infrastructure maintenance) or PE 5 (monitoring). The indicators for which “negative and meaningfully measured” effects were concluded are:

- Water Temperature
- Sediment/turbidity
- Substrate embeddedness
- Large woody debris
- Refugia

7.3.3.1 EFFECTS ON WATER TEMPERATURE

Water temperature is an important factor affecting distribution and abundance of salmonids within the action area. Water temperatures influence water chemistry, as well as every phase of salmonid life history. Optimal temperatures for steelhead are 52° to 59° F (11° to 15° C), and the lethal temperature is approximately 77° F (25° C) (NMFS 2009). Stream temperatures are of particular concern within the John Day Subbasin. This finding is highlighted in the John Day Subbasin Plan (NPCC 2005) as well as the MCR Steelhead recovery plan (NMFS 2009). Degraded water quality, which includes elevated water temperatures, is identified as a Limiting Factor in both plans.

Temperature monitoring information for MNF lands within the allotments is derived from five sites that were monitored sporadically from 1995 to 2005: Fields, Wickiup, Buck Cabin, and Tex creeks in the Fields Peak allotment and Riley Creek of the Deadhorse allotment. Fields and Tex creeks are on the Oregon Department of Environmental Quality 303(d) list for water temperature (ODEQ 2010). Analysis of the temperature monitoring information at all sites determined they were NPF under NMFS MPI criteria (see Section 6.1.3). Within the action area, high stream temperatures occur near the end of July or the beginning of August and coincide with low stream flows and warm daytime temperatures. By the end of August, stream temperatures are typically dropping.

Stream temperature is driven by the interaction of site conditions, weather, riparian vegetation, and the input of radiant energy to a stream system. Energy exchange that affects a change in water temperature may involve solar radiation, long wave radiation, evaporative heat transfer, convective heat transfer, conduction, and advection (Lee 1980; Beschta and Weathered 1984) (Figure 8). Solar radiation is the most important source of radiant energy affecting stream temperature (Brown 1969; Beschta 1997). With the exception of solar radiation that only delivers heat energy, all the other processes are capable of both introducing and removing heat from a stream. While the process of introducing and removing heat from a stream is complex, certain processes are more important than others in determining how stream temperature is affected by

solar inputs (Beschta et al. 1987). In terms of water temperature increases, the principal source of heat energy is solar radiation directly striking the stream (Brown 1972) (Figure 8.⁹)




FIGURE 8. HEAT EXCHANGE BETWEEN A STREAM AND ITS ENVIRONMENT.

Canopy density and height are the dominant factors in the ability of streamside vegetation to intercept incoming solar radiation and reduce the rate of warming. Decline in the abundance and vigor of riparian plants in a floodplain may also cause streams to become shallow and wide, which increases the surface area that is exposed to solar radiation. Platts (1981) cited studies by Claire and Storch (1977) and others that found that removal of streamside vegetation contributed to increases in water temperatures in small headwater streams as well as influencing suspended sediment concentrations. Small streams (especially Rosgen C and E channels such as Tex and Murderers creeks) are more susceptible to warming because they have a lower volume of water to absorb solar energy. They are also more susceptible to warming because grazing impacts herbaceous vegetation and shrubs that typically provide shade to the stream channel.

Effective shade is the total solar radiation blocked from reaching the stream over a twenty-four hour period, expressed as a percentage of the total solar radiation:

$$\text{Effective Shade} = \frac{\text{Total Solar Radiation} - \text{Total Solar Radiation Reaching the Stream}}{\text{Total Solar Radiation}}$$

Effective shade is provided by features such as topography and vegetation (ODEQ 2010). Effective shade is influenced by slope steepness, vegetation species composition, tree height, vegetation density, tree distance from the stream bank, and stream width. Thus, although riparian vegetation is a physical barrier between the stream and incoming solar radiation, only a portion of the riparian canopy contributes to effective shade. The relationship of variables

⁹ Stream temperature is an expression of heat energy per unit volume, which in turn is an indication of the rate of heat exchange between a stream and its environment.

influencing effective shade can be simplified, to some degree, using geometry and computer models that simulate shade (Park 1993, Boyd 1996).

Figure 9 and Table 25 illustrate the relationship between shade and stream channel width. As stream channel width increases beyond the point where vegetation is not tall enough to cast a shadow across the stream channel, shade values decrease. The model analysis results in Table 25 are based on the shadow cast by vegetation at a distance of 1 foot and farther from the edge of the channel.




FIGURE 9. SHADE PROVIDED BY 150-FOOT TALL CONIFERS (PLATTS ET AL 1987), (PARK, 1993).

TABLE 25. EFFECTIVE SHADE PROVIDED BY THREE HEIGHTS OF GREENLINE VEGETATION AT VARYING ACTIVE STREAM CHANNEL WIDTHS.¹

Active Channel Width (feet)	Percent Effective Shade at Varying Vegetation Heights		
	0.5 feet	2 feet	3 feet
1	0	46	57
3	0	22	41
7	0	9	18
10	0	7	12
12	0	5	10
14	0	5	9
16	0	4	8
18	0	4	7

20	0	3	6
22	0	0	3
24	0	0	1
26	0	0	0

¹Effective shade values are based on the shadow cast by continuous vegetation growing at a distance of 1 foot and farther from the edge of the channel

Figure 9, Table 25 and the discussion above illustrate that decreasing vegetation height will reduce effective shade, and as a result increase solar radiation to the surface of the water in the stream. If the potential height of ungrazed vegetation is in the 2-3 feet range, then a considerable loss of effective shade takes place when the vegetation is grazed to an end-point of 0.5 feet (six inches) or less. The potential for stream warming is greatly increased, particularly for streams with narrow active channel widths.

Ungrazed height for 13 grass or grass-like plant species in the MNF ranged from 14 to 36 inches, with a mean of 26 inches (MNF 2007). The vegetation heights in Table 25 of two and three feet encompass the approximate mean ungrazed height and maximum ungrazed height of the 13 species. An end-point of 6 inch (0.5 feet) stubble height is used when livestock are grazed, which also reflects use by wild ungulates. In meadow streams with narrow channels, grass/grass-like species often are the plants that provide stream shade. Model results presented in Table 25 indicate that managing to a six-inch vegetation height will reduce effective shade to zero for channel widths that are 1 foot and greater when the model criterion is for vegetation beginning at 1 foot distance from the active channel, and to no more than 18 percent if modeled for vegetation at the edge of the active channel. This is considerably less than effective shade provided by potential vegetation heights of two to three feet for similar active channel widths.

A logical conclusion is that implementation of PE 1 (livestock use) will reduce effective shade for a bank distance sufficient to result in a measurable water temperature increase. These impacts are expected to be generally confined to narrow stream channels with grass/grass-like vegetation providing shade. In the action area, this would include portions of Tex, Murderers, Basin, Charlie Mack, and White creeks. Murderers Creek temperature data gathered in the Murderers Creek pasture in the Fields Peak allotment during the late 1990s showed summer mean maximum temperature of about 60 °F, documenting this section of stream provides important coldwater summer rearing habitat for MCR steelhead (MNF unpub data. Murderers Creek above White Creek).

Platts (1991) states that grasses are too short to keep much solar radiation from reaching the water, except along very small streams (stream orders 1 and 2). Wright and Li (2002) measured wetted widths in late July and early August 1996-1998 for five 1st and 2nd order streams in the MFJD River drainage. These are likely to have similar dimensions to streams of the same orders in the UMJD River drainage. The mean wetted width was 2.0 meters with a standard error of 0.5 meter (6.6 feet with a standard error of 1.6 feet). There is 90% confidence that the true mean wetted width is within 6.6 +/- 3.4 feet (3.2 to 10.0 feet). The greatest probability of measurable temperature increases as a result of livestock grazing is therefore likely to occur in channels less

than 10 feet wide. Portions of Murderers and Tex creeks in the action area have summer wetted widths less than 10 feet wide even though these streams are at least 3rd order.

Figure 9 and Table 25 also indicate that effective shade is reduced as channel width increases. Grazing by large hooved animals has the potential to increase channel width by bank alteration (Armour et al. 1991; Clary and Webster 1989; Kaufmann and Kruger 1984). However, the analysis presented for effects to width-depth ratio in Table 23 for the Floodplain Connectivity habitat feature of PCE 2 (rearing critical habitat) concluded that the livestock use PE is anticipated to have a negative, but not meaningfully measured effect to the indicator. The potential for increases in width-depth ratio is mitigated because of implementation of endpoint indicators for livestock grazing (which includes all use by wild ungulates) and adaptive management. For example, McDowell and Mowry (2002) found little difference in width to depth ratios between grazed areas and exclosures along Tex Creek, although they did find a lower width to depth ratio in an exclosed section of Murderers Creek when compared to a grazed section. However, due to endpoint indicators and adaptive management, effective shade will not be measurably reduced as a result of increasing the width-to-depth ratio indicator.

Conclusion. The discussion above described negative effects to habitat and vegetation characteristics from the livestock use PE. Effects to these characteristics result in negative impacts (increases) to summer water temperature. It is probable that livestock use will result in small, but measurable increases in water temperature in streams with narrow channels (<10 feet) where grass/grass-like vegetation is providing stream shade such as portions of Tex, Murderers, and possibly Laycock creeks (Fields Peak and Hanscomb allotments). This will occur as a direct result of reducing the height of shade-producing vegetation such as grasses and sedges by grazing.

These effects to water temperature will be minimized by use of endpoint indicators and PDC. These indicators were developed to meet PACFISH grazing standards and guidelines as well as water quality BMPs. The assumption is that meeting these endpoint indicators would move key riparian and stream channel elements (bank stability, w/d ratio, woody species regeneration) towards their Desired Conditions and meet Riparian Objectives. If monitoring fails to show this upward trend, adaptive management and administrative actions would be implemented to continue to minimize adverse effects to CH and the listed MCR steelhead. Some impacts from past management activities (e.g. logging, roads, grazing) will persist over the life of this consultation and likely much longer in some cases.

7.3.3.2 EFFECTS ON SEDIMENT/TURBIDITY, SUBSTRATE AND SUBSTRATE EMBEDDEDNESS

Grazing by large herbivores such as domestic cattle can result in hoof shear to streambanks, and trampling and consumption of streamside vegetation. The result is a potential increase in the supply of fine sediment available for transport. This can occur when grazing results in compacted soils and bare areas; and when grazing results in decreased bank stability through mechanical damage to streambanks or reductions in rooting strength of streambank stabilizing vegetation. Both result in an increase in erosion rates and subsequent increases in fine sediment levels in streams.

Small amounts of fine sediment are also likely to enter streams where livestock access streams to cross or water. Small amounts of fine sediment are likely to become deposited in substrate that

can decrease egg-to-fry survival and slightly reduce available substrate cover for juveniles and macroinvertebrates.

Increased fine sediment is detrimental to MCR steelhead through increased turbidity and sediment deposition in the substrate (NMFS 2009). Increases in fine sediment lead to greater substrate embeddedness and a decrease in the interstitial spaces between gravel substrate important for salmonid spawning. Successful salmonid spawning requires clean gravels with low fine sediment content (Spence et al. 1996). Well-oxygenated water must be able to reach eggs and pre-emergent fry during incubation and emergence. Suffocation of these life stages may occur if redds become covered with fine sediment. Emerging fry may be physically blocked from escaping a redd. Increased sediment load is also detrimental to juvenile salmon by introducing suspended particulate matter that interferes with feeding and territorial behavior (Berg and Northcote 1985). Increased fine sediment deposition in the substrate is likely to decrease MCR steelhead egg-to-fry survival (Reiser and White 1988, Spence et al. 1996).

In addition, inputs of fine sediment resulting from livestock trampling banks can reduce benthic invertebrate abundance and lead to a shift from aquatic insects to mollusks, which are less palatable to salmonids. Studies have shown that sediment inputs resulting in substrate embeddedness of greater than one-third can result in a decrease in benthic invertebrate abundance and thus a decrease in food available for juvenile salmonids (e.g. Waters 1995). Other field studies have shown small amounts of fine sediment and corresponding embeddedness have measurable effects on juvenile steelhead growth, behavior, and survival (Suttle et al. 2004).

There are no streams in the proposed action area that have been identified on the 303(d) list for sedimentation. Fine sediment levels vary across the allotments depending on local geology, stream type, and management history. Currently the PIBO information within the allotments show percent fines < 2mm in the pool tail-outs as ranging from 6.2 to 23.8 percent (sample size of 2). Fine sediments have been reduced in both sites between 2003 and 2008 (9.3 % vs 23.8 in Riley Creek and 6.2% vs 23.6 in Fields Creek, Tables 18 and 19) suggesting improving management practices. Amendment 29 of the LMRP sets a desired condition of <20% embedded. The most current level II stream survey data available reported that the majority of reaches within the allotments (19 of 26 stream reaches) have substrate embeddedness greater than 20% (See Appendix J for PIBO and Stream Survey results).

Conclusion. The livestock use PE will likely result in fine sediment entering stream channels where MCR steelhead are present in areas such as, but not limited to, Tex and Murderers creeks in the Fields Peak allotment. It will likely also result in sediment entering areas where steelhead may not currently be present but its CH is designated such as Laycock Creek in the Hanscomb Allotment (MNF 1995d). It is unlikely the livestock use PE would introduce meaningfully measurable sediment into Vance Creek in the Seneca Allotment due to its heavy brush and tree-lined riparian area or Ingle and Riley creeks in the Deadhorse allotment. The probability is not discountable that sediment would be meaningfully identifiable from the livestock use PE in the Fields Peak and Hanscomb allotments. The mechanisms include: 1) mechanical bank damage from hoof chisel and trampling; 2) trailing; and, 3) impacts to soil-holding vegetation by being eaten and trampled. These mechanisms negatively impact bank stability, resulting in increased erosion and sedimentation. The increases in fine sediment will negatively and measurably affect the Sediment/Turbidity and Substrate Embeddedness NMFS MPI indicators. This analysis understands that wild ungulate use may also significantly and negatively affect the sediment

indicator in some of these streams and some of the effects from cattle use observed in the early and mid-1990s (particularly unauthorized use in Ingle Creek, MNF 1995c) in the stream survey reports are unlikely under the current proposed action.

These effects to the Sediment/Turbidity and Substrate Embeddedness indicators will be minimized by use of endpoint indicators and PDC. If pre-season monitoring indicates that wild ungulate use is resulting in measurements near or exceeding an endpoint indicator, cattle will not be turned-out into that specific pasture. These indicators were developed to meet PACFISH grazing standards and guidelines as well as water quality BMPs. The assumption is that meeting these endpoint indicators would move key riparian and stream channel elements (bank stability, w/d ratio, woody species regeneration) towards their Desired Conditions and meet Riparian Objectives. If monitoring fails to show this upward trend, adaptive management and administrative actions would be implemented to continue to minimize adverse effects to designated CH and the listed MCR steelhead. Some impacts from past management activities (e.g. logging, roads, grazing) will persist over the life of this consultation and likely much longer in some cases.

7.3.3.3 EFFECTS ON REFUGIA

The concept of “Refugia” is not described in detail in the NMFS MPI (NMFS 1996) (see Table 14 earlier in this document). The definition provided therein is: “important remnant habitat for sensitive aquatic species.” The availability of various types of habitat refugia are described as limiting factors in the recovery plan for the Oregon steelhead populations of the MCR Steelhead DPS (e.g., loss of side-channels that provided high flow refugia; cold water refugia provided by Columbia River tributary streams such as the Deschutes River (NMFS 2009). In the action area, MSRAs may qualify as refugia because these areas have some of the highest steelhead habitat potential due to their low gradient and other geomorphic qualities.

The analysis of effects to PCEs of CH provided in Table 23, and summarized in Table 24, indicate that the livestock use PE will have negative and meaningfully measured effects to several of the MPI indicators that correlate to components of PCEs. Specifically, they are “Water Temperature,” “Sediment/Turbidity,” and “Substrate Embeddedness.” This may occur in stream reaches providing refugia conditions for one or more of these habitat characteristics (areas with cooler water temperatures, low levels of sediment in substrate or the water column, and low levels of substrate embeddedness). Therefore, PE 1 will have a negative effect to the refugia indicator.

Conclusion. The livestock use PE will result in negative and meaningfully measured impacts to several habitat indicators associated with refugia in the Hanscomb and Fields Peak allotments. Consequently, there will be negative and meaningfully measured, evaluated or detected impacts to the refugia indicator. The effects are not expected to be distributed evenly across the ESA action area because stream reaches providing characteristics of refugia are not ubiquitous. The effects will likely be observed in areas where adequate cattle forage overlaps MSRAs or other refugia such as seeps or springs used by steelhead. Negative impacts to the refugia indicator will be minimized by use of the endpoint indicators and PDC.

7.3.3.4 EFFECTS ON LARGE WOODY DEBRIS

Large woody debris (aka large wood) is one of the most important habitat components in many fish-bearing streams (Gurnell et al. 2002). Large wood helps provide cover, scour pools, stabilize banks, retain spawning gravels, create off-channel habitats, and provide habitat for macroinvertebrate production (Gregory et al. 2003).

In streams within the action area, large wood is usually provided by fallen conifers that have no effect from the project elements. However, in some areas where hardwoods—particularly black cottonwood and quaking aspen—play an important role in riparian species composition, ungulate grazing can prevent future large wood recruitment by limiting sapling regeneration and large tree recruitment. Young cottonwoods are desirable forage to both domestic and wild ungulates (Braatne et al. 1996).

Kaufmann et al. (1983) found late season riparian cattle grazing retarded regeneration of black cottonwood saplings in northeastern Oregon. Another study found when cattle were removed from a riparian pastures, but wild ungulates were not excluded, the number of black cottonwood seedlings/saplings increased 56% 3 years after livestock removal (Case and Kaufmann 1997). Clearly, cattle grazing can influence the abundance of black cottonwoods in a riparian area, which can have measurable and foreseeable future effects to riparian structure and future large wood recruitment. Beschta and Ripple (2005) surveyed a 40-mile reach of the Middle Fork John Day River nearby the action area for cottonwood abundance and stand structure and found very little cottonwood seedling/sapling regeneration or recruitment into large trees and described wild and domestic ungulate browsing as the primary causal factor.

The analysis of effects to PCEs of CH provided in Table 23, and summarized in Table 24, indicate that the livestock use PE will have negative and meaningfully measured effects to the “Large Woody Debris” MPI indicator that correlates to components of PCEs. Therefore, PE 1 will have a negative effect to the large woody debris indicator.

Conclusion. The livestock use PE will likely result in negative effects to future large woody debris recruitment. The effects will likely be observed in areas where adequate cattle forage overlaps low-gradient stream sections such as MSRAs that have relatively open canopy and have potential to develop a cottonwood gallery forest. The mechanisms include: 1) browsing on young cottonwoods seedlings/saplings, 2) retarding cottonwood succession and large tree recruitment; and, 3) reduction in future levels of instream large wood. These mechanisms will negatively and measurably affect the large woody debris NMFS MPI indicators in the Hanscomb and Fields Peak Allotments. Negative impacts to the large woody debris indicator will be minimized by use of the endpoint indicators and PDC.

8 ESA EFFECT DETERMINATIONS

ESA effect determinations are presented in Table 2. The determinations are “May Affect, Likely to Adversely Affect” for MCR Steelhead and its designated CH in the Hanscomb and Fields Peak Allotments. For the Seneca and Deadhorse allotments, the determinations are “May Affect, Not Likely to Adversely Affect” for MCR Steelhead and its designated CH. There are no bull trout or its designated critical habitat in the action area. Consequently, the effect determinations across all allotments for bull trout are “No Effect.”

8.1 RATIONALE

The PCEs are the physical or biological features of critical habitat essential to the conservation of the species. For PCE 1 (Freshwater spawning sites), the analysis determined that there were **negative and measurable effects** to the *temperature* and *sediment* indicators corresponding to the *water quality* feature of the PCE, and the *substrate embeddedness* indicator corresponding to the *suitable substrate* feature of the PCE, as diagrammed below:

- PCE1: Freshwater spawning sites.
 - Water quality PCE feature
 - Temperature indicator
 - Sediment indicator
 - Suitable substrate PCE feature
 - Substrate embeddedness indicator

In addition, for PCE 2 (Freshwater rearing sites), the analysis determined that there were **negative and measurable effects** to the *temperature* and *sediment* indicators corresponding to the *water quality* feature of the PCE, the *substrate embeddedness* indicator corresponding to the *forage* feature of the PCE, and the *refugia* and *large woody debris* indicator for both the *forage* and *natural cover* features of the PCE, as diagrammed below:

- PCE2: Freshwater rearing sites.
 - Water quality PCE feature
 - Temperature indicator
 - Sediment indicator
 - Forage PCE feature
 - Substrate embeddedness indicator
 - Large woody debris indicator
 - Refugia indicator
 - Natural cover PCE feature
 - Large woody debris indicator
 - Refugia indicator

Negative measurable effects do not meet the definition of “insignificant” effects and they are not “discountable” because the effects are likely to occur. Consequently, the effect determination for MCR Steelhead designated CH overall for the Hanscomb and Fields Peak allotments is “May Affect, Likely to Adversely Affect.”

The same NMFS MPI indicators determined to have negative, measurable effects during the PCE analysis were brought forward in the analysis of effects to the species. The mechanisms by which the livestock use (PE 1) would affect habitat characteristics that would in turn result in measurable increases in water temperature, increased sediment and turbidity, increased substrate embeddedness, and decreased large woody debris were described in detail. The biological consequences to MCRS Steelhead were also described. The conclusion was that the effects to the indicators would result in negative effects to each indicator that were measurable, and therefore did not meet the definition of “insignificant” effects. They are not “discountable” because the effects are likely to occur.

In addition, there are likely to be direct effects to individual MCR steelhead from the livestock use PE, because livestock will have access to streams during the spawning, incubation, and rearing periods of the MCR steelhead life cycle. Direct impacts are likely to occur if livestock wade into a stream and disturb rearing juveniles or spawning adults, and/or step on redds.

Juveniles in close proximity to stream crossings or watering sites are likely to move out of an area when livestock enter or approach the stream. Juveniles forced into open water could have increased risk of predation. It is likely that spawning behavior will be interrupted forcing adults to retreat to nearby cover and that redds will be at risk of being stepped on. However, these risks will be minimized with the implementation of the *Malheur National Forest Strategy to Minimize Redd Trampling “Take” of Steelhead and Bull Trout* (Appendix G), which incorporates PDC 5 and 6 (Section 4.1.2). Additionally, MNF staff (range and aquatic specialists) will take extra effort to monitor these sites when they are in the field. If active redds are located, mitigation actions consistent with PDC 6 (Section 4.1.2) will be taken to eliminate or significantly minimize the potential for redd trampling,

In summary, because the proposed action will result in measurable negative effects to components of MCR steelhead habitat, with indirect effects to the species, and it is likely there will be direct negative effects to adults, juveniles and possibly incubating embryos in redds, the effect determination is “May Affect, Likely to Adversely Affect” for the species.

9 ESA CUMULATIVE EFFECTS

ESA cumulative effects are those effects of future State, tribal, local or private activities that are reasonably certain to occur in the area of the Federal action subject to consultation. Future Federal actions that are unrelated to the proposed action are not considered in this section because they are subject to separate consultation pursuant to section 7 of the ESA. There are several future State or private activities that are reasonably certain to occur.

9.1 ODFW STATE LANDS

Land managed by the state of Oregon is located adjacent to the Fields Peak Allotment, including stream frontage along Murderers Creek and Tex Creek near their confluence. Livestock grazing may continue on these state lands at similar levels to recent years. The effects of livestock grazing on state lands are similar to those described in the effects to MCR steelhead CH section by the proposed action: (1) increased sediment in stream channels resulting in increased turbidity, substrate embeddedness, a reduction in macroinvertebrate production, and reduced quality of spawning gravel; (2) increase in water temperature as a result of shade loss along stream channels from grazing/browsing of riparian vegetation; (3) potential loss of future large woody debris.

9.2 ODFW ELK AND DEER MANAGEMENT

The ODFW manages Rocky Mountain elk and mule deer populations in the action area. The action area is located entirely within the state of Oregon’s Murderers Creek Wildlife Management Unit (WMU).

Elk and mule deer utilize streamside vegetation differently. Both animals eat riparian vegetation, but have different forage preferences. The diets of elk, mule deer, and cattle are very different during early summer and become increasingly similar during late summer. Cattle diets have more grasses, deer diets have more shrubs and forbs, and elk diets are in between those of cattle and deer. (USDA 2006). There is overlap between what each species will eat dependent upon season and availability. Additionally, Coe et al. (2005) found a cascading effect of larger

ungulates displacing smaller ungulates. They found that the presence of livestock displaced smaller ungulates including mule deer and elk, and that livestock chose resources such as forage before smaller ungulates.

Table 26 presents Rocky Mountain elk and mule deer management objectives (MO) and population estimates from 2004-2010 for the Murderers Creek WMU that entirely encompasses the Murderers Creek Allotment. The mule deer population MO was obtained from ODFW (2003), available online at:

http://www.dfw.state.or.us/wildlife/management_plans/docs/MuleDeerPlanFinal.PDF. Mule deer population estimates, and Rocky Mountain elk MOs and population estimates, were obtained from ODFW wildlife biologist Ryan Torland (pers. comm. 2011).

TABLE 26. ROCKY MOUNTAIN ELK AND MULE DEER MANAGEMENT OBJECTIVES AND WINTER POPULATION ESTIMATES FROM 2004-2010 FOR THE MURDERERS CREEK WILDLIFE MANAGEMENT UNITS IN OREGON.

Year	Murderers Creek Wildlife Management Unit	
	Elk Mgmt. Objective = 1,700	Deer Mgmt. Objective = 9,000
2004	1,700	6,695
2005	1,800	6,968
2006	1,800	6,820
2007	2,273	5,207
2008	1,900	6,820
2009	1,900	6,968
2010	1,900	6,695

ODFW has managed the elk population of the Murderers Creek WMU at or above the population MO. Since 2005, the elk population has exceeded the MO for six consecutive years. The mule deer population MO was not exceeded during 2004-2010 in the WMU. It is not known what elk or mule deer population level in the WMU would result in detectable effects to the PCEs of CH for MCR Steelhead.

There are data available regarding combined wild ungulate use (i.e., deer and elk) in specific pastures nearby the allotments within the action area that were not grazed in 2009 and 2010. The MNF proposed grazing use endpoints were approached in a few instances. In 2010 in the Belshaw Riparian Pasture of the Mt. Vernon Allotment, the percent woody browse use was estimated at 37.6 percent when the endpoint is 40 percent. Similarly in 2010 for the Lower Creek Pasture of the John Day Allotment, the percent woody browse use was estimated at 38.6

percent. Percent streambank alteration was estimated at 15 percent at the Bear Creek Pasture in 2010, which was five percentage points below the proposed MNF endpoint of 20 percent. Had livestock been turned out in these allotments, their use would have been limited, otherwise the endpoints might have been exceeded.

Because these data were from areas nearby the Seneca, Deadhorse, Hanscomb, and Fields Peak allotments, it is likely wild ungulate use also occurs in these allotments. Indeed, monitoring observations from late August 2011 showed bank trampling and greenline vegetation grazing from wild ungulates (mule deer, elk, feral horse) in MSRAs along Murderers and Tex creeks within the action area that were rested from cattle grazing (Reid personal observation 2011).

There is a potential for cumulative effects to MCR Steelhead designated CH from use by wild ungulates. Such effects are identical to those described in the effects to MCR Steelhead CH section: (1) increased sediment in stream channels resulting in increased turbidity, substrate embeddedness, a reduction in macroinvertebrate production, and reduced quality of spawning gravel; (2) and an increase in water temperature as a result of shade loss along stream channels from grazing/browsing of riparian vegetation.

9.3 UNAUTHORIZED LIVESTOCK GRAZING

Unauthorized livestock grazing has occurred in the allotments, and is reasonably certain to occur in the future. However, it is not a large problem in the Murderers Creek Allotment. Typically, it involves a few cow/calf pairs every few years. The unauthorized use has been resolved in a timely manner once MNF staff notified livestock owners. As long as the MNF takes timely action whenever trespass occurs, habitat degradation is likely to be minimized.

9.4 ACTIONS ON PRIVATE PROPERTY

The ESA action area includes no private property but there is one private in-holding upstream of MNF land in the Fields Peak allotment. There is the potential for this property to be developed. However, there is no information on specific proposals at this time. The effects to PCEs of CH (if any) of activities on private property, such as cattle grazing, are expected to continue at the same rate as they have been.

10 ESSENTIAL FISH HABITAT FOR CHINOOK SALMON

The Magnuson-Stevens Act, as amended by the Sustainable Fisheries Act of 1996 (Public Law 104-267), established procedures designed to identify, conserve, and enhance Essential Fish Habitat (EFH) for those species regulated under a Federal fisheries management plan. The MSA requires Federal agencies to consult with NOAA Fisheries on all actions, or proposed actions, authorized, funded, or undertaken by the agency, that may adversely affect EFH.

The Pacific Fisheries Management Council (PFMC) is one of eight regional fishery management councils established under the Magnuson-Stevens Act. PFMC develops and carries out fisheries management plans for salmon, groundfish and coastal pelagic species off the coasts of Washington, Oregon, and California, and recommends Pacific halibut harvest regulations to the International Pacific Halibut Commission.

As required by the Magnuson-Stevens Act, the PFMC described and identified Essential Fish Habitat (EFH) in each of its fisheries management plans. The EFH includes “those waters and

substrates necessary to fish for spawning, breeding, feeding, or growth to maturity.” All streams, lakes, ponds, wetlands, and other water bodies currently, or historically accessible to salmon in Washington, Oregon, Idaho, and California are designated as EFH for affected salmon stocks with management plans. The Upper John Day 4th field HUC (HUC 17070201), which encompasses the project area, has been designated as EFH for Chinook salmon (73 FR 200:60987 October 15, 2008). However, finer resolution of what constitutes waters “currently or historically accessible to salmon” is dependent upon local information.

The Magnuson-Stevens Act (MSA) also established an EFH consultation process. Federal agencies are required to consult with NMFS on all actions that may adversely affect EFH. The NMFS interprets the scope of these consultations to include actions by Federal agencies that occur outside designated EFH, such as upstream or upslope, but which nonetheless may have an adverse effect on habitat conditions necessary for the long-term survival of the species within EFH. NMFS must provide conservation recommendations for any Federal or State activity that may adversely affect EFH. Within 30 days of receiving EFH conservation recommendations from NMFS, Federal agencies must conclude EFH consultation by responding to NMFS with a written description of conservation measures the agency will use to avoid, mitigate or offset the impact of its action on EFH. If the Federal agency selects conservation measures, which are inconsistent with the conservation recommendations of NMFS, the Federal agency must explain in writing its reasons for not following NMFS recommendations.

The MNF searched for information to determine if the action under EFH consultation includes areas currently or historically accessible to Spring Chinook salmon. The MNF could not find any references that Chinook salmon historically utilized any streams within the allotments. Regarding current use, an ODFW website provides access to maps titled *Spring Chinook Habitat: Seneca Quad*, *Spring Chinook Habitat: Aldrich Gulch Quad*, *Spring Chinook Habitat: Canyon Mountain Quad*, and *Spring Chinook Habitat: John Day Quad*. The Chinook salmon distribution maps depict use of the UMJD River, SFJD River, and Canyon Creek in the vicinity of the three allotments. No spring Chinook use is predicted for any of the streams that drain the action area, including larger streams such as Murderers Creek. The ODFW maps are accessible at: http://nrimp.dfw.state.or.us/nrimp/default.aspx?pn=chs_dist.

The analysis of effects to designated CH for MCR Steelhead concluded that there were measurable negative effects to several PCEs. These effects would be a proxy for effects to Chinook salmon EFH as they have similar habitat requirements. However, there is no information to support that Chinook salmon historically or currently use any stream in the five allotments. Furthermore, the nearest Chinook salmon habitat is about 1 mile downstream from the MNF boundary of the five allotments (Canyon Creek downstream of Vance Creek in the Seneca Allotment). There is no likelihood that any effects (NLAA in Vance Creek) to designated CH for MCR steelhead (as a proxy for effects to EFH for spring Chinook salmon) will be detectable at those distances in Canyon Creek, SFJD River, or the UMJD River, where historic and current use by Chinook salmon is documented. Consequently, the MNF concludes that the proposed action will not adversely affect EFH for MSA-managed Chinook salmon (Table 2).

11 REFERENCES

- Al-Chokhachy, R., B. B. Roper, and E. Archer. 2010. Using a multimetric approach to evaluate the abiotic condition of streams in the upper Columbia and Missouri river basins. *Transactions of the American Fisheries Society* 139:1041–1059.
- Alvarado, R. 2011. Personal communication. Wildlife Program Manager, Pacific Northwest Region, Regional Office, USDA Forest Service, Portland, OR. June 7, 2011.
- Armour, C.L., D.A. Duff and W. Elmore. 1991. The effects of livestock grazing on riparian and stream ecosystems. *Fisheries* 16(1): 7–11.
- Benda, L. E., D. Miller, T. Dunne, J. Agee, and G. H. Reeves. 1998. Dynamic landscape systems. Pages 261-288 in R. J. Naiman and R. E. Bilby eds. *River ecology and management: lessons from the Pacific Coastal Region*. Springer Verlag, New York.
- Bengeyfield, P. 2006. Managing cows with streams in mind. *Rangelands* 28(1): 3–6.
- Bengeyfield, P. and D. Svoboda. 1989. Determining allowable use levels for livestock movement in riparian areas. Specialty Conference on Rangeland Management and Water Resources. Proceedings. American Water Resources Association. Reno, NV.
- Berg, L. and T.G. Northcote. 1985. Changes in territorial, gill-flaring, and feeding behavior in juvenile coho salmon (*Oncorhynchus kisutch*) following short-term pulses of suspended sediment. *Canadian Journal of Fisheries and Aquatic Sciences* 42:1410-1417.
- Beschta, R.L. 1997. Riparian shade and stream temperature: an alternative perspective. *Rangelands*. 19(2): 25-28.
- Beschta, R.L. and J. Weatherred. 1984. A computer model for predicting stream temperatures resulting from the management of streamside vegetation. USDA Forest Service. WSDG-AD-00009.
- Beschta, R.L., R.E. Bilby, G.W. Brown, L.B. Holtby, and T.D. Hofstra. 1987. Stream temperature and aquatic habitat: Fisheries and forestry interaction. Pp. 191-232. University of Washington, Institute of Forest Resources, Contribution No. 57.
- Beschta, R.L., and W. J. Ripple. 2005. Rapid assessment of riparian cottonwoods: Middle Fork John Day River, northeastern Oregon. *Ecological Restoration* 23: 150-156.
- Bjornn, T.C., and D.W. Reiser. 1991. Habitat requirements of salmonids in streams. Pages 83-138, In W.R. Meehan (editor) *Influences of forest and rangeland management on salmonid fishes and their habitats*. Special Publication 19. American Fisheries Society.
- Boyd, M.S. 1996. Heat Source: stream temperature prediction. Master's Thesis. Department of Civil and Bioresource Engineering, Oregon State University, Corvallis, Oregon.
- Braatne, J.H., S.B. Rood and P.E. Heilman. 1996. Life history, ecology, and conservation of riparian cottonwoods in North America. Pages 57-85 in R.F. Stettler, H.D. Bradshaw, Jr. and T.M. Hinkley (eds.), *Biology of Populus and its Implications for Management and Conservation*. Ottawa, Ontario, Canada: National Research Council.
- Brown, G.W. 1969. Predicting temperatures of small streams. *Water Resour. Res.* 5(1):68-75.

- Brown, G.W. 1972. An improved temperature model for small streams. Water Resour. Report 16, Oregon State University, Corvallis, Oregon.
- Bunte, K. and S.R. Abt. 2001. Sampling surface and sub-surface particle sizedistributions in wadeable gravel- and cobble-bed streams for analysis in sediment transport, hydraulics, and streambed Monitoring. USDA Forest Service, Rocky Mountain Experiment Station, General Technical Report, RMRS-GTR-74. 450 pp.
- Bureau of Land Management. 1996. Utilization Studies and Residual Measurements. Interagency Technical Reference. BLM/RS/ST-96/004+1730.
- Bureau of Land Management, Prineville District (PD BLM). 2006. Biological assessment: LAA grazing actions on the lower John Day River for 2006 and beyond. Prineville BLM District Office, Prineville, Oregon.
- Burton, T.A., E.R. Cowley, and S.J. Smith. 2008. Monitoring Stream Channels and Riparian Vegetation—Multiple Indicators Version 5.0 – 2008 BLM/ID/GI-08/001+1150
- Burton, T.A., S.J. Smith, and E.R. Cowley. 2011. Riparian area management: Multiple indicator monitoring (MIM) of stream channels and streamside vegetation. Technical Reference 1737-23. BLM/OC/ST-10/003+1737. U.S. Department of the Interior, Bureau of Land Management, National Operations Center, Denver, CO. 155 pp.
- Case and Kauffman. 1997. Wild ungulate influences on the recovery of willows, black cottonwood and thin-leaf alder following cessation of cattle grazing in northeastern Oregon. Northwest Science. 1997; 71(2): 115-126.
- Claire, E. W. and R. L. Storch. 1977. Streamside management and livestock grazing: an objective look at the situation. In: *Proc. Symp. Livestock and Wildlife-Fisheries Relationships in the Great Basin*. Sparks, Nevada. May 3-5, 1977.
- Clary, W. P. 1999. Stream channel and vegetation responses to late spring cattle grazing. *Journal of Range Management*. 52: 218-227.
- Clary, W. P. and B. F. Webster. 1989. Managing grazing of riparian areas in the Intermountain Region. General Technical Report INT-263, U.S. Dept. of Agriculture, USFS, Intermountain Research Station, Ogden, Utah. 11 p.
- Clary, W. P., C. I. Thornton and S. R. Abt. 1996. Riparian stubble height and recovery of degraded streambanks. *Rangelands*. 18: 137-140.
- Clary, W.P., and W.C. Leininger. 2000. Stubble height as a tool for management of riparian areas. *J. Range Management* 53(6): 562-573.
- Coe, P. K., B. K. Johnson, K. M. Stewart, and J. G. Kie. 2005. Spatial and Temporal Interactions of Elk, Mule Deer, and Cattle. In: *Transactions of the 69th North American Wildlife and Natural Resources Conference*: 656-669.
- Coulloudon, B., K. Eshelman, J. Gianola, N. Habich, L. Hughes, C. Johnson, M. Pellant, P. Podborny, A. Rasmussen, B. Robles, P. Shaver, J. Spehar, J. Willoughby. 1999. Sampling Vegetation Attributes. BLM Technical Reference 1734-4, Denver, CO.

- Cowley, E.R. 2002. Guidelines for Establishing Allowable Levels of Streambank Alteration. USDI, Bureau of Land Management, Idaho State Office. Information Bulletin No. ID-2002-172. Boise, Idaho.
- Cowley, E.R. and T.A. Burton. 2005. Monitoring Streambanks and Riparian Vegetation – Multiple Indicators. Tech. Bull. No. 2005-002. USDI, BLM, Idaho State Office. Boise, ID. http://www.id.blm.gov/techbul/05_02/doc.pdf
- Gregory, J. S., and B. L. Gamett. 2009. Cattle trampling of simulated bull trout redds. *North American Journal of Fisheries Management* 29:361–366.
- Gregory, S.G., K. L. Boyer, and A.M. Gurnell, editors. 2003. The ecology and management of wood in world rivers. American Fisheries Society Publication. 444 pp.
- Gurnell, A., H. Piegay, F. J. Swanson, and S. V. Gregory. 2002. Large wood and fluvial processes. *Freshwater Biology* 47:601-619
- Hall, F.C and L. Bryant L. 1995. Herbaceous Stubble Height as a Warning of Impending Cattle Grazing Damage to Riparian Areas. USDA Forest Service Gen Tech Rep PNW-362. 10 p.
- Heitke, J. D., R. C. Henderson, B. B. Roper, and E. K. Archer. 2008. Evaluating livestock grazing use with streambank alteration protocols; challenges and solutions. *Rangeland Management and Ecology* 61:647–655.
- Henderson, R. C., E. K. Archer, B. A. Bouwes, M. S. Coles-Ritchie, and J. L. Kershner. 2005. PACFISH/INFISH Biological Opinion (PIBO): Effectiveness Monitoring Program seven-year status report 1998 through 2004. General Technical Report RMRS-GTR-162. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station Fort Collins, Colorado.
- Interior Columbia Basin Technical Recovery Team. 2003. Independent Populations of Chinook, Steelhead, and Sockeye for Listed Evolutionarily Significant Units Within the Interior Columbia River Domain. Working Draft, July 2003. Northwest Fisheries Science Center, NMFS Northwest Region.
- Kauffman, J. B. and W. C. Krueger. 1984. Livestock impacts on riparian ecosystems and streamside management implications - a review. *Journal of Range Management* 37(5):430-438.
- Kauffman, J. B., W. C. Krueger, and M. Vavra. 1983. Effects of late season grazing on riparian communities. *Journal of Range Management* 36(6):685-691.
- Kershner, J.L., B.B. Roper, N. Bouwes, R. Henderson, and E. Archer. 2004. An Analysis of Stream Habitat Conditions in Reference and Managed Watersheds on Some Federal Lands within the Columbia River Basin. *North American Journal of Fisheries Management* 24: 1363-1375.
- Kershner, J. and B. Roper. 2010. An evaluation of management objectives used to assess stream habitat conditions on Federal lands within the Interior Columbia Basin. *Fisheries* 35(6):269-278.

- Lee, R. 1980. Forest hydrology. Columbia University Press, New York.
- Li, H.W. G.A. Lamberti, T.N. Pearsons, C.K. Tait, and J.L. Li. 1994. Cumulative effects of riparian disturbances along high desert trout streams of the John Day Basin, Oregon. Transactions of the American Fisheries Society. 123:629-640.
- Lisle, T.E. 1987. Using "Residual Depths" to monitor pool depths independently of discharge. Research Note PSW-394. U.S. Department of Agriculture, Pacific Southwest Forest and Range Experiment Station. 4 pp.
- Malheur National Forest (MNF). 1990. Malheur National Forest Land and Resource Management Plan.
- Malheur National Forest (MNF). 1992. Fields Creek level II stream survey report.
- Malheur National Forest (MNF). 1993 Vance and South Fork Vance Creek level II stream survey report.
- Malheur National Forest (MNF). 1995a. Hanscombe Creek level II stream survey report.
- Malheur National Forest (MNF). 1995b. Riley Creek level II stream survey report.
- Malheur National Forest (MNF). 1995c. Ingle Creek level II stream survey report.
- Malheur National Forest (MNF). 1995d. Laycock Creek level II stream survey report.
- Malheur National Forest (MNF). 2004. Malheur National Forest Roads Analysis Report.
- Malheur National Forest (MNF). 2006. Malheur National Forest Range Monitoring Guidelines. October 16, 2011.
- Malheur National Forest (MNF). 2007. Biological assessment for grazing activities on the Rail Creek Allotment. Prairie City Ranger District. September 28, 2007. 23 p.
- Malheur National Forest (MNF). 2009a. 2009 end of year grazing report: Blue Mountain Ranger District, Malheur National Forest. December 16, 2009.
- Malheur National Forest (MNF). 2009b. 2008 end of year grazing report for the Blue Mountain Ranger District of the Malheur National Forest. January 7, 2009.
- Malheur National Forest (MNF). 2011. 2010 end of year grazing report: Blue Mountain Ranger District, Malheur National Forest. January 6, 2011.
- McDowell, P.F. and A. Mowry. 2002. Geomorphic Response to Exclosures *in* Kaufman, J, P. McDowell, P. Bayley, H. Li, R. Beschta, "Research /Evaluate Restoration of NE Oregon Streams", Project No. 2000-05100, 93 electronic pages, (BPA Report DOE/BP-00006210-1)
- National Marine Fisheries Service (NMFS). 1996. Making ESA Determinations of Effect for Individual or Grouped Actions at the Watershed Scale. NOAA Fisheries, Environmental and Technical Services Division, Habitat Conservation Branch, 525 NE Oregon Street, Portland, Oregon. 28 p. (Available @ www.nwr.noaa.gov under Habitat Conservation Division, Habitat Guidance Documents).
http://www.nwr.noaa.gov/Publications/Guidance-Documents/upload/matrix_1996.pdfU3T

- National Marine Fisheries Service (NMFS). 2000. Biological Opinion for the effects to anadromous salmonids from continued implementation of land and resource management plans and resource management plans as amended by the interim strategy for managing fish producing watersheds in eastern Oregon and Washington, Idaho, western Montana, and portions of Nevada (INFISH), and the interim strategy for managing anadromous fish-producing watersheds in eastern Oregon and Washington, Idaho, and portions of California (PACFISH).
- National Marine Fisheries Service (NMFS). 2004. Consultation on Remand for Operation of the Columbia River Power System and 19 Bureau of Reclamation Projects in the Columbia Basin. NOAA Fisheries Log Number: F/NWR/2004/00727. November 30.
- National Marine Fisheries Service (NMFS). 2005. Final Assessment of NOAA Fisheries Critical Habitat Analytical Review Teams for 12 Evolutionarily Significant Units of West Coast Salmon and Steelhead (August 2005), including Appendix J: Initial CHART Assessment for the Middle Columbia River Steelhead.
- National Marine Fisheries Service (NMFS). 2007a. Letter of concurrence for the Blue Mountain expedited Section 7 consultation process to the Forest Supervisor of the Malheur National Forest. NMFS no. 2007/02970.
- National Marine Fisheries Service (NMFS). 2007b. Endangered Species Act - Section 7 Consultation Biological Opinion and Magnuson-Stevens Fishery Conservation and Management Act Essential Fish Habitat Consultation: Malheur National Forest 2007-2011 Administration of Thirteen Grazing Allotments, North Fork John Day Subbasin (HUC 17070202), Middle Fork John Day Subbasin (HUC 17070203), Upper John Day Subbasin (HUC 17070201), Grant County, Oregon. NMFS No. 2007/01290.
- National Marine Fisheries Service (NMFS). 2008. Endangered Species Act Section 7 Formal Programmatic Consultation and Magnuson-Stevens Fishery Conservation and Management Act Essential Fish Habitat Consultation for Fish Habitat Restoration Activities in Oregon and Washington. NMFS No. FS: 2008/03505. NMFS No. BLM: 2008/03506.
- National Marine Fisheries Service (NMFS). 2009. Middle Columbia River Steelhead ESAR Recovery Plan. Available at: <http://www.nwr.noaa.gov/Salmon-Recovery-Planning/Recovery-Domains/Interior-Columbia/Mid-Columbia/upload/Mid-C-Prop-Plan.pdf>
- Northwest Power and Conservation Council (NPCC). 2005. John Day Subbasin Plan. Prepared by Columbia-Blue Mountain Resource Conservation and Development Area for The NPCC. <http://www.nwcouncil.org/fw/subbasinplanning/johnday/plan/PlanRevised.pdf>
- ONDA v. Tidwell, Civ. No. 07-1871-HA, Docket #235.
- Oregon Department of Environmental Quality (ODEQ). 2010. John Day River Basin Total Maximum Daily Load (TMDL) and Water Quality Management Plan (WQMP). November 2010.
- Oregon Department of Fish and Wildlife (ODFW). 2003. Oregon's Mule Deer Management Plan. February 2003. 29 p.

- Oregon Department of Fish and Wildlife (ODFW). 2010. Conservation and Recovery Plan for Oregon Steelhead Populations in the Middle Columbia River Steelhead Distinct Population Segment.
- Park, C.S. 1993. SHADOW stream temperature management program. USDA, USFS, Pacific Northwest Region.
- Platts, W. S. 1981. Influence of forest and rangeland management on anadromous fish habitat in western North America -effects of livestock grazing. USDA Forest Service Gen. tech. Report PNW-124. 25 p.
- Platts, W. S. 1991. Livestock grazing. pp. 389-424 in Meehan, ed., Influences of Forest and Rangeland Management on Salmonid Fishes and Their Habitats. American Fisheries Soc., Bethesda, Maryland. 751 p.
- Platts, W.S., C. Armour, G.D. Booth, M. Bryant, J.L. Bufford, P. Cuplin, S. Jensen, G.W. Lienkaemper, G.W. Minshall, S.B. Monsen, R.L. Nelson, J.R. Sedell, and J.S. Tuhy. 1987. Methods for evaluating riparian habitats with application to management. General Technical Report INT-221. Intermountain Research Station, U.S. Department of Agriculture Forest Service. Ogden, UT.
- Powell, Russ, Kenneth Delano, "John Day River Subbasin Fish Habitat Enhancement Project", 2002-2003 Annual Report, Project No. 198402100, 31 electronic pages, (BPA Report OE/BP-00005632-3)
- Prichard, D., C. Bridges, R. Krapf, S. Leonard, and W. Hagenbuck. 1994. Riparian Area Management: Process for Assessing Proper Functioning Condition for Lentic Riparian-Wetland Areas. TR 1737-11. Bureau of Land Management, BLM/SC/ST-94/008+1737, Service Center, CO. 37 pp.
- Prichard, D., J. Anderson, C. Correll, J. Fogg, K. Gebhardt, R. Krapf, S. Leonard, B. Mitchell, and J. Staats. 1998. Riparian Area Management TR 1737-15. A User Guide to Assessing Proper Functioning Condition Under the Supporting Sciences for Lotic Areas. National Business Center, BC-650B, P.O. Box 25047, Denver, Colorado.
- Reiser, D. W. and R. G. White. 1988. Effects of two sediment-size classes on survival of steelhead and Chinook salmon eggs. North American Journal of Fisheries Management 8:432-437.
- Rosgen, D. 1996. Applied River Morphology. Wildland Hydrology. Pagosa Springs, Colorado.
- Sheer, Mindi B. et. al. 2008. Development and Management of Fish Intrinsic Potential Data and Methods: State of the IP 2008 Summary Report.
- Spence, B.C. and G.A. Lomnický, R.M. Huges, R.P. Novitzki. 1996. An Ecosystem Approach to Salmonid Conservation. TR-4501-96-6057. Management Technology. 356 pp.
<http://www.nwr.noaa.gov/1habcon/habweb/habguide/ManTech/front.htmU3T>
- Stewart, K. M., R. T. Bower, J. G. Kie, B. L. Dick, M. Ben-David. 2003. Niche partitioning among mule deer, elk, and cattle: do stable isotopes reflect dietary niche? Ecoscience 10(3): 297-302.
- Suttle, K. B., M. E. Power, J. M. Levine, and C. McNeely. 2004. How fine sediment in riverbeds impairs growth and survival of juvenile salmonids. Ecological Applications 14:969-974.

- Tappel, P. D. and T. C. Bjornn. 1983. A new method of relating size of spawning gravel to salmonid embryo survival. *North American Journal of Fisheries Management* 3:123-135.
- Torland, Ryan. 2011. Personal communication. Oregon Department of Fish and Wildlife, John Day Oregon.
- University of Idaho Stubble Height Study Team. 2004. University of Idaho Stubble Height Study Report. Submitted to Idaho State Director, BLM, and Regional Forester, Region 4, US Forest Service. University of Idaho Forest, Wildlife and Range Experiment Station, Moscow, ID. 26p.
- USDA Forest Service, Regions 1, 4 and 6. 1991. Columbia River Basin Anadromous Fish Habitat Management Policy and Implementation Guide. 30 p.
- USDA Forest Service. 2001. Forest Roads - A Synthesis of Scientific Information. General Technical Report GTR-509. May 2001.
- USDA Forest Service and USDA Bureau of Land Management. 1994. Environmental assessment for the implementation of interim strategies for managing anadromous fish-producing watersheds in Eastern Oregon and Washington, Idaho, and portions of California (PACFISH).
- U.S. Environmental Protection Agency (EPA). 1993. Monitoring protocols to evaluate water quality effects of grazing management on western rangeland streams. Region 10, Seattle, WA. 179 p.
- Waters, T. 1995. Sediment in streams: sources, biological effects and control. *American Fisheries Society Monograph* 7.
- Winward, A. H. 2000. Monitoring the vegetation resources in riparian areas. Gen. Tech. Rep. RMRS-GTR-47. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station.
- Wright, K. and J. Li. 2002. From continua to patches: examining stream community structure over large environmental gradients. *Canadian Journal of Fisheries and Aquatic Sciences* 59: 1404-1417.