


2011 Aerial Insect and Disease Survey

USGS 100K Quad: VANCOUVER - E145122; 3G



Mortality Agents		
Code	Damaging Agent	Primary Host
1	Douglas fir beetle	Douglas fir
2	Douglas fir engraver	Douglas fir
3	Spineless knitter	True fir
4	Pine engraver	True fir
5	Western balsam bark beetle	Sub-alpine fir
6B	Mountain pine beetle	Whitebark pine
6L	Mountain pine beetle	Lodgepole pine
6P	Mountain pine beetle	Ponderosa pine
6S	Mountain pine beetle	Sugar pine
6W	Mountain pine beetle	Western white pine
7	Wet pine beetle	Ponderosa, longleaf pines
8	Western pine beetle	Ponderosa pine
9	Wet pine beetle	Pine-bark, ponderosa pine
9B	Wet pine beetle	Bark beetle
9L	Wet pine beetle	Leaf beetle
9S	Wet pine beetle	Silver fir, true fir
9W	Wet pine beetle	Western white pine
FL	Flatheaded woodborer	Douglas fir
RD	Black stain root disease	Douglas fir, ponderosa pine
RL	Pine Olfactor cedar root disease	Pine Olfactor cedar
RO	Root disease	Cedar
WATR	Water Damage	All species

Other Damaging Agents		
Code	Damaging Agent	Primary Host
AB	Balsam woolly adelgid	True fir
AC	Cooley spruce gall adelgid	Spruce, Douglas fir
AM	Leaf discoloration	Maple
BR	Bleeder rust	True fir
CC	Chrysomelid canker	True fir
DI	Dying hemlock	All species
DRE	Fire	All species
HD	Hardwood decline	Hardwoods
HDA	Hardwood decline	Aspen
HDC	Hardwood decline	Oak
MN	Aravae nut flower - non host	All species
MTN	Aravae nut flower - host	All species
PAID	Pacific madrone decline	Pacific madrone
RF	Rust	Maple
RS	Rust leaf	All species
SLD	Silver	All species
WATR	Water damage	All species
WIND	Windthrow	All species
WINT	Winter damage	All species

Defoliators		
Code	Damaging Agent	Primary Host
BS	Western spruce budworm	True fir, Douglas fir, spruce
CH	Larch casebearer/typhodermata	Western larch
HL	Western hemlock looper	Western hemlock
LC	Needle cast	Lodgepole pine
LS	Black pine leaf scale	Ponderosa pine
ML	Larch budmoth	Western larch
PH	Pine butterfly	Ponderosa pine
PC	Pine needle cast	Ponderosa pine
PC	Needle cast	Cedar
SP	Sawfly	True fir
SM	Satin moth	Aspen
SNC	Swiss needle cast	Douglas fir
SP	Sawfly	Ponderosa pine
TA	Tent caterpillar, alder	Alder
TM	Douglas fir tussock moth	Truefir, Douglas fir

USGS 100K Quad: VANCOUVER - E145122; 3G
2011 Aerial Insect and Disease Survey
Map Scale: 1:100,000
Date: 13 December 2011

Legend

	Defoliating Agents		Areas Not Flown
	Mortality Agents		2011 Large Fires
	Other Damage		Source: Northwest Interagency Coordination Center
	2011 Special Swiss Needle Cast Survey		

More information about this special survey and the related data is located here: <http://www.oregon.gov/ODF/privateforests/ff/Maps.shtml>

The TOPOI maps are seamless, scanned images of United States Geological Survey (USGS) paper topographic maps. For more information on this map, visit us online at http://gto.arcgisonline.com/maps/USA_Topo_Maps

A data dictionary, digital copies of this map and Arctis insect and disease data are available at www.fs.usda.gov/gto/r6/fhp/ads

How the Aerial Surveys Are Conducted

Data represented on this map are based on trees visibly affected by forest insects and diseases detected and recorded during aerial survey flights conducted by the USDA Forest Service, the Washington Department of Natural Resources and the Oregon Department of Forestry. Observers have just a few seconds to recognize the color difference between healthy and damaged trees of different species; diagnose causal agents correctly; estimate intensity; delineate the extent of damage; and precisely record this information on a georeferenced, digital map. Air turbulence, cloud shadows, distance from aircraft, haze, smoke and observer experience can all affect the quality of the survey. These data summaries provide an estimate of conditions on the ground and may differ from estimates derived by other methods.

The aerial survey provides information on the current status for many causal agents, and is important when examining insect activity trends by comparing historical and current survey data over large areas.

Overview surveys are a 'snap shot' in time and therefore may not be timed to accurately capture the true extent or severity of a particular disturbance activity. Specially designed surveys with modified flight patterns and timing may be conducted to more accurately delineate the extent and severity of a particular disturbance agent. Separate surveys, such as Swiss needle cast surveys, are conducted when resources are available to address situations of sufficient economic, political or environmental importance.

WASHINGTON STATE DEPARTMENT OF Natural Resources

OREGON DEPARTMENT OF FORESTRY

U.S. FOREST SERVICE

DIRECT ALL INQUIRIES TO:

Washington State Department of Natural Resources
 Resource Protection Division
 Forest Health
 1111 Washington St. SE
 MS 47037
 Olympia, WA 98504

-- OR --

Oregon Department of Forestry
 Forest Health Management
 2600 State Street
 Salem, OR 97310

-- OR --

USDA Forest Service, Region 6
 Natural Resources
 Forest Health Protection
 PO Box 3623
 Portland, Oregon 97208

DISCLAIMER

Forest Health Protection (FHP), Washington Department of Natural Resources (WONR) and Oregon Department of Forestry (ODF) strive to maintain an accurate Aerial Detection Survey (ADS) Dataset, but due to the conditions under which the data are collected FHP, WONR and ODF shall not be held responsible for missing or inaccurate data. ADS are not intended to replace more specific information. An accuracy assessment has not been done for this dataset; however, ground checks are completed in accordance with local and national guidelines. <http://www.fs.usda.gov/foresthealth/operations/qualityassurance.shtml>. Maps and data may be updated without notice. Please cite: "USDA Forest Service, Forest Health Protection, Washington Department of Natural Resources, Resource Protection Division, and Oregon Department of Forestry, Forest Health Management" as the source of this data.