


2011 Aerial Insect and Disease Survey

USGS 100K Quad: LAKE ABERT - E142120; 5M



Mortality Agents			Other Damaging Agents		
Code	Damaging Agent	Primary Host	Code	Damaging Agent	Primary Host
1	Douglas fir beetle	Douglas fir	AB	Balsam woolly adelgid	True fir
2	Douglas fir engraver	Douglas fir	AC	Cooley spruce gall adelgid	Spruce, Douglas fir
3	Spineless bark beetle	Spineless bark beetle	AM	Leaf miner	Various
4	Pine engraver	True fir	AN	Leaf miner	Various
5	Western balsam bark beetle	Sub-alpine fir	BR	Bitter root	Various
6B	Mountain pine beetle	Whitebark pine	CC	Chrysomelid canker	True fir
6L	Mountain pine beetle	Lodgepole pine	DC	Dying hemlock	Various
6P	Mountain pine beetle	Ponderosa pine	DI	Dwarf mistle	All species
6S	Mountain pine beetle	Sugar pine	DR	Dwarf mistle	Various
6W	Mountain pine beetle	Western white pine	FD	Fern root disease	Various
7	Wet spruce beetle	Pacific silver pine	HC	Hardwood decline	Hardwoods
8	Western pine beetle	Ponderosa pine	HCO	Hardwood decline	Hardwoods
9	Western pine beetle	Pine-needle scale	HN	Arava nut flower - non host	Various
9A	Western pine beetle	Pine-needle scale	MTN	Arava nut flower - host	Various
9B	Western pine beetle	Pine-needle scale	PAID	Pacific madrone decline	Pacific madrone
9C	Western pine beetle	Pine-needle scale	PR	Pine root rot	Various
9D	Western pine beetle	Pine-needle scale	RS	Rail rust	Various
9E	Western pine beetle	Pine-needle scale	SLD	Shade	All species
9F	Western pine beetle	Pine-needle scale	WATR	Water damage	All species
9G	Western pine beetle	Pine-needle scale	WIND	Windthrow	All species
9H	Western pine beetle	Pine-needle scale	WNTN	Winter damage	All species
9I	Western pine beetle	Pine-needle scale			
9J	Western pine beetle	Pine-needle scale			
9K	Western pine beetle	Pine-needle scale			
9L	Western pine beetle	Pine-needle scale			
9M	Western pine beetle	Pine-needle scale			
9N	Western pine beetle	Pine-needle scale			
9O	Western pine beetle	Pine-needle scale			
9P	Western pine beetle	Pine-needle scale			
9Q	Western pine beetle	Pine-needle scale			
9R	Western pine beetle	Pine-needle scale			
9S	Western pine beetle	Pine-needle scale			
9T	Western pine beetle	Pine-needle scale			
9U	Western pine beetle	Pine-needle scale			
9V	Western pine beetle	Pine-needle scale			
9W	Western pine beetle	Pine-needle scale			
9X	Western pine beetle	Pine-needle scale			
9Y	Western pine beetle	Pine-needle scale			
9Z	Western pine beetle	Pine-needle scale			

USGS 100K Quad: LAKE ABERT - E142120; 5M
2011 Aerial Insect and Disease Survey
Map Scale: 1:100,000
Date: 14 December 2011

Legend

- Defoliating Agents
- Mortality Agents
- Other Damage
- Areas Not Flown
- 2011 Large Fires

Source: Northwest Interagency Coordination Center

The cause of damage is described by a symbol above and is followed by: number of trees affected; number of trees (example: SA) or intensity of damage (L - Light, M - Moderate, H - Heavy).

The TOPO! maps are seamless, scanned images of United States Geological Survey (USGS) paper topographic maps. For more information on this map, visit us online at http://gto.arcgis.com/arcgis/rest/services/USA_Top_Maps

A data dictionary, digital copies of this map and Arctics insect and disease data are available at: www.fs.usda.gov/gto/r6/fhp/ads

How the Aerial Surveys Are Conducted

Data represented on this map are based on trees visibly affected by forest insects and diseases detected and recorded during aerial survey flights conducted by the USDA Forest Service, the Washington Department of Natural Resources and the Oregon Department of Forestry. Observers have just a few seconds to recognize the color difference between healthy and damaged trees of different species; diagnose causal agents correctly; estimate intensity; delineate the extent of damage; and precisely record this information on a georeferenced, digital map. Air turbulence, cloud shadows, distance from aircraft, haze, smoke and observer experience can all affect the quality of the survey. These data summaries provide an estimate of conditions on the ground and may differ from estimates derived by other methods.

The aerial survey provides information on the current status for many causal agents, and is important when examining insect activity trends by comparing historical and current survey data over large areas.

Overview surveys are a 'snap shot' in time and therefore may not be timed to accurately capture the true extent or severity of a particular disturbance activity. Specially designed surveys with modified flight patterns and timing may be conducted to more accurately delineate the extent and severity of a particular disturbance agent. Special surveys, such as Swiss needle cast surveys, are conducted when resources are available to address situations of sufficient economic, political or environmental importance.

DIRECT ALL INQUIRIES TO:

Oregon Department of Forestry
Forest Health Management
2600 State Street
Salem, OR 97310

-- OR --

USDA Forest Service, Region 6
Natural Resources
Forest Health Protection
PO Box 3623
Portland, Oregon 97208

DISCLAIMER
 Forest Health Protection (FHP), Washington Department of Natural Resources (WDNR) and Oregon Department of Forestry (ODF) strive to maintain an accurate Aerial Detection Survey (ADS) Dataset, but due to the conditions under which the data are collected, FHP, WDNR and ODF shall not be held responsible for missing or inaccurate data. ADS are not intended to replace more specific information. An accuracy assessment has not been done for this dataset; however, ground checks are completed in accordance with local and national guidelines. <http://www.fs.fed.us/foresthealth/ads/>
 quality assurance sheet. Maps and data may be updated without notice. Please cite: "USDA Forest Service, Forest Health Protection, Washington Department of Natural Resources, Resource Protection Division, and Oregon Department of Forestry, Forest Health Management" as the source of this data.