



United States
Department of
Agriculture

Forest
Service

**Southwestern
Region**



Forest Products Specialist Report

Forest Plan Revision DEIS

Submitted by: /s/ *Monica Boehning* _____

Monica Boehning
Forest Silviculturist
Apache-Sitgreaves National Forests
November 30, 2012

Contents

Forest Products Specialist Report..... 4

Introduction	4
Relevant Laws, Regulations, and Policy that Apply to Silvicultural Actions.....	4
Federal Statutes.....	4
Regulations	6
Forest Service Directives	6
Programmatic Agreements.....	7
Methodology and Analysis Process	9
Revision Topics Addressed in this Analysis	17
Summary of Alternatives	18
Affected Environment.....	18
Lands Tentatively Suitable for Timber Production	21
Adjustments to the suitable timberland acreage within the Wallow Fire and other high-severity fires may be appropriate in the next 10 years during the scheduled review and update of the forest suitability classification process.	23
Allowable Sale Quantity (ASQ)	23
Long-Term Sustained-Yield Capacity (LTSYC).....	24
Wood and Tree Products Availability.....	26
Environmental Consequences of Alternatives	29
Suitable Timberlands	30
Allowable Sale Quantity (ASQ)	33
Long-Term Sustained-Yield Capacity.....	37
Total Wood Products.....	41
Relationship of Short-Term Management to Long-Term Productivity	52
Climate Change Considerations.....	54
Cumulative Environmental Consequences	56
Unavoidable Adverse Environmental Consequences	60
Irreversible and Irretrievable Commitment of Resources	60
Adaptive Management.....	60
Other Planning Efforts.....	61
References	62
The following information sources were reviewed and/or consulted for this analysis.	62
Appendices - Forest Products.....	68

Forest Products Specialist Report

Introduction

This report evaluates and discloses the potential environmental and social consequences to the forest, piñon-juniper woodland, and great basin grassland resources, with respect to lands suitable for timber production (hereafter referred to as “suitable timberlands”), vegetation management practices, and potential wood and tree products contributing to the local communities, which may result with the adoption of a revised land management plan. It examines, in detail, four different alternatives for revising the 1987 Apache-Sitgreaves NFs land management plan (1987 forest plan).

Relevant Laws, Regulations, and Policy that Apply to Silvicultural Actions

Federal Statutes

A Federal statute, or law, is an act or bill which has become part of the legal code through passage by Congress and approval by the President (or via congressional override). Although not specified below, many of these laws have been amended.

Anderson-Mansfield Reforestation and Revegetation Act of October 11, 1949 - Provides for the reforestation and revegetation of National Forest System lands and other lands under the administration or control of the Forest Service.

Federal Land Policy and Management Act of October 21, 1976 - Requires that public lands be managed in a manner that will protect the quality of scientific, scenic, historical, ecological, environmental, air and atmospheric, water resource, and archeological values; that, where appropriate, will preserve and protect certain public lands in their natural condition; that will provide food and habitat for fish and wildlife and domestic animals; and that will provide for outdoor recreation and human occupancy and use. Also states that the United States shall receive fair market value of the use of the public lands and their resources unless otherwise provided for by law.

Forest and Rangeland Renewable Resources Planning Act of August 17, 1974, as amended by the National Forest Management Act (NFMA) of October 22, 1976 - The National Forest Management Act reorganized, expanded, and otherwise amended the Forest and Rangeland Renewable Resources Planning Act of 1974, which called for the management of renewable resources on National Forest System lands. The National Forest Management Act requires the Secretary of Agriculture to assess forest lands, develop a management program based on multiple-use, sustained-yield principles, and implement a resource management plan for each unit of the National Forest System. It is the primary statute governing the administration of National Forests.

Healthy Forests Restoration Act (HFRA) of 2003 (H.R. 1904)- Purposes are to reduce wildfire risk to communities and municipal water supplies through collaborative hazardous fuels reduction projects; to assess and reduce the risk of catastrophic fire or insect or disease infestation; to enhance efforts to protect watersheds and address threats to forest and rangeland health (including wildfire) across the landscape; to protect, restore, and enhance forest ecosystem components such as biological diversity, threatened/endangered species habitats, enhanced productivity.

Knutson-Vandenberg Act of June 9, 1930 - Authorizes the Secretary of Agriculture to establish forest tree nurseries; to deposit monies from timber sale purchasers to cover the costs of planting young trees, sowing seed, removing undesirable trees or other growth, and protecting and improving the future productivity of the land; and to furnish seedlings and/or young trees for the replanting of burned-over areas in any National Park.

Multiple-Use Sustained-Yield Act (MUSY) of June 12, 1960 -States that it is the policy of Congress that the national forests are established and shall be administered for outdoor recreation, range, timber, watershed, and wildlife and fish purposes, and authorizes and directs the Secretary of Agriculture to develop and administer the renewable surface resources of the national forests for the multiple-use and sustained-yield of products and services.

National Environmental Policy Act of January 1, 1970 - Directs all Federal agencies to consider and report the potential environmental impacts of proposed Federal actions, and established the Council on Environmental Quality.

Organic Administration Act of June 4, 1897 - Authorizes the President to modify or revoke any instrument creating a national forest; states that no national forest may be established except to improve and protect the forest within its boundaries, for the purpose of securing favorable conditions of water flows, and to furnish a continuous supply of timber for the use and necessities of citizens of the United States. Authorizes the Secretary of Agriculture to promulgate rules and regulations to regulate the use and occupancy of the national forests.

Supplemental National Forest Reforestation Fund Act of September 18, 1972 - Directs the Secretary of Agriculture to establish a supplemental national reforestation fund, and states that money transferred to this fund shall be available to the Secretary for the purpose of supplementing programs of tree planting and seeding on National Forest System lands determined by the Secretary to be in need of reforestation.

Stewardship End Result Contracting Projects (16 U.S.C. 2104) - Grants the Bureau of Land Management and the Forest Service ten-year authority to enter into stewardship contracts or agreements to achieve agency land management objectives and meet community needs.

Sustained Yield Forest Management Act of March 29, 1944 - Authorizes the Secretaries of Agriculture and the Interior to establish by formal declaration cooperative sustained-yield units which shall consist of federally owned or administered forest land under their jurisdiction and, in addition thereto, land which reasonably may be expected to be made the subject of one or more of the cooperative agreements with private landowners authorized by section 2 of the Act in order to promote the stability of forest industries, of employment, of communities, and of taxable forest wealth through continuous supplies of timber and forest products; and in order to secure the benefits of forests in the maintenance of water supply, regulation of stream flow, prevention of soil erosion, amelioration of climate, and preservation of wildlife.

Timber Export Act of March 4, 1917 - Permits the Secretary of Agriculture to allow timber or other forest products to be cut or removed from a national forest and exported from the state or territory in which that national forest is situated.

Timber Exportation Act of April 12, 1926 - Authorizes the exportation of lawfully cut timber from the state or territory where grown if the supply of timber for local use will not be endangered, and authorizes the Secretary to issue rules and regulations to carry out the provisions of the Act.

Tribal Forest Protection Act of 2004 - Authorizes the Secretary of Agriculture and the Secretary of the Interior to enter into an agreement or contract with Indian tribes meeting certain criteria to carry out projects to protect Indian forest land or rangeland, including a project to restore Federal land that borders on or is adjacent to Indian forest land or rangeland.

Twenty-Five Percent Fund Act of May 23, 1908 - Provides that twenty-five percent of all monies received from the sale of timber or other forest products shall be paid to the state in which such forest is located to be expended as the state may prescribe for the benefit of public schools and roads.

Wood Residue Utilization Act December 19, 1980 - Enacted to develop, demonstrate, and make available information on feasible methods that have the potential for commercial application to increase and improve utilization in residential, commercial, and industrial or power plant applications of wood residues resulting from timber harvesting and forest protection and management activities occurring on public and private forest lands, and from the manufacture of forest products, including wood pulp.

Regulations

Below is a partial listing of relevant regulations. Federal executive departments and administrative agencies write regulations to implement laws. Regulations are secondary to law. However, both laws and regulations are enforceable.

36 CFR 221 Timber Management Planning - Sets forth the requirements for management plans for National Forest timber resources.

36 CFR 223 Sale and Disposal of National Forest System Timber - Sets forth the requirements relating to the sale and disposal of National Forest System timber.

Forest Service Directives

The Forest Service Manual (FSM) and Forest Service Handbooks (FSH) contains legal authorities, goals, objectives, policies, responsibilities, instructions, and the necessary guidance to plan and execute assigned programs and activities.

FSM 2000 National Forest Resource Management

- **FSM 2020** Ecological Restoration and Resilience
- **FSM 2070** Biological Diversity
 - **FSM 2070.3** Vegetation Ecology (use of native plants in re-vegetation, rehabilitation, and restoration)

FSM 2400 Timber Management, Southwestern Region and Apache-Sitgreaves NFs supplements

- **FSM 2430** Commercial Timber Sales, Southwestern Region and Apache-Sitgreaves NFs supplements, Small Sales and Commercial/Personal Use Permits of Timber, Fuelwood, and other forest products
- **FSM 2470** Silvicultural Practices
- **FSH 2409.13** Timber Resource Planning Handbook, Ch. 30 – Timber Sale Scheduling, WO Amendment 2409.13-92-1, Effective 8/3/92. Provides optional agency direction as recommended guidance.

Programmatic Agreements

Memorandum of Understanding among the Arizona Game and Fish Department, New Mexico Game and Fish Department, U.S.D.A. Animal and Plant Health Inspection Service/Wildlife Services, U.S.D.A Forest Service, U.S. Fish and Wildlife Service, White Mountain Apache Tribe, Arizona Counties of Graham, Greenlee, and Navajo, New Mexico Counties of Catron and Sierra, and the New Mexico Department of Agriculture

Glossary

Terms and Abbreviations Used Throughout This Report and Appendices:

ASNFs – Apache-Sitgreaves National Forests

ASQ – Allowable Sale Quantity

BA – Basal Area. A unit of stand/forest density measure, using the cross-sectional area of all tree boles standing in a given land area. Expressed as square feet per acre.

Bole - The cylindrical portion of a tree trunk.

CF – Cubic feet of wood volume (used as a per-acre unit measure).

CCF - 100 Cubic Feet of wood volume (used as a total unit measure across all acres treated).

CWPP - Community Wildfire Protection Plan.

DBH – Diameter Breast Height (tree bole measurement at 4.5 feet above ground).

GIS - Geographical Information System.

FIA - Forest Inventory and Analysis, permanently located plots across National Forest Service system lands.

FFE - Fire-Fuels Extension of the FVS model.

FVS - Forest Vegetation Simulator Model (and all its submodels and extensions).

Gross growth – Ingrowth plus accretion. A measurable increase in wood volume due to the addition of new trees per acre added or grown into size classes which count toward total stand volume (ingrowth), plus added increases in tree diameter increment and height of trees already existing in those same size classes (accretion).

LTSYC – Long Term Sustained Yield Capacity

MA - Management Area (in the forest plan).

NEPA - National Environmental Policy Act.

Net growth - Gross growth in forest wood volume minus natural (non-cut) mortality volume.

Non-declining even flow (of ASQ volume) – Concept of long-term sustained yield which legally prohibits the ASQ volumes harvested from suitable timberlands from declining from one decade to the next (see additional handbook definitions below).

MMBF - Million Board Feet (of timber)

Planning period –Length of time (15 years) the revised plan is expected to be in effect.

PNVT - Potential Natural Vegetation Type (see Vegetation Specialist report)

Regulated (forest) – A balanced progression of age/size classes are present at consistent growth rates to ensure a regular harvest of relatively similar volume each entry, for long-term sustained yield.

TIM – Timber Information Manager (wood/tree product sales program and database)

TPA - Trees per Acre.

VDDT - Vegetation Dynamic Development Tool model.

WMSP - White Mountain Stewardship Project

In context of provisions of the 1982 planning rule, a number of key terms and definitions are presented and explained here. They are not necessarily arranged here in alphabetical order, but intentionally flow in a logical process to enable the reader to digest the concepts. Certain key words are emphasized in italics here for the reader’s benefit. (Source: Appendix A1 contains all definitions provided in the 1992 WO amendment to FSH 2409.13, zero code.)

Suitable Timberland and Allowable Sale Quantity Definitions:

Timber production is strictly defined as “the purposeful growing, tending, harvesting, and regeneration of regulated crops of trees for cutting into logs, bolts, or other round sections for industrial or consumer use. For purposes of forest planning, timber production does not include *fuelwood* or harvests from *unsuitable lands*.” (FSM 1900)

Fuelwood is strictly defined as “wood that is round, split, or sawn and/or otherwise generally refuse material cut into short lengths or chipped for burning.”

Unsuitable lands are defined as “forest land not managed for *timber production* because: (a) Congress, the Secretary, or the Chief has withdrawn it; (b) it is not producing or capable of producing crops of *industrial wood*; (c) technology is not available to prevent irreversible damage to soils productivity, or watershed conditions; (d) there is no reasonable assurance based on existing technology and knowledge, that it is possible to restock lands within 5 years after final harvest, as reflected in current research and experience; (e) there is, at present, a lack of adequate information about responses to timber management activities; or (f) timber management is inconsistent with or not cost efficient in meeting the management requirements and multiple-use objectives specified in the forest plan.

Tentatively suitable is defined as “forest land that is producing or is capable of producing crops of *industrial wood* and: (a) has not been withdrawn by Congress, the Secretary, or the Chief; (b) existing technology and knowledge is available to ensure timber production without irreversible damage to soils productivity, or watershed conditions; (c) existing technology and knowledge, as reflected in current research and experience, provides reasonable assurance that it is possible to restock adequately within 5 years after final harvest; and (d) adequate information is available to project responses to timber management activities.” This is essentially the same definition also used for “commercial forest land.”

Industrial wood is strictly defined as “all commercial *roundwood products*, except *fuelwood*.”

Roundwood products are strictly defined as “logs, bolts, or other round sections cut from trees.”

Suitable forest land is defined as “land to be managed for *timber production* on a regulated basis.” Throughout this document, is referred to as “suitable timberlands”.

Allowable Sale Quantity (ASQ) Definitions:

Allowable sale quantity (ASQ) - The quantity of timber that may be sold from the area of suitable land covered by the forest plan for a time period specified by the plan. This allowable sale quantity (ASQ) is usually expressed on an annual basis as the “average annual allowable sale quantity.” For timber resource planning purposes, the allowable sale quantity applies to each decade over the planning horizon and includes only *chargeable volume*. Consistent with the definition of timber production, do not include fuelwood or other nonindustrial wood in the allowable sale quantity.

Chargeable volume - All volume included in the growth and yield projections for the selected management prescriptions used to arrive at the allowable sale quantity, based on regional utilization standards. Consistent with the definition of timber production, planned production of fuelwood is not included in the allowable sale quantity and therefore is nonchargeable.

Base sale schedule - A timber sale schedule formulated on the basis that the quantity of timber planned for sale and harvest for any future decade is equal to or greater than the planned sale and harvest for the preceding decade and that this planned sale and harvest for any decade is not greater than the *long-term sustained-yield capacity*. This definition expresses the *principle of non-declining flow*.

Long-term sustained yield capacity (LTSYC) - The highest uniform wood yield from lands being managed for *timber production* that may be sustained, under specified management intensity, consistent with multiple-use objectives.

Non-declining flow - is defined in the regulations by the statement “see base sale schedule.”

Sale schedule is further defined as “the quantity of timber planned for sale by time period from the area of suitable land covered by a forest plan. The first period, usually a decade, of the selected sale schedule provides the allowable sale quantity. Future periods are shown to establish that long-term sustained yield will be achieved and maintained. For timber resource planning purposes, consider the sale schedule and allowable sale quantity to be synonymous for all periods or decades over the planning horizon.”

Departure - A sale schedule that deviates from the *principle of non-declining flow* by exhibiting a planned decrease in the sale schedule at any time during the planning horizon. A departure is characterized by a temporary increase, usually in the beginning decade(s) of the planning horizon, over the base sale schedule originally established. This increase does not impair the future attainment of the *long-term sustained yield capacity*.

Also see Appendices A2 and F for more explanation of how these terms and other directives are pertinent to this analysis.

Methodology and Analysis Process

Beyond this section, brief disclosure of analysis data and methodology are also provided where appropriate in the affected environment and environmental consequences sections, to assist the reader with concepts and conclusions discussed therein. Several appendices were created to help describe and demonstrate methods of analysis. Due to the size and nature of the appendices, they are all available in the Plan set of documents as separate electronic files. They are listed at the end of this report as an index, and are referenced throughout this report when needed.

This report examines how the plan alternatives contribute to local communities through the availability of wood and tree products. It provides an estimate of how much land (acreage) is available for timber production, cutting levels (CCF per acre) in relation to long-term sustained yield, and allowable sale quantity (CCF) for the first decade. And finally, it estimates the volumes (CCF) of wood products that could be removed from both lands suitable and not suitable for timber production. CCF of sawtimber/pulpwood/poles, CCF of fuelwood/posts, and tons of biomass (converted to CCF) are the only units of wood volume measure analyzed here for a meaningful basis to compare alternatives.

This report only addresses the effects of managing those potential natural vegetation types, (PNVTs) which currently have trees on them, and were modeled in VDDT with cut volume estimates. These PNVTs are:

- Ponderosa pine forested PNVT (includes both the ponderosa pine bunchgrass and ponderosa pine-Gambel oak types);
- Dry mixed conifer forested PNVT (frequent fire regime mixed conifer forest);
- Wet mixed conifer forest with aspen PNVT (infrequent fire regime mixed conifer forest type);
- Spruce-fir forested PNVT (low elevation spruce-fir mixed forest type);
- Piñon-juniper woodland PNVT (piñon-juniper grass woodland). This type is mostly savanna with some persistent woodland; and
- Great Basin grassland PNVT(which is occupied with tree encroachment).

Existing vegetation classification of forest, woodland, and riparian PNVTs and amounts of VDDT structural states were originally mapped using mid-scale technology in 2005-2006. This classification was reassessed with a 2011 (post-Wallow wildfire) update of mid-scale mapping. This update did not address changes in vegetation structure since 2006 due to recent fuels reduction treatments, prescribed burns, tree planting, and other smaller wildfires outside of the Wallow fire. However, it is assumed that it does generally reflect current forest-wide conditions including the major impacts of the 2002 Rodeo-Chediski and 2011 Wallow wildfires. (See further details in the Vegetation Specialist Report.)

The 1987 plan mapped forest and woodlands using the outdated cover type classification, which is based on the visibly dominant tree or plant species at the time of mapping. This analysis uses potential natural vegetation types (PNVTs), which may not be the currently dominant vegetation. For example, what appears to be wet mixed conifer forest (infrequent fire type) may actually be dry mixed conifer (frequent fire type) where shade-tolerant species have become established in the absence of frequent fires. Alternatives B, C, and D are based on a large body of research (see US Forest Service, 2008-ESR; and supporting Vegetation Specialist's analyses and bibliographies) that shows the need to reclassify our lands as PNVTs. These types are based on what vegetation species should be dominating there if all natural processes and disturbances were functioning normally. Without this PNVT concept, true ecological restoration is not possible.

Total forest covert type acres used for alternative A (current plan) were re-calculated to match the PNVT acreages for the action alternatives, in order to provide a fair basis for comparison.

This report relies heavily on results from modeling conducted for this analysis. Two models were used to estimate volumes of wood cut under each alternative: the Vegetation Dynamic Development Tool (VDDT), and the Forest Vegetation Simulator (FVS). FVS gives per-acre averages, while VDDT can handle as many acres as the analysis area contains in any given

PNVT. A major limitation of the FVS and VDDT models is that neither one is a spatial model. Both are dynamic, density- and time-dependent, and each reacts in its own way to changes in existing condition parameters as a result of natural growth and disturbances, including management actions. VDDT simulations were run out for fifty years.

This report utilizes and summarizes the VDDT model results to compare estimated cutting volumes produced from various treatments. The VDDT model only addresses vegetation development (structural) states for each PNVT based on various combinations of three structural attributes: predominant tree bole diameter range (seedling/sapling, small, medium, very large); canopy closure (open, closed); and number of canopy layers (single-storied, multi-storied). See Appendix B3 to this report, and the Vegetation Specialist Report for detailed explanation of the VDDT model, most of its limitations, and methodology for its use in forest planning analysis.

To calibrate VDDT for volumes of wood cut by various treatments, as well as calibrations for post-treatment progression to other VDDT structural states in key PNVTs, the FVS model and FVS sub-models were used extensively in 2010-2011 by the USFS Region 3 Timber Calculations Working Group (Weisz et al., 2012).

Appendices B1 through B5, C, E1, and E2 for this report demonstrate how the VDDT model was calibrated by Region-3 and used by the Apache-Sitgreaves NFs to simulate the different mixtures of prescribed tree cutting, tree planting, and prescribed burning treatments designed to reflect the emphasis of each Apache-Sitgreaves planning alternative for comparative analysis. Silviculture cutting methods and intensities of prescribed burns were modeled in FVS to simulate a range of possible treatments designed to move forests and woodlands toward the desired conditions. (For details of fire simulations conducted in FVS, see Weisz et al., 2012, and the ASNFs Fire Specialist Report.)

In an attempt to move acres of vegetation structural states toward desired conditions for each PNVT, all cutting methods were examined with respect to how well they can reduce percentages of the landscape currently in surplus of the desired amount for that structural state. Similarly, how well each cutting method can create extra acres of those desired states currently in deficit from the desired percentages was examined (see Appendix B3 for an example). Regionally-derived structural state transitions from the FVS model provide this information by percentages (see Appendices B1 and B4). Each alternative model run in VDDT then uses a different mix of those cutting methods to simulate the alternative's different management emphasis (see Appendices B2 and B5). Then different acreage mixtures (based on the high and low range of treatment objectives) for various prescribed cutting and prescribed burning treatments were input into the VDDT model (Appendices C and B5), to approximate just one of several plausible treatment schemes commensurate with the vegetation management emphasis of each alternative.

Product demands (or lack thereof) did not influence methods of cuts used to model alternatives; nor should certain treatments be chosen to implement primarily because they will give the greatest dollar return or the greatest output of timber (NFMA, Sec. 219.27 (a)(12)(b)(3)).

Live green tree total wood volumes cut for restoration treatments by PNVT in each alternative, for decades 1 through 5, were taken directly from VDDT model outputs sorted by industrial species, non-industrial species, size and product classes, measured in CCF, and tons. They were added together for all forested PNVTs in additional spreadsheets. (See Appendices D1 and D2).

Additionally, the total allowable sale quantity (ASQ) volumes from VDDT were adjusted to include conservative average estimates of annual pole permit sales, and of routine small salvage sales that have become a part of the annual forest management program (removals of roadside hazard and developed recreation site hazard trees for public safety, new construction and/or Forest Products Specialist Report - Apache-Sitgreaves NFs Plan Revision DEIS

maintenance of power-line clearings, road realignment and right-of-way clearings), as well as more frequent insect outbreaks and increasing numbers of small wildfires occurring in a warmer/drier climate. They do not include additional salvage volumes resulting from unanticipated large scale wildfire events.

ASQ volume estimates only include chargeable volumes of industrial wood. The ASQ calculation includes restoration green tree cutting volumes from the VDDT model which vary by alternative; plus extra small sales and permits sold for live and/or dead poles and miscellaneous salvage timber which are considered to be a constant addition across all alternatives.

Apache-Sitgreaves NFs average annual dead fuelwood sales sold the public, were added to the live green fuelwood volumes cut in VDDT for restoration treatments, to provide total fuelwood volumes estimates that could be offered.

These total ASQ volumes, as well as total fuelwood and biomass volumes, for Decade 1 were provided to the economics specialist for further analysis in the SocioEconomic Report (see Appendix D2).

Also not included in the wood volume estimates are the additional dead tree volumes indirectly created by intentionally using moderate-intensity fire as a tree-thinning tool (both planned and unplanned ignitions). Only volumes of live trees cut in VDDT were modeled for this analysis, with incidental salvage volumes of small wildfire-killed trees added manually from salvage sale cruise records. The VDDT model was not calibrated to provide volumes of fire-killed snags. If salvage volumes of prescribed fire-killed trees were included, then Alternatives B and D might possibly approach Alternative C, for total wood products available to offer, at least in the first few decades. The Regional FVS-FFE prescribed fire model runs do report numbers of tree per acre and size classes killed by the fire intensity/severity. However, FFE does not calculate the wood volumes of those fire-killed trees. If desired, estimates of this extra prescribed fire-created wood volume would require further analysis of the R3 FVS-FFE fire simulation results.

Regional average long-term sustained yield calculations were derived in FVS by Region 3 for plan revision analysis. See Appendix F (Youtz and Vandedriesche, Sept. 2012); and also see assumptions below.

Wood products resulting from restoration treatments on riparian/broadleaf hardwood and grasslands other than Great Basin PNVTs are not available because they lack adequate tree data to run timber growth and yield models to obtain cutting volumes. Also, a wide range of existing conditions and different restoration treatments may be needed, which vary greatly with respect to conifer trees or other species to be removed to meet the desired conditions for these PNVTs. Thus, estimating volumes requires too much conjecture for reliable comparisons. Any volumes resulting from cuts in these PNVTs would not affect ASQ estimates.

The consequences between using different cutting methods emphasized in the alternatives were also considered, as well as the different emphases on using prescribed fire between the alternatives. Those implementation differences were included in VDDT modeling, and are summarized in Appendices B2, B5, E1 and E2. In general, Alternatives A, B, and C would use a wide variety of mechanical treatment methods to meet ecological needs. Alternative B, within the ponderosa pine type, would emphasize the retention of more large/old trees than Alternative C would; therefore some cutting methods would be limited. Alternative D would retain all large/old trees practically everywhere, so that cutting options under this alternative would be much more limited than those under the other alternatives.

Regional average natural regeneration rates are designed to automatically occur in each VDDT model run, depending on the amounts of appropriate structural states present. Additionally, artificial reforestation efforts were modeled in the VDDT simulations for those structural states that need tree cover replaced as a desired condition. Annual acres of tree planting were assigned commensurate with each alternative's overall management emphasis, and were kept within realistic levels possible according to current workforce and funding capabilities. Successful plantation establishment was assumed.

The annual planned cutting acres input into VDDT in year 1 were found to not always be the same acreage treated internally within the model. This is due to the dynamic nature of this model, which continually is tracking acreage transitions of structural states to acres of other states. Some acres, or portions of acres, may fit a particular cutting prescription at a certain point in time, while others may not. This is realistic in the sense that any given project area will propose to treat a certain number of acres, but the number of acres on which trees are actually cut varies a bit when implemented on the ground.

The total acres of suitable timberlands for each PNVT used in all VDDT model runs do not necessarily match the final acres classified as suitable. This is because VDDT modeling was given higher priority in the analysis schedule than completion of the cost-efficiency step for timberland suitability classification. Therefore, the acres removed from suitability due to cost-inefficiency are included in the total suitable acreage programmed in each VDDT model run. However, this total number turns out to be inconsequential because only the prescribed annual cutting acres on suitable lands were used to provide the resulting volumes and ecological trends provided by VDDT. Likewise, this same modeling inconsistency exists across all alternatives equally, so it makes no difference for the decision-maker in choosing between alternatives.

In the case of this analysis, neither model was used to simulate changes in methods of cut from one treatment entry to the next on the same acre. Treatments can be turned off in VDDT, using the "time-since-disturbance" feature, as was done for single-entry clearcuts done on grasslands and cuts implemented in Alternative D. But in cases where several cutting entries are expected on the same land, like on forested acres in Alternatives A, B, and C, the exact same prescribed cutting method (or prescribed burn severity) input into VDDT to implement for year 1 is repeatedly implemented again every time the same acre is due for another treatment some number of years later.

So a logical sequence of different silvicultural treatments that would normally be prescribed for any piece of ground over time was not modeled in our use of VDDT. For example, if an intermediate thinning cut was input for a certain number of acres in Entry1, that same thinning method and target basal area was replicated in VDDT for those same acres again in each subsequent entry. No future regeneration cut or conversion to the group selection cutting method was possible for those same acres in the model runs.

Alternatives A and C were found to comply with the non-declining even-flow legal requirement by continuing the same treatment strategy each decade as was used in the initial level of VDDT modeling. In the case of alternative B however, the initial VDDT model runs which repeated the same treatment strategy in subsequent decades after this planning period produced ASQ volumes that consistently declined each decade, while staying below the long term sustained yield capacity (LTSYC). Therefore, additional analysis at a more refined level of VDDT modeling revealed that treatment strategy would need to change after the 15-year planning period for alternative B.

In order to sustain a non-declining even flow of ASQ volumes on suitable timberlands in alternative B, additional VDDT modeling revealed that the restoration strategy for decades 2 through 5 (years 16 through 50) would need to do the following: Increase treatment acreages in

closed canopy transition vegetation states in the ponderosa pine and dry mixed conifer PNVTs; and shift to using only low-severity prescribed fire as a maintenance tool which just thins the seedling/sapling sizes.

It makes sense to constrain the model this way by targeting more aggressive treatments in the closed-canopy states on suitable timberland acres in decades 2-5, because they will need the most restoration at that point in time. Decade 1 focuses partly on the community forest intermix and wildland urban interface areas, and acres already treated recently. Past/recent treatments and the Rodeo-Chediski/Wallow fires have created more open-canopy acres in these PNVTs. By decade 2 the untreated and more-dense portions of treated/burned areas left dense will need heavier and/or more frequent thinning to open them up. This focus on closed-canopy states is additional rationale for meeting the requirement to keep ASQ non-declining. Further discussions of the modeling methodology adjustments made are addressed in the “Environmental Consequences – LTSYC” and the “Relationship of Short-term Management to Long-term Productivity” sections, where explanations are more logical for the reader.

A few other model constraint scenarios were also studied for alternative B to maintain non-declining ASQ such as: (A) Target cutting all states more heavily/more often in decades 2-5, rather than just targeting the closed-canopy states. (B) Keeping some moderate-severity prescribed burn acres in decades 2-5. Both scenarios resulted in considerably more additional cutting acres needed annually beyond the minimum needed to meet the NFMA law, and possibly beyond the maximum treatment acreage objective set for Alternative B. They also shift the alternative B mix of cutting to burning from about a 50/50 mix, more toward the Alternative C mix of 70% cutting/30% burning which deviates too much from the alternative B management theme (i.e. we didn't need another alternative that ends up being so similar to an alternative that we've already analyzed).

Such a logical treatment sequence shift was not simulated in VDDT for Alternatives A and C. See additional modeling methodology explanations in the following two sections on LTSYC in this report.

Therefore given these various limitations of our VDDT modeling, the results are less reliable when enough decades have passed to expect the next re-entry onto the same acres. That timeframe varies by alternative, but generally occurs at about year 30. Thus, beyond the third decade, this specialist is skeptical about the VDDT model's reliability with respect to precisely measured outputs. General trends beyond the third decade are still assumed to be reasonably reliable outcomes at the programmatic level.

Assumptions

In the analysis for this resource, the following assumptions have been made:

- All four alternatives share the same desired conditions for the resources of the Apache-Sitgreaves NFs.
- Volumes of wood products potentially available from the Apache-Sitgreaves NFs are considered the by-products of treatments that move vegetation towards desired conditions. Management of all PNVTs would be needed to meet desired conditions.
- The revised plan is intended to be more strategic rather than prescriptive. The cuts and volumes presented in this report are estimates, and are not intended to dictate actual project-level treatment methods. According to the proposed plan objectives, it is assumed that work targets will be driven more by acres treated for restoration, rather than by maximized volume harvests.

- For estimating the consequences of alternatives at the programmatic forest plan level, the assumption has been made that the kinds of resource management activities allowed under the prescriptions will in fact occur to the extent necessary to achieve the goals and objectives of each alternative toward reaching the desired conditions. It is assumed the budgets and workforce needed to implement the specific activities will be available. However, the actual locations, design, and extent of such activities are generally not known at this time. That will be a site-specific (project-by-project) decision. Thus, the discussions here refer to the potential for consequences to occur, realizing that in many cases, these are only estimates.
- Southwestern regional FVS modeling is assumed to be representative of PNVTs on the Apache-Sitgreaves NFs. The ASNFs FIA plots were included in the Region-3 modeling data set.
- The VDDT model results are assumed to be reliable for the existing conditions and resulting conditions at year 15 to represent reasonable estimates for comparison of alternatives for this planning period. Longer-term VDDT results may not be as reliable, due to numerous model limitations.
- Appropriate cutting methods and other forms of treatments to be used are many, and vary by the site-specific objectives and existing condition. Decisions about which treatment methods to use, and on which acres, are project-level determinations that will be made by the project silviculturist after site-level field diagnosis, then working in cooperation with a full interdisciplinary team for NEPA analysis, line officer approval, and implementation.
- Harvest volumes from VDDT model runs reflect cutting intensities and acres which have incorporated the multiple-use objectives of each alternative. Model results include fairly realistic estimates of mixtures of approved vegetation management practices, for at least the first one to two decades. The particular silvicultural systems and cutting methods used in modeling for this analysis do not indicate which treatments would actually be applied at the project-level. A full range of all silvicultural treatment systems and methods would be available for project-level planning and decision making. See the standard silvicultural vegetation treatment options table provided in Appendix B1, which is also an appendix in the draft plan.
- USFS Region 3 FVS modeling indicates that to maintain the desired open-canopy structure on average site conditions, thinning re-entries onto the same forested acres sooner than 30 years after the prior cut typically may not enable tree growth rates adequate to provide an economical harvest. (Youtz and Vandendriesche, 2010, 2012; Weisz, et.al., 2012). Because site productivity is not directly used by the VDDT model for growth rates of structural state transitions, it is assumed that treatments using cuts on suitable timberlands would need to occur at about a 30-year re-entry cutting cycle, on average, to agree with harvest volumes calibrated into VDDT model treatments by regional FVS modeling.
- USFS Region 3 FVS modeling assumed that treatments using cuts to maintain piñon-juniper woodlands would need to occur at about a 50-year cutting cycle, on average due to slower growing species usually located on dryer and poorer sites than suitable timberlands. (Youtz and Vandendriesche, 2010; Weisz, et.al., 2012).
- “Industrial” (ASQ) timber 9+” DBH and pulp 5 to 9” DBH species would include: ponderosa pine, Douglas-fir, white fir, southwestern white pine, blue spruce, Engelmann spruce, corkbark fir, when cut from suitable lands. Although not included as “industrial” products under the provisions of the 1982 planning rule, the < 5” DBH sized materials, including tops and limbs, cut from these trees may be utilized as non-ASQ biomass.

- “Non-industrial” (non-ASQ) species include all sizes of: aspen, all junipers, all piñon pines, Chihuahua pine, all oaks cut from any lands; and any “industrial” species when cut from non-suitable timberlands. Wood cut from this category may be utilized as fuelwood and/or biomass.
- Markets would exist for all cut materials. At least 5 percent of cut materials may remain on the ground as broken logs and limbs, and/or debris left for soil stability, productivity, and wildlife needs. About 95 percent of the cut materials would be offered for removal from the site. The adjusted removal volumes constitute the amounts provided for the Economics Specialist Report analysis, as actual product volumes that could be made available for the local economy.
- Whatever future levels of increasing public and small market demands might be for fuelwood and incidental miscellaneous permits and small salvage sales of sawlogs/posts/poles/novelty woods/pulp volumes, they are assumed to be relatively similar for all alternatives. Current sale volumes (averaged TIM records for the 6 year period of 2005 to 2010) of timber salvage, fuelwood, poles and posts contribute to the local economy in measurable amounts and are itemized in the SocioEconomic specialist report (see Appendix D2 of this report, and the Plan set of documents).
- Increasing public and small market demands for fuelwood and small salvage sales of sawlogs, posts, poles, and/or pulp volumes would be similar under all alternatives. Incidental tree products like Christmas trees, wildings, seed cones, novelty woods, and other miscellaneous items would continue to be made available to meet public demand, and would be constant across all alternatives.
- In order to meet desired conditions, some form of management would be needed to meet desired conditions, regardless of whether lands are classified as suitable timberlands or not. Timber harvesting, salvage sales, and tree cutting may be conducted on non-suitable timberlands as needed to move those lands toward desired conditions for multiple resource objectives. The National Forest Management Act (NFMA) Section 6. (k) states (bold emphasis added):
 - *In developing land management plans pursuant to this Act, the Secretary shall identify lands within the management area which are not suited for timber production, considering physical, economic, and other pertinent factors to the extent feasible, as determined by the Secretary, and shall assure that, **except for salvage sales or sales necessitated to protect other multiple-use values, no timber harvesting shall occur on such lands for a period of 10 years. Lands once identified as unsuitable for timber production shall continue to be treated for reforestation purposes, particularly with regard to the protection of other multiple-use values.***
- Long-term sustained yield capacity (LTSYC) calculations include Region 3 FVS modeling that simulates treatments to move toward desired conditions. It uses data with the currently available tree growth and mortality rates, and incorporates maximum size limits of openings (in the group selection method), and minimum utilization standards with Region 3 average defects, into the determination of the annual sustained yield volume available to cut on a non-declining even-flow basis from suitable timberlands. These regional results are assumed adequate for the Apache-Sitgreaves NFs to establish a base sale offering schedule, using the projected future regulated inventory volume that could establish perpetual timber harvest.
- Numerous assumptions were made in the USFS Region 3 FVS Timber Calculations Modeling process to calibrate VDDT for cutting volumes and post-cut vegetation structural destination states. All methodology, assumptions, and rationale are documented

in the Region 3 Timber Calculations Working Group White Papers by Weisz et al. (2012). Some methodology is summarized in appendices of this report to assist the reader with understanding the basic procedural assumptions, silvicultural rationale, and modeling shortcomings (weaknesses/inconsistencies) of merging FVS model results into the VDDT modeling approaches used. For more detailed explanation of the VDDT model (including limitations), see the Vegetation Specialist Report.

- The allowable sale quantity (ASQ) cutting volume displayed by alternative would establish the sale offering schedule, as the average cut planned annually from suitable timberlands in the first decade.
- Because prescribed fire is now proposed in the action alternatives as a silvicultural thinning tool to intentionally kill trees of various sizes, it is addressed in this report. It is assumed that such use of fire would affect available amounts of green wood and dead wood harvest volumes and market values.
- Low severity fire (both planned and unplanned ignitions) on suitable timberlands would be used to reduce ground fuels and remove slash, and it would be used to maintain or move towards desired conditions (i.e., the age class distributions desired for uneven-aged structure, regulated forest, and sustained volume yields).
- The use of fire (e.g., moderate and high severity burns, planned and unplanned ignitions) as a thinning tool on suitable timberlands may occur when necessary to meet resource objectives. Moderate and/or high severity fire may not achieve the age class distributions desired for uneven-aged structure, regulated forest, and sustained volume yields.
- Due to fire's unpredictable nature under moderate burning conditions, it is assumed that moderate severity fire used for thinning trees over 5 inches diameter has far less control and assurance than tree cutting to achieve precise age class distributions for creating uneven-aged structure, regulated forest, and sustained volume yields.
- Due to fire's unpredictable nature under low burning conditions, it is assumed that low severity fire used for thinning reforestation (seedlings/sapling size class under 5" diameter) has less control and assurance than tree cutting to achieve precise tree stocking levels for creating uneven-aged structure, regulated forest, and sustained volume yields.
- Prescribed fire as a silvicultural tool will only be used under carefully prepared burn prescriptions developed by certified fire/fuels specialists working in cooperation with certified silviculturists; and fire would only be applied when the burning conditions fit the prescription to meet site-specific burn objectives toward meeting land management plan desired conditions.
- Because acres of topsoil loss and/or reduced site productivity due to recent uncharacteristic severe wildfire and subsequent erosion are not yet known, all lands classified as suitable for timber production remain in the suitable timberland base for this analysis.
- Currently deforested acres would not need thinning during this planning period.
- Funding will be available to collect needed amounts of green tree seed and to plant acres proposed for artificial reforestation.

Revision Topics Addressed in this Analysis

This report analyzes consequences of the alternatives for the listed revision topic and issues, using the following indicators and units of measure for a meaningful basis to compare alternatives:

Revision Topic 3: Community-Forest Interaction

Issue: Contribution to Local Communities

1. Suitable Timberlands and Allowable Sale Quantity
 - Lands suitable for timber harvest (in Acres)
 - Allowable sale quantity (ASQ) of timber (in CCF/Year)
 - Long-term sustained yield capacity (LTSYC, in CCF/Acre)

2. Potentially Available Wood Products From Suitable and Non-suitable Timberlands
 - Timber cut volumes (in CCF)
 - Fuelwood cut volumes (in CCF)
 - Biomass cut volumes (both in Tons and converted to CCF)
 - Total woody products (all in CCF)

Summary of Alternatives

A summary of alternatives, including the key differences among alternatives, is outlined in the Draft Environmental Impact Statement.

Affected Environment

This section describes existing conditions. Of the total national forest land area managed by the Apache-Sitgreaves, approximately 945,753 acres (47%) are classified as vegetation PNVTs in which coniferous tree species should dominate. Riparian hardwood deciduous forest, woody shrub types, and most grassland PNVTs are addressed in the Riparian and Vegetation Specialist reports. The great basin grassland is only addressed in this report because so many acres are encroached with trees that need mechanical reduction (fire alone will not accomplish this objective).

Table 1. PNVTs and Acreages addressed in this report.

PNVT	Total GIS Acres	Approximate Percent of Coniferous Forest PNVT Acres
Ponderos pine (about 30% has Gambel oak as a co-dominant species)	602,206	63%
Dry Mixed Conifer (frequent fire type)	147,885	16 %
Wet Mixed Conifer with Aspen (infrequent fire type)	177,995	19 %
Mixed Spruce –Fir (low elevation mixed species type)	17,667	2 %
Pinyon-Juniper (Grassland savanna and persistent woodland)	222,166	NA
Great Basin grassland	185,523	NA

Past forest growth and mortality, previous active and passive management, and disturbance patterns have produced the current forest tree species composition, sizes, densities, and conditions, which affect the species and volumes of wood and other tree products available for cutting treatments now and in the future.

Across the Apache-Sitgreaves NFs, the annual net and gross forest growth have has far exceeded cutting levels. Rogers (2003) and O’Brien (2002) both illustrate this condition with Forest Inventory Analysis (FIA) plots. Rogers shows that annual gross growth in Apache-Sigreaves NFs Forest Products Specialist Report - Apache-Sitgreaves NFs Plan Revision DEIS

ponderosa pine non-reserved lands has been as much as 12 times the annual mortality. For local Douglas-fir stands, growth has exceeded mortality by at least 8 times. O'Brien shows that mortality and cutting levels combined are far below gross growth rates for trees inventoried in Arizona national forests non-reserved timberlands. In other words, cutting rates have been far less than net growth rates (please see Glossary).

In the past 30 years, an average of about 720 cubic feet (CF) of volume per acre has generally been added as surplus net growth, in addition to the desired sustainable volume. This surplus needs to be removed each entry to maintain desired conditions. Every three decades that pass without treatment, a backlog of overgrowth continues to be added. On the Apache-Sitgreaves NFs, several acres of suitable forest land have not been thinned in over 40 to 50 years, leaving a current surplus of over 1,000 CF on some acres. Acres which have this backlog suffer from conditions which contribute to the departure from desired conditions. These areas are at risk of accelerated stand mortality due to intense tree competition, weakened tree vigor, disease intensification and spread, epidemic insect attacks, and/or uncharacteristic wildfire.

This forest growth has outpaced all mortality (including natural mortality and by cutting) for many years, resulting in an extreme imbalance, which is not sustainable. Normal disturbance regimes which used to act as natural thinning agents have been altered (primarily characteristic wildfire), giving rise to overgrown conditions. Abnormal disturbances (like uncharacteristic wildfire and unprecedented insect/disease outbreaks) have produced undesirable stand-replacement conditions across large areas (US Forest Service, Dec.2008-ESR; Lynch et al., 2010) Noticeable (non-fire) losses of large/old trees have been seen recently in all forested PNVTs (Boehning, 1982-2012 professional field observations validated with FIA plot data, see Appendix H), especially due to competition in overgrown stands. Overgrowth increases competition among trees for nutrients, water, and growing space, which in turn reduces individual tree and forest stand growth, vigor, and the ability to endure bark beetle attacks and drought years (Korb et.al., 2012; Covington et al. 1997; Friederici, 2006b).

Two very different existing condition categories now occur across the forested PNVTs of the Apache-Sitgreaves NFs (see below and table 2):

- **Forested/Overgrown¹** - Approximately 71 percent of the current forested types have tree stocking and growth levels which require some degree of tree thinning to restore and/or maintain desired conditions. Without additional severe disturbances accelerating immediate and complete tree mortality, these areas can contribute industrial cutting volumes in the first decade and beyond.
- **Deforested/Early Development** - Conversely, an average of about 29 percent of the current forest type vegetation structural states are now temporarily in deforested² states (22 percent) or early developmental³ states (7 percent) that require reforestation and growth for restoration to desired conditions. This condition is primarily attributed to the 2011 Wallow Fire, the 2002 Rodeo-Chediski Fire, and other fires that caused tree mortality that exceeded or eliminated net growth. These areas cannot provide industrial cutting volumes in the next one to three decades.

Footnotes:

¹ "Forested/Overgrown" acres are all remaining structural states not identified as deforested/early development acres.

² "Deforested acres" are those in excess of desired condition percentages for the following PNVTs: Ponderosa pine and dry mixed conifer structural states A and N; wet mixed conifer and spruce-fir structural states A and K.

³ “Early developmental” acres are those in the seedling/sapling (<5” diameter) structural states, which for some forest type PNVTs also include the small size class (5-9” diameter).

Table 2. Acres and percent by forest type PNVT in forested/overgrown or deforested/early development condition

Forest Type (PNVT)	Forested/Overgrown Acres (Percent)	Deforested/Early Development Acres (Percent)
Ponderosa Pine	475,743 (79%)	126,463 (21%)
Dry Mixed Conifer	100,562 (68%)	47,323 (32%)
Wet Mixed Conifer	89,005 (50%)	88,998 (50%)
Spruce-Fir	8,127 (46%)	9,540 (54%)
TOTAL	673,437 (71%)	272,316 (29%)

On the forested/overgrown lands, net growth is expected to outpace natural (non-fire) mortality, such that regular thinning (prescribed cutting and/or burning) will be necessary to reduce overgrowth, develop desired uneven-aged forest structure, and/or prevent growth stagnation and movement away from desired conditions. Where moderate-severity fire has occurred on these lands, natural mortality levels are expected to continue to remain high for approximately the next six years, due to fire-related tree stress, sudden exposure to weather extremes, weakened roots, greater exposure to lightning and/or prevailing winds, and greater susceptibility to insect/disease attack. Once surviving trees have stabilized, they are expected to again need thinning for maintenance of desired forested conditions.

For now, deforested lands remain in the suitable timberland base; but without natural regeneration or artificial reforestation, they may not return to timber productivity for many years. Limestone soils are common across the Sitgreaves NF, while basalt soils are more common on the Apache NF (ASNFs TES, Laing et al.,1987). Natural conifer regeneration rates and numbers on limestone soils across northern Arizona are known to survive better than on basalt soils (Puhlick et al. 2012, Richardson 2012; Boehning, 1982-2012). Natural aspen root-sucker regeneration occurs quickly after fire events where aspen roots exist in good health. Natural aspen seedling establishment can also occur in burned areas where aspen roots do not pre-exist. Successful establishment and long-term survival of aspen regeneration will mostly depend on the amount of domestic and wild ungulate herbivory damage that occurs within the first 2-6 years post-burn (Shepperd and Fairweather, 1994; Rogers, 2008 and 2011; Lynch et al, 2010).

A large portion of the natural regeneration acres, especially in the wet mixed conifer and spruce-fir PNVTs, now have prolific aspen root-sucker regeneration occurring in response to fire stimulation. Those young, aspen-dominated acres will temporarily provide no industrial harvest volumes until suitable conifer regeneration becomes established and grows to commercial size. New aspen stands which survive to maturity may begin to provide some fuelwood or biomass products in about 20+ years.

However, a portion of those acres in need of reforestation may not return to a productive forested condition. A body of research in the southwestern ponderosa pine forest type has shown that, on average, approximately 50% of severely burned acres tend to convert to grass/forb/shrub/rock lands that can take a very long time to return to forest, possibly centuries, if ever (Roccaforte et al., 2012; ERI, 2011; Savage and Mast, 2005; Strom and Fule, 2007). Artificial reforestation is needed to reduce that percentage (Higgins, 2008; ERI, 2012b). This tendency is noted for mixed conifer by Jones, 1974. Climax spruce-fir and upper elevation (wet) mixed conifer forests have been documented as also undergoing very long-term type conversions to grass/forb/shrubland following severe fire (Alexander 1974).

Thus, the deforested lands (approximately 203,378 acres) can be further divided into three categories:

- Lands which can be expected to successfully regenerate native tree species naturally (estimated at approximately 54 percent or 110,629 acres) at low management cost. This is approximately 12 percent of all forest type PNVT acres.
- Lands which would need artificial tree planting to restore forest cover (estimated at approximately 19 percent or 37,695 acres) at high management cost. This is approximately 4 percent of all forest type PNVT acre.
- Lands that are likely to undergo site conversions to long-term grass/forb/shrub/rock cover rather than return to tree cover (estimated at approximately 27 percent or 55,054 acres) at the expense of lost forest/timber production acres. This is approximately 6 percent of all forest type PNVT acres. Artificial tree planting can help mitigate this condition by accelerating post-fire succession back to desired tree cover.

Newly established conifer stands (both natural and planted) will require regular thinning entries once they are past the seedling state, to promote tree vigor and facilitate maximum growth for faster return to desired forest conditions.

Lands Tentatively Suitable for Timber Production

Timber production is defined as the purposeful growing, tending, harvesting, and regeneration of regulated crops of trees for cutting into logs, bolts, or other round sections for industrial or consumer use. Timber production does not include fuelwood or products harvested from unsuitable lands. Suitable timberland does not dictate tree cutting. It means that all cutting treatments done on suitable lands would be subject to the annual ASQ volume as a control to prevent over-cutting.

Lands are identified as suitable or not suitable for timber production (referred to as suitable and non-suitable timberlands) during the plan revision process. Appendix A2 details the steps used in the suitability determination. The first step of the suitability determination is to identify those lands that are tentatively suitable for timber production.

Table 3 displays the criteria used to identify lands as tentatively suitable timberlands. The Apache-Sitgreaves NFs have approximately 808,368 acres considered tentatively suitable. Suitable timberland does not dictate tree cutting. It means that all cutting treatments done on suitable lands would be limited by the ASQ volume (see following section on Allowable Sale Quantity).

Table 3. Criteria used to determine tentatively suitable timberlands in all alternatives

Criteria	Acres	Total Acres
TOTAL Apache-Sitgreaves NFs		2,110,196
Non-NFS Land	94,844	
Total NFS Lands		2,015,352
Non-forest Lands		1,039,258
Areas not defined as forest land ¹	4,250	
quarry, urban/agriculture, water		
Grasslands	344,033	
Great Basin, montane/subalpine, semi-desert		
Woodlands	617,094	
Madrean pine-oak, piñon-Juniper		
Interior chaparral	55,981	
Wetland/cienega	17,900	
Forested lands withdrawn from timber production²		87,190
Designated Wilderness	20,628	
Bear Wallow, Escudilla, Mount Baldy		
Blue Range Primitive Area	43,258	
Research Natural Area	219	
Eligible or suitable wild and scenic river corridors or areas classified as wild	23,085	
Irreversible resource damage likely		23,952
Unsuited/unstable soils (sensitive and unstable)	23,952	
Inadequate restocking		56,584
Low reforestation potential based on soil properties	56,584	
LANDS Tentatively Suitable for Timber Production		808,368³
¹ Forest land is defined as having greater than 10 percent overstory canopy cover at stand maturity ² Some categories overlap areas already withdrawn in non-forest lands ³ The tentatively suitable lands in alternative A equal 807,289 acres. There are more acres in research natural area (1,882 acres)		

The above table reflects the same step 1 common to all action alternatives.

Acres of “unsuited/unstable soils” and “low reforestation potential” were derived from the Apache-Sitgreaves NFs Terrestrial Ecosystem Survey as “inadequate for restocking”. They were not modified after the 2011 Wallow Fire, because the Forest Soil Scientist believes it is too early (in 2012) to determine accurate estimates of soil productivity losses due to fire consumption of the organic layers and/or subsequent erosion of topsoil. The fire area soils, watersheds, and ground cover have not yet stabilized post-burn. This is a site-specific determination that will need to be made at the project-level and based on soils monitoring over time. Any estimates made of possible site conversion from forested PNVTs to grass/rock/shrubland in the Forest Products Specialist report for this analysis are purely estimates based on a search of relevant literature, which will also require on-site monitoring for local validation.

Adjustments to the suitable timberland acreage within the Wallow Fire and other high-severity fires may be appropriate in the next 10 years during the scheduled review and update of the forest suitability classification process.

Allowable Sale Quantity (ASQ)

The allowable sale quantity volume control concept enacted by law (National Forest Management Act of 1976) was intended to prevent excessive tree losses due to over-cutting beyond sustainable forest levels on suitable timberlands. The ASQ is the quantity of timber that may be sold from suitable timberland within the Apache-Sitgreaves NFs for a time period specified by the plan. ASQ volume is expressed as the average annual allowable sale quantity. For timber resource planning purposes, the allowable sale quantity applies to each decade during the planning horizon period and includes only chargeable volume. ASQ volume does not include fuelwood or other nonindustrial wood.

ASQ volume estimates only include chargeable volumes¹ of industrial wood². The ASQ calculation includes estimated green tree cutting volumes from the VDDT model which vary by alternative; extra small sales and permits sold for live and/or dead poles; and miscellaneous salvage timber. Small sales, permits, and miscellaneous salvage are considered to be a constant addition across all alternatives.

ASQ and a timber base sale schedule were published with the original 1987 forest plan, along with the timber suitability determination. They were specific and prescriptive. The original ASQ volume of 119 million board feet (MMBF) was subsequently reduced to an interim ASQ of 99 MMBF (198,000 CCF³) of sawtimber per year in response to a plan settlement agreement and through forest plan amendment one. That ASQ and related sale offering schedule were never modified again after plan amendment number 4, issued in 1991. During that reanalysis of the ASQ for the settlement agreement, the Forest Silviculturist documented additional ecosystem management concerns beyond the ASQ debate at that time (Shafer, 1993). His concerns are still valid today.

The ASQ volume for alternative A has been recalculated in this analysis for consistency across all alternatives, and is based on current vegetation conditions.

During the planning period (next 15 years) on-going monitoring would evaluate cutting levels compared to the ASQ. According to law and policy, the suitable timberland classification would be updated as conditions and/or management emphasis/strategies change.

¹ Chargeable volume of industrial wood is from tree species that are saleable as sawtimber logs, pulpwood bolts, poles, or other roundwood sections (excluding fuelwood) based on regional utilization standards and cut from suitable timberlands.

² Industrial timber species (5 inches and greater) include ponderosa pine, Douglas-fir, white fir, southwestern white pine, blue spruce, Engelmann spruce, and corkbark fir.

³ CCF = one hundred cubic feet.


Long-Term Sustained-Yield Capacity (LTSYC)

Long-term sustained yield (LTSY) is the calculated annual volume of wood per acre that can be harvested from suitable timberlands, which does not exceed annual net growth volume per acre after desired conditions have been met for multiple resource objectives. LTSY multiplied by the total suitable timberland acreage derives the long-term sustained-yield capacity (LTSYC). This concept is one means of measuring forest sustainability, consistent with ecological desired conditions as well as sustainable harvest volumes for society.

On a per-acre basis, the true problem today on many forested acres in fact is not over-cutting, but rather a backlog of overgrowth that has consistently been undercut annually. In a regulated forest, annual cut equals annual net growth, such that the entire forest never becomes completely overgrown or stagnant for very long. Historically, regular fire intervals were one of nature's methods for removing excess growth. On suitable timberlands, management actions like tree cutting and fire-use may be used to achieve the same purpose.

From Youtz and Vandendriesche's September 2012 regional work, an average of 24 cubic feet (CF) per acre per year is used to represent the net growth across all acres of forested PNVTs on the ASNFs that are suitable for timber production. This net per-acre average accounts for annual gross growth after annual mortality is deducted, based on regional average growth and mortality rates calibrated in the Forest Vegetation Simulator (FVS) model. The following simplified figure (not drawn to any scale) conceptually demonstrates this problem.

Figure 1. Conceptual diagram of ideal cutting level for a sustainable forest and sustainable harvests (i.e. "regulated forest")



Because uneven-aged forest structure is an important part of the ecological desired conditions, Youtz and Vandendriesche (2012) emphasized it in their modeling to determine regional average CF per-acre-per-year LTSY values for timber-producing PNVTs. Their LTSYC analyses which used the 6-age group 30-yr. cutting cycle prescription for ponderosa pine, dry and wet mixed conifer is the only one of their silviculture scenarios that was used by the Southwestern Region to also model FVS Group Selection cuts for calibrating uneven-aged cutting volumes in the VDDT

model. Therefore, the values used in this report’s analysis are taken from their 30-year cutting cycle scenarios, and it is assumed here that a return cutting entry of about 30 years is needed on average under the Group Selection system for uneven-aged management to maintain a regulated sustainable forest for desired conditions (Youtz and Vandendriesche, 2010).

A weighted average long-term sustained yield (LTSY) has been calculated as 24 cubic feet (CF) per acre per year for all Apache-Sitgreaves NFs forested PNVTs in suitable timberlands based on current regional data (see Youtz and Vandendriesche, 2012 in Appendix F). This is slightly higher than the 20 CF per acre per year capacity identified in the 1987 plan. This preliminary LTSY was one criterion used in the determination of tentatively suitable timberlands, based on local soil productivity ratings (see Appendix A2).

For simplification of analysis, the long-term sustained yield of 24 cubic feet per acre per year used is a rounded, weighted average value for all suitable timberlands, using the regional model run results for each PNVT, based on the proportional acres of each forested PNVT present on the Apache–Sitgreaves NFs suitable land base. Only the Southwestern Region’s high-site model run for the ponderosa pine/grass type was used in this calculation, because soils not capable of producing at least 20 cubic feet/acre/year (approximately site index of 70 or greater) were eliminated from the tentatively suitable land base with the Apache-Sitgreaves NFs’ soils assessment (see Appendix A2). Because acres of suitable timberland vary by PNVT, a weighted average was used to verify the correct average to be used for all analyses of all PNVTs combined. Table 4 below shows how this average was derived mathematically.

Table 4. Average LTSY calculation for all suitable timberland PNVTs on the Apache-Sitgreaves NFs by alternative

PNVT	Suitable Acres in Alternative A ³	LTSY in cubic feet/acre/year ¹	Multiplication Product
Ponderosa pine ²	503,412	23.6	11,880,523
Dry Mixed Conifer	108,208	22.9	2,477,963
Wet Mixed Conifer	148,072	24.7	3,657,378
Spruce-Fir	5,180	0	0
Totals	764,872	71.2	18,015,864
Alternative A Weighted Average: 18,015,864 / 764,872 = 23.6, rounded to 24 cubic feet/acre/year			
PNVT	Suitable Acres in Alternative B ³	LTSY in cubic feet/acre/year ¹	Multiplication Product
Ponderosa pine ²	445,440	23.6	10,512,384
Dry Mixed Conifer	65,086	22.9	1,490,469
Wet Mixed Conifer	86,217	24.7	2,129,560
Spruce-Fir	0	0	0
Totals	596,743	71.2	14,132,413
Alternative B Weighted Average: 14,132,413 / 596,743 = 23.7, rounded to 24 cubic feet/acre/year			
PNVT	Suitable Acres in Alternative C ³	LTSY in cubic feet/acre/year ¹	Multiplication Product
Ponderosa pine ²	451,179	23.6	10,647,824
Dry Mixed Conifer	65,778	22.9	1,506,316
Wet Mixed Conifer	87,789	24.7	2,168,388
Spruce-Fir	0	0	0
Totals	604,746	71.2	14,322,528
Alternative C Weighted Average: 14,322,528 / 604,746 = 23.7, rounded to 24 cubic feet/acre/year			
¹ From Youtz and Vandendriesche, 2012.			
² Only the regional ponderosa pine/grass type high site index LTSY model result was used.			

³ See Forest Products Specialist Report Appendix A-2 for additional information.

Because this net growth average of 24 cubic feet per acre per year does not vary by alternative, it was used in all LYSTC calculations for all alternatives. This LTSYC level becomes a baseline for comparison of estimated short-term wood product cutting levels by alternative, as presented later in this report.

LTSY multiplied by the total suitable timberland acreage derives the long-term sustained-yield capacity (LTSYC). This concept is one means of measuring forest sustainability, consistent with ecological desired conditions. Ideally, ASQ volume should equal or fall just short of the LTSYC once desired conditions are met. After desired conditions are achieved, management on suitable timberlands would need to be consistent with the LTSYC level (no cutting departure above the LTSYC); annual cutting levels would not exceed annual net growth rates. On suitable timberland acres, the LTSYC is a way to further incorporate the social and economic desired condition of providing a long-term, dependable source of wood products, while maintaining desired multiple-use objectives.

So after every 30 years of net growth 720 CF of volume is added which needs to be removed each entry. Every entry that passes by untreated continues to add to a backlog of overgrowth. On the ASNFs, several acres of forestland have not been thinned in over 40-50 years (personal knowledge and district records), leaving an excess backlog of over 1,000 CF on some acres. Acres which have this backlog suffer from conditions that contribute to an ecological imbalance which must be corrected. If accelerated cutting is not done to correct that imbalance, then nature will correct it by means of extreme stand mortality due to weakened tree vigor, disease intensification and spread, epidemic insect attacks, and/or uncharacteristic wildfire. This understanding of overgrowth backlog applies to both suitable and non-suitable timberlands, although LTSYC and forest regulation are only formally applied to suitable timberlands.

Moreover, to comply with legal requirements of the National Forest Management Act (NFMA) and Multiple Use Sustained Yield Act (MUSYA), long-term sustained yield also means that ASQ volumes harvested from suitable timberlands cannot decline from one decade to the next. More explanation of the non-declining ASQ concept is presented later in the Environmental Effects section of this report.

Wood and Tree Products Availability

Wood products can be provided from both suitable and non-suitable timberlands. The most common wood products (e.g., industrial⁴ and non-industrial⁵, live and dead wood) on the Apache-Sitgreaves NFs include sales and permits for:

- Ponderosa pine, Douglas-fir, southwestern white pine, white fir, blue spruce, Engelmann spruce, and corkbark fir

⁴ The less than 5 inch diameter size materials, including tops and limbs from timber species may be utilized as non-ASQ biomass.

⁵ Non-industrial (non-ASQ) species include aspen, junipers, piñon pines, Chihuahua pine, oaks, and any industrial species cut from non-suitable timberlands. Wood cut as non-industrial may be used as fuelwood and/or biomass.

- Sawtimber and house logs (9 inches or greater diameter)
- Pulpwood or Roundwood (5 to 8.9 inch diameter)
- Poles, posts, vigas, latillas, rails
- Laminated beams
- Paneling and trim moulding
- Fuelwood
- Biomass (chips)
- Furniture
- Piñon pine and all juniper species
 - Poles, posts, vigas, latillas, rails
 - Fuelwood
 - Biomass
 - Furniture and novelty wood
- Aspen, Gambel-oak, and other oak species
 - Fuelwood
 - Furniture and novelty wood
 - Poles, posts, rails
 - Interior paneling
 - Evaporative cooler pad excelsior
 - Livestock bedding

Major non-wood tree products of various species have also been made available for the following other uses: Christmas trees, live seedling/wilding transplants, green seed cones for nurseries, green holiday decorative boughs and wreaths, and piñon nuts. Ceremonial wood, tee-pee poles, and other native American items have also been provided only to tribal members by special request.

Within the life of the 1987 plan, annual harvest volumes have varied from 5,000 to 100,000 thousand board feet (MBF) with annual treatment acreages ranging from 2,500 to 30,000 with an average of 9,400 acres (US Forest Service, Dec. 2008-CER). Harvest volumes have stayed under the original 1987 plan ASQ of 119,000 MBF (or 119 MMBF). The 100,000 MBF maximum harvest level achieved converts to 200,000 CCF. Annual acreages treated under the White Mountain Stewardship project (2004 to present) are slightly more than the prior average at approximately 12,182 acres used to represent the current trend under the existing land management plan (Drury, February 2012).

A brief summary of vegetation management pertaining to past cutting practices follows.

With the implementation of the 1987 plan, vegetation management of forest types emphasized even-aged cutting methods: seed cuts, final overstory removals, intermediate thinnings, and a few clearcuts/seedling plantations. Sanitation/salvage cuts have also been used. The concept of “integrated stand management” promoted the policy to leave numerous timber stands untreated as “deferrals” intermixed between treated stands across all projects for creating diversity, and leaving dense key wildlife habitat and un-thinned old growth. Most vegetation management was accomplished through timber sales that focused on cutting trees over 9 inch diameter (or over 12 inch diameter in some cases). Multi-product sales (sales which offer both sawtimber and pulp

sizes) targeted trees in the 5- to 9-inch class as well as larger trees, but the lack of markets for the smaller size resulted in many projects not being completed (defaulted sales).

When the 1987 plan was amended in 1996 to specifically address management for Mexican spotted owl, Northern goshawk, and old growth, it initiated direction to emphasize uneven-aged cutting methods (e.g., group selection, individual tree selection). But implementation was met with varying degrees of success, due to various factors. Only thinning of marketable size trees (usually 9 inch diameter and larger) was successfully implemented. Thus many acres became further over-grown with trees under 9 inch diameter which can act as understory ladder fuels.

Uncharacteristic wildfires in the early 2000s highlighted the need for fuels reduction projects. Treatment of all vegetation types, regardless of timberland suitability, became a priority near communities, private lands, and developed recreation areas. The treatment emphasis on removing understory ladder fuels led to the use of a diameter cap (an upper cutting size limit) as the way to focus on removing the over-abundant, small diameter trees.

The Healthy Forests Restoration Act of 2003 fostered the development of community wildfire protection plans that incorporated programmatic and widespread use of diameter caps (limiting cuts to smaller diameter trees) (Logan Simpson Design, 2004a, 2004b, 2005). Meanwhile, the Stewardship End Result Contracting Projects statute (16 U.S.C 2104) opened a new era of cutting small trees to reduce wildfire hazards under 10-year long stewardship service contracts. On the Apache-Sitgreaves NFs, this meant that most wood volume was no longer removed through timber sales. Service contractors are being paid in the White Mountain Stewardship Project to cut and remove the vast majority of resulting wood volume; a government strategy used to reduce fuels near communities until local small-tree markets are established and small tree wood value offsets treatment cost (ERI, 2007).

Major local markets cannot afford the current gasoline and transportation costs to move raw cut materials much over a maximum of 70 miles to their processing plants (Drury, 2011). Raw wood values are so low that very little of the cut volume is sold to purchasers (ERI, 2007). Where haul distances are too far or transportation costs too high, a portion of the cut volume is left on-forest as waste material that must be disposed of at additional cost to the government (Hunt, 2012; Sitko and Hurteau, 2010; Drury, 2011).

Many of the most easily operable, accessible, and shortest haul distance acres have already been thinned with numerous fuels reduction projects. Most wildland urban interface (WUI) areas have already been treated in the past 10 years, and /or are still under contract with White Mountain Stewardship Project to be treated under existing approved NEPA decisions. Many of the acres left to be treated during this next 15 year planning period are more difficult ground operationally, and/or have longer haul distances to markets (Drury, 2011).

Black, fire-charred wood has been rejected by some local markets, like pulpwood for paper manufacturing and the wood heating stove pellet industry. Even though other markets like commercial firewood, shipping pallets, lumber sawmills, and bio-electric energy plants will take this material, the majority of fire-salvaged logs and poles have been sold at minimum base rates (\$1.00 per CCF for all species). High haul distance costs continue to be the deciding factor for the demand and value of this raw material (Drury, May 2012). Fire-charred wood could tend to stay at a very low market value, even if green wood values improve. The SocioEconomic Specialist Report summarizes more local/regional wood product economic, market, and workforce conditions.

Using even-aged cutting methods and diameter caps has resulted in a longer timeframe to produce uneven-aged forests and woodlands. In areas where even-aged management or removal of small Forest Products Specialist Report - Apache-Sitgreaves NFs Plan Revision DEIS

diameter trees have occurred, the next treatment (cutting entry) may be limited to large trees (16 inch and greater diameter) (Triepke et al., 2011). The effects of stand-replacement fire result in essentially even-aged regeneration areas for at least 60 years after new tree establishment. In all cases, subsequent entries could provide variable harvest volumes and product types; while conversion from even-aged to uneven-aged structure proceeds over time.

Environmental Consequences of Alternatives

The land management plan provides a programmatic framework that guides site-specific actions but does not authorize, fund, or carryout any project or activity. Because the land management plan does not authorize or mandate any site-specific projects or activities (including ground-disturbing actions) there can be no direct effects. However, there may be implications, or longer term environmental consequences, of managing the forests under this programmatic framework.

All alternatives would propose various mixtures of three basic vegetation treatments during the 15-year planning period to move toward desired conditions. These include:

- Tree cutting on some forested/overgrown areas, followed by low severity fire to reduce ground fuels and/or thinning-created slash.
- Moderate and/or high severity burning to thin other forested/overgrown areas and reduce ground fuels
- Tree planting on some deforested areas.

All three forms of treatment indirectly impact the amount and availability of sustainable wood products. The number of total annual cutting and burning treatment acreages by alternative (regardless of timber suitability classification) were analyzed in the table below.

Table 5. Annual cutting and burning treatment acres for all PNVTs, suitable and non-suitable timberlands, by alternative

Alternative	High Treatment Acres		Low Treatment Acres		Average Treatment Acres		
	Cutting Treatment	Burning Treatment	Cutting Treatment	Burning Treatment	Cutting Treatment	Burning Treatment	Total Cutting and Burning
A	-	-	-	-	12,182	6,844	19,026
B	30,327	43,771	8,852	14,087	19,591	28,930	48,521
C	42,651	22,586	5,342	3,124	23,997	12,857	36,854
D	25,440	78,772	6,465	19,079	15,954	48,927	64,881

Under any alternative, regular tree cutting on suitable timberlands would be necessary to move the forests toward an uneven-aged (regulated) balance of age classes, and then maintain this condition once it is achieved. It also is very costly. Increasing use of prescribed fire to intentionally kill excess trees is expected to become necessary for treating additional acres as a more cost-effective thinning tool.

The following table displays the relative amounts of prescribed fire severity modeled for each alternative, by PNVT. While these amounts do not in any way dictate that these levels would be used on any given project, they do reveal the general concept for emphasizing more fire as part of the overall restoration activity focus, especially during the planning period. Sources of this data are Appendices E1 and E2 of this report.

Table 6. Annual prescribed burning average objective acres modeled by forest type PNVT (suitable and non-suitable timberlands) and by burn severity, for each alternative

Forest PNVT	Alternative A Burn Severity		Alternative B Burn Severity		Alternative C Burn Severity		Alternative D Burn Severity	
	Low	Moderate &/or High	Low	Moderate &/or High	Low	Moderate &/or High	Low	Moderate &/or High
Ponderosa Pine	2,836	316	2,205	4,095	1,965	3,649	4,438	8,242
Dry Mixed Conifer	720	80	396	1,268	363	1,162	805	2,576
Wet Mixed Conifer	855	1,047	633	1,268	575	1,150	1,273	2,551
Spruce-Fir	90	10	115	231	164	329	185	370
Subtotals :	4,501	1,453	3,349	6,862	3,067	6,290	6,701	13,739
Totals:	5,954		10,211		9,357		20,440	

In addition, attempting to use fire, and/or natural tree thinning agents like insects/diseases, instead of tree cutting on suitable timberlands would be risky because there is no assurance that such methods would create the regulated forest needed for non-declining even-flow of harvest yields, neither in the short-term nor in the long-term. Much debate exists in forestry science on this matter. A study in southwestern ponderosa pine forest indicates that restoration of that ecosystem might be achievable solely with high and/or moderate severity prescribed fire (Fule et al., 2004). As of yet, no conclusive textbooks have been written precisely on how to achieve a regulated forest with non-declining even-flow harvest levels by using fire as the primary or sole silvicultural thinning tool. Much literature makes a good case for using prescribed fire after prescribed cutting practices on the same acres (Arno and Fiedler, 2005; Stephens et al., 2009; Schwilk et al., 2009; Cram et al., 2006; Korb et.al., 2012; Friederici, 2006a). For more information on use of prescribed burning and fire-use as the only tree-thinning method, see Appendix E2, and the Fire Specialist Report.

Suitable Timberlands

Timberland suitability was determined for each alternative, based on the detailed description of the suitable timberland analysis process and criteria found in Appendix A2. Acres of suitable timberlands would vary by alternative because the boundaries and acres of management areas differ between alternatives. The next table shows criteria used to identify lands suitable or not suitable for timber production, based on area objectives which limit industrial timber management activities.

Table 7. Lands suitable and not suitable for timber production, by area

Management Area	Timber Production	
	Suitable	Not Suitable
General Forest	X	
Community-Forest Intermix	X	
High Use Developed Recreation Area		X
Energy Corridor		X
Wild Horse Territory	X	
Wildlife Quiet Area	X	
Natural Landscape		X

Management Area	Timber Production	
	Suitable	Not Suitable
Recommended Research Natural Area		X
Research Natural Area		X
Primitive Area		X
Recommended Wilderness		X
Wilderness		X
Other Areas		
Areas with soil condition rating of unsuited/inherently unstable		X
Lands not cost efficient in meeting timber production objectives		X
Grasslands, woodlands, interior chaparral, and riparian forests		X
Communication sites		X
Developed recreation and administrative sites		X
Eligible or suitable wild and scenic river		X
MSO protected lands		X

Timberland suitability was determined for each alternative. The original 1987 plan suitability classification did not clearly follow the same criteria and classification as outlined below. For a basis of comparison across all alternatives, **alternative A** was recalculated using the same process and PNVT concept as the **action alternatives**. In addition to these criteria, other considerations (e.g., timber production cost efficiency) were used to further eliminate acres from suitability classification (see Appendices A3 and A3Roads.)

Results of the suitability determinations are provided in the following table. **Alternative A** would have the most acres suitable for timber production, followed by **alternatives C and B**. **Alternative D** would have no suitable acres.

Table 8. Lands suitable and not suitable for timber production by alternative

Classification	Acres			
	Alternative A	Alternative B	Alternative C	Alternative D
Lands Tentatively Suitable for Timber Production	807,289	808,368		
Lands where Management Area Direction Precludes Timber Production	12,258	65,497	27,321	145,118
Lands where Management Objectives Limit Timber Harvest	30,159	76,537	91,067	663,250
Lands that are Not Economically Cost Efficient	0	69,590	85,234	N/A
Lands Not Appropriate for Timber Production	42,417	211,624	203,622	808,368
Lands Suitable for Timber Production	764,872	596,744	604,746	0
Lands Not Suitable for Timber Production	1,250,480	1,418,608	1,410,606	2,015,352

Suitable timberland maps for **alternatives A, B, and C** are shown in Appendix A4. **Alternative D** would have no suitable timberlands.

Steep slopes (over 40%) were included in the original plan as suitable timberlands for cable harvest, with no suitability update conducted after Plan Amendment #6 adopted the Mexican Spotted Owl (MSO) Recovery Plan Protected Habitat restrictions for steep slopes and MSO Protected Activity Centers. **Alternatives B and C** do not include any steep slopes, or MSO Protected Habitat acres in the dry and wet mixed conifer, spruce-fir, and pine-oak (30 percent of the ponderosa pine) types. Today these acres are neither economically feasible, nor possible when managed under the MSO Recovery Plan requirements. Thus for all even-aged treatments appropriately used in the spruce-fir type, the NFMA requirement to select the rotation age at culmination of mean annual increment for sustained yield would not apply.

Alternative A would provide the most acres of suitable timberlands; while **alternative C** would provide slightly more acres of suitable timberlands than **alternative B**. Steep slopes (over 40 percent) were included in the 1987 plan (**alternative A**) as suitable timberlands for cable harvest.

Alternatives B and C would not include any steep slopes because these areas are not economically feasible. Spruce-fir forest was included in the 1987 plan, but was classified as non-suitable for this analysis in **all alternatives** because the majority of it is located in withdrawn lands, not cost-efficient, and/or is located in MSO protected habitat as defined in the current MSO recovery plan (USDI-FWS, 1995).

Alternative A would have the most acres of suitable timberland and the most regulated acres managed for long-term sustained yield of wood products. **Alternatives C and B** would have fewer acres managed for long-term sustained yield of wood products.

In **alternatives A, B, and C**, use of moderate and/or high severity fire for tree thinning and density reduction, especially across large areas of suitable timberland, would increase the risk that those acres could not become regulated with the balanced and sustainable progression of age classes needed to ensure non-declining even-flow⁶ of future harvest volumes.

In keeping with its “natural processes” management emphasis, **Alternative D** provides no suitable timber acres because fire would be relied upon as a primary tool to thin the majority of acres, even in the first treatment entry. Moreover in theory, after the first cutting entry on the few mechanical thinning treatment acres, more natural agents like using fire and endemic insect/disease levels would primarily be relied upon to continue thinning all acres thereafter. The consequence of having no suitable timberlands, is that no long-term sustained yield of wood products could be assured. This does not mean that no volume would be available to supply markets. It only means that industrial volumes of traditional sawtimber and pulpwood would not be ensured for truly long-term sustained yield. See the Total Wood Products section below. Additionally, **Alternative D** allocates the most land to the Recommended Wilderness Management Area which precludes timber management. **If** desired tree cover is destroyed or degraded on these lands, limited reforestation activities would be emphasized to move those acres toward plan desired conditions.

⁶ Non-declining even flow is a policy governing the volume of timber removed from a national forest, which states that the volume planned for removal in each succeeding decade will equal or exceed that volume planned for removal in the previous decade.

Of the **action alternatives**, suitable timberlands may occur within the Community Forest Intermix Management Area in **Alternatives B and C**, but not in **Alternative D**. **Alternative C** provides slightly more acres of suitable timberlands than **Alternative B**.

Under **Alternative C**, mostly ponderosa pine, dry mixed conifer, and wet mixed conifer PNVs would be treated because they dominate the suitable timberland areas. These acres would also receive more mechanical treatments and less treatment by prescribed fire, in order to provide the most wood and tree products for the local economy. Cutting treatments on grassland acres would not be emphasized, because sustained industrial tree production is not appropriate there.

Alternative B emphasizes treatments on both suitable and non-suitable timberlands by including objectives to treat Great Basin grasslands mechanically and use more fire treatments in the piñon-juniper woodlands. Fewer acres of suitable timberland mean this alternative would have fewer regulated acres mandated for long-term sustained yield.

Under **all alternatives**, if proposed plan desired conditions are met and maintained by the cutting practices used, the non-suitable timberland acres should provide long-term sustainable tree cover, although they would not be subject to the ASQ volume or LTSYC controls.

Under **all alternatives**, monitoring of soil productivity loss, and vegetation type conversions following severe burns will be needed during the next 1+ decades, to facilitate adjustments (reductions) in the acres classified as “suitable timberlands”.

Allowable Sale Quantity (ASQ)

Industrial wood volumes from annual cuts on suitable timberlands that were modeled in VDDT are summarized for decades 1 through 5 in the next table (table 9).

Table 9. VDDT Model estimated industrial harvest volumes (in CCF) cut annually from all suitable timberlands for first five decades.

ALTERNATIVE A Suitable Lands											
AVERAGE Option Volumes in CCF	Decade 1	Decade 2	Decade 3	Decade 4	Decade 5	Alternative A has no high or low treatment objective options.					
Ponderosa Pine	48,116	46,636	45,953	45,300	44,503						
Dry Mixed Conifer	8,617	8,240	8,074	7,947	7,850						
Wet Mixed Conifer	9,192	9,901	10,685	11,546	12,199						
Spruce-Fir Mix	0	0	0	0	0						
Annual Cut Totals:	65,925	64,777	64,712	64,793	64,552						
ALTERNATIVE B Suitable Lands											
HIGH Option Volumes in CCF	Decade 1	Decade 2	Decade 3	Decade 4	Decade 5	LOW Option Volumes in CCF	Decade 1	Decade 2	Decade 3	Decade 4	Decade 5
Ponderosa Pine	83,865	84,081	84,141	84,315	84,665	Ponderosa Pine	12,920	13,044	13,096	13,313	13,355
Dry Mixed Conifer	14,299	14,607	14,748	14,768	14,860	Dry Mixed Conifer	2,203	2,226	2,257	2,290	2,315
Wet Mixed Conifer	14,960	15,188	15,736	17,111	17,720	Wet Mixed Conifer	1,995	1,892	1,825	1,808	1,879
Spruce-Fir Mix	0	0	0	0	0	Spruce-Fir Mix	0	0	0	0	0
Annual Cut Totals:	113,124	113,876	114,625	116,194	117,245	Annual Cut Totals:	17,118	17,162	17,178	17,411	17,549
ALTERNATIVE C Suitable Lands											
HIGH Option Volumes in CCF	Decade 1	Decade 2	Decade 3	Decade 4	Decade 5	LOW Option Volumes in CCF	Decade 1	Decade 2	Decade 3	Decade 4	Decade 5
Ponderosa Pine	178,278	125,155	107,457	102,491	100,786	Ponderosa Pine	21,219	20,238	19,519	19,204	19,105
Dry Mixed Conifer	33,576	25,347	24,462	25,077	26,049	Dry Mixed Conifer	4,013	4,007	4,081	4,210	4,339
Wet Mixed Conifer	48,032	43,253	42,536	42,455	42,861	Wet Mixed Conifer	4,823	4,324	4,107	4,052	4,161
Spruce-Fir Mix	0	0	0	0	0	Spruce-Fir Mix	0	0	0	0	0
Annual Cut Totals:	259,886	193,755	174,455	170,023	169,696	Annual Cut Totals:	30,055	28,569	27,707	27,466	27,605
ALTERNATIVE D has ZERO Suitable Lands = zero volumes cut would contribute to the ASQ.											

ASQ is expressed as an annual average of industrial wood cutting volumes from suitable timberlands to meet multiple-resource objectives and public demand. Next, table 10 displays the total ASQ volume estimates for the first decade needed to implement restoration treatments (as modeled in VDDT) and to also meet average public demand for sawtimber, pulp, and pole sales (not modeled in VDDT). Because the modeling done only represents one possible green-tree cutting scenario to implement each alternative, the resulting volume outputs shown previously are too precise for a forestwide programmatic assessment. Therefore, all ASQ values are rounded to the nearest thousand CCF for alternative comparison.

Table 10. ASQ volume from suitable timberlands for the first decade (rounded to the nearest thousand)

	Alternative A ²	Alternative B		Alternative C		Alternative D	
Cutting Treatment Objective Level	Average ³	High	Low	High	Low	High	Low
Annual Range of ASQ, in CCF ¹	N/A	122,000	26,000	268,000	39,000	0	0
Annual Average ASQ, in CCF ¹	73,000	74,000		153,000		0	
¹ All volumes shown above for all alternatives include green tree harvests plus extra annual average chargeable volume sold on incidental public permits and small salvage sales, to reflect the total commercial timber management program. Because incidental permits and small sales are based on public demand, their additional volumes are assumed to be constant across all alternatives. This extra sales amount is estimated at 8,467 CCF per year, based on district and forest records.							
² Alternative A's ASQ was recalculated using the same methodology as the other alternatives.							
³ Alternative A has no high to low range, only the annual average.							

Alternatives A, B, and C would have different ASQs because they were based on the expected level of cutting treatments on suitable timberlands (see table11). **Alternative D** would have no ASQ volume because there are no suitable timberlands.

ASQ volume for **alternative B** would be 122,000 CCF per year, as the maximum allowable sale quantity from suitable timberlands. Under **alternative C**, it would be 268,000 CCF maximum for any given year. For comparison, the 1987 Plan Amendment #1 ASQ volume was 198,000 CCF. The highest total harvest in recent years, occurred in 2011, and was approximately 103,000 CCF (Drury, May 2012).

Consequences of implementing the alternatives are that **alternative C** would offer the most ASQ volume of traditional sawtimber and pulpwood offerings for sale to the markets that may desire these products.

Alternative A would offer less ASQ volume than **alternatives B and C** because it does not have a high treatment objective. This amount or some higher amount could become the replacement ASQ for the 1987 plan if there had been no other needs for change to revise the plan (see chapter 1).

Suitable timberland acres are capable of producing greater annual harvest volumes than those shown in the above table. However, use of planned and unplanned fire under **alternatives B, C and A** would reduce available green harvest volumes, because varied amounts of moderate and/or high severity fire would be used to intentionally thin the forests.

Alternative B would reduce the most green volume because it proposes using more moderate and/or high severity fire to thin trees on forested lands, including some suitable timberlands. **Alternative C** would use less moderate and/or high severity fire on suitable lands to thin trees, resulting in more green volume available for cutting and industrial wood. **Alternative A** would reduce the least green volume killed by fire, because it uses the least moderate and/or high severity fire.

Trees intentionally killed by use of fire treatments may or may not be salvaged. Salvage of such trees would require removal within approximately 3 to 4 years post-burn before wood borers and other decay agents reduce their amount of solid wood fiber and market utility. Under **all alternatives**, unanticipated large-scale salvage volume (e.g., wildfire-kill, blow-down, insect/disease mortality) does not count towards the ASQ per the National Forest Management Act.

The ASQ volume control concept enacted by law is intended to prevent excessive tree losses due to “over-cutting” management beyond sustainable forest levels on suitable timberlands. Annual monitoring of cutting levels compared to the ASQ and regular reviews of the suitable timberland acreage classification are required, in order to reveal adjustments that may be needed before too many acres are over-cut in the long-run. However, ASQ does not adequately prevent excessive cutting on non-suitable lands, nor for losses due to the use of fire as an intentional tree-reduction tool. This analysis does not investigate “over-burning” management beyond sustainable forest levels, because no similar review process is required by law. Therefore, careful annual monitoring of prescribed fire-kill volumes on suitable timberlands can only be strongly recommended here, for comparison to the LTSYC in the first decade. Use of unplanned ignitions as “resource benefit” managed fires especially merit this monitoring.

Base Sale Schedule

This planning effort emphasizes proposed management *outcomes* rather than *outputs*. The desired outcome is to restore the forest type PNVTs toward desired ecological conditions, while also providing wood products to the economy as a *byproduct* of the restoration activities. Therefore, listing definite site-specific volume outputs tied to individual sales listed for each of ten years is not appropriate to provide here as a forest program target. Each revision action alternative offers a flexible range of annual cutting volumes, based on the realistic objective levels that help to frame the alternative. Annual volume levels offered for sale will vary as budgets, market demand, and opportunities occur.

For example, the annual cutting level for **Alternative B** may vary from one year to the next between the high and low range of ASQ volumes shown in table 10 above, provided the decade total does not exceed the annual average times ten. Therefore, forest-wide ASQ cutting volumes could fluctuate between 121,591 CCF and 25,585 CCF each year, provided that the total maximum volume of all cuts in the decade would not exceed 735,880 CCF for the 10-year total ASQ. The same interpretation would apply for **alternative C** should it be selected.

In contrast, **alternative D** would have neither of these limitations placed on its acres proposed for cutting; and **alternative A** has no high to low range, only the annual average.

If funding and workforce capacity should somehow become available to offer the high level volume of the selected alternative every year for the first decade, then that level would not be permissible within the ASQ decade total. However, given the high degree of “ecological departure” situation that these lands are currently in, a plan amendment could be justified to re-evaluate the ASQ for an increase to meet multiple-use objectives.

ASQ volumes from suitable timberlands only constitute a fraction of the total wood products that would result from cutting treatments implemented to restore forested acres toward the ecological desired conditions. In reality, a majority of industrial tree species in the traditional sawtimber and pulp size classes are no longer sold as these products. They are currently sold as fuelwood, and/or extracted from the forest and scaled as tons of biomass, which are not included in the definition of ASQ volume. This trend is expected to increase, as the nation continues to emphasize alternative energy (heat and electricity) generation from green biomass.

Therefore, any administrative ten-year action plan developed for forest vegetation management projects across all ASNFs lands (both suitable and non-suitable) is expected to reflect much higher total wood volumes than the ASQ volume levels stated in the Forest Plan base sale schedule stated above.

Long-Term Sustained-Yield Capacity

The LTSY level of 24 CF per acre per year becomes a baseline for comparison of estimated wood product outputs (volumes) by alternative. Total acres of suitable timberland in each alternative x 24 CF per acre, then converted to CCF derives the LTSYC. When rounded to the nearest thousand, the LTSYC for **alternative A** is approximately 184,000 CCF, **alternative B** is 143,000 CCF, and **alternative C** is 145,000 CCF. See table 11 below.

Under **all alternatives except D**, planned, scheduled entries of tree cutting on suitable timberlands would be necessary to move the forests toward an uneven-aged (regulated) balance of age classes and then to maintain the desired condition.

To comply with legal requirements of the National Forest Management Act (NFMA) and Multiple Use Sustained Yield Act (MUSYA), long-term sustained yield also means that ASQ volumes harvested from suitable timberlands cannot decline from one decade to the next. Ideally, harvest volumes below the LTSYC should continue increasing to eventually reach the LTSYC and then level off at or near that regulated value. The only exception to this rule is if the cutting volumes are departed above the LTSYC, in which case they would be expected to decline toward the LTSYC over time.

Long-term Sustained Yield Capacity in Relation to Estimated Industrial Harvest Volumes

As each alternative has different amounts of suitable timberlands, the LTSYC varies accordingly. Table 11 displays the annual ASQ volumes based on the high cutting treatment level for decades one to five, compared to the LTSYC for each alternative.

Table 11. Estimated annual ASQ volume by decade by alternative (rounded to the nearest thousand)

Annual High¹ Cutting Treatment Volumes in CCF	Alternative A Annually Cuts 10,041 average acres² of 764,872 acres of suitable timberland LTSYC = 184,000	Alternative B Annually Cuts 14,037 high acres of 596,743 acres of suitable timberland LTSYC = 143,000	Alternative C Annually Cuts 31,893 high acres of 604,746 acres of suitable timberland LTSYC = 145,000	Alternative D Annually Cuts only on non- suitable lands LTSYC = 0
ASQ Decade 1	73,000	122,000	268,000	0
ASQ Decade 2	73,000	122,000	202,000	0
ASQ Decade 3	73,000	123,000	183,000	0
ASQ Decade 4	73,000	125,000	178,000	0
ASQ Decade 5	73,000	126,000	178,000	0
¹ Alternative A only has an average ² Acres are based on the estimated cutting treatments modeled for ponderosa pine and dry and wet mixed conifer forests on suitable lands only				

As seen above, when cut at the highest treatment objective levels modeled, **alternatives A and B's** volumes would not decline and would remain below the LTSYC. **Alternative A's** ASQ unrounded volumes for decades one through five are all within 1 to under 2 percent of each other, which indicates a flat line of sustained yield harvests. By cutting at a relatively level trend across all five decades as modeled, **alternative A** would fail to reduce any backlog of overgrowth by just barely keeping up with new growth each decade. VDDT methodology used in this analysis did not permit the ability to model the most logical changes in cutting methods for subsequent re-entries on acres previously treated with the model inputs. By decade three, less intermediate thinning treatments to cut smaller-sized trees would be used; instead more uneven-aged group selection cuts which require cutting bigger trees would be used, thus producing greater harvest volumes than those shown here for decades three through five.

Alternatives A and B comply with legal requirements by cutting at levels which do not decline and are below the LTSYC. The first five decades of VDDT modeling do not produce substantially increasing harvest volumes that ramp up closer to the LTSYC, due to predicted cutting levels on suitable timberlands according to budget and workforce estimates for these alternatives in this planning period. In a regulated (sustainable) forest, annual cut equals annual net growth, such that the forest never becomes overgrown or stagnant. **Alternatives A and B**, because they produce harvest volumes below the LTSYC, would continue to result in overgrown forests that would be more susceptible to uncharacteristic disturbances (e.g., severe wildfire, insect/disease outbreaks). These undesired events could result in additional deforested acreages. **Alternative B** would make more progress than **alternative A**, but still would not reach the LTSYC by decade five.

In contrast, the high objective ASQ volumes for **alternative C** would decline while exceeding the LTSYC in a departure situation. **Alternative C** would reduce the most amount of overgrowth backlog in the first three decades, and it would continue to remove more backlog in decade four until it levels off at decade five. Any departure (exceedance) level of cutting above the LTSYC such as seen in **alternative C**, especially in the early planning decades, is justifiable when any of the following situations occurs: 1) a need to drastically reduce or prevent high tree mortality losses from any cause; 2) it is possible to improve timber age class distribution, thereby facilitating the attainment of LTSYC; 3) implementation of the corresponding base sale schedule

would have a substantial adverse impact upon a community in the local economic area; and 4) it is reasonable to expect that it would be better to attain overall multiple-use objectives in other ways; (source: see Appendix F for directives). All four situations apply to the Apache-Sitgreaves NFs:

- Situation 1: High tree mortality losses would continue to occur in connection with: uncharacteristic wildfires; insect outbreaks abnormal in either intensity or acreages affected; elevated disease levels; new disease arrivals; and accelerated tree stress and deaths resulting from over competition with each other.
- Situation 2: Current age class distributions are skewed. Many acres are even-aged with certain age classes missing. A majority of acres still have far too many small-medium diameter trees that act as ladder fuels and aggressive competition for larger trees.
- Situation 3: Several “small tree” based industries have recently emerged to utilize the surplus numbers of saplings and pulp-sized trees that used to go to the paper pulp mill in Snowflake, AZ. Government grant dollars have helped these industries become established as collaborative partners to begin restoring our imbalanced forest conditions. Numerous new jobs have resulted or returned to this area (Hunt, 2012; ERI, 2007; Sitko and Hurteau, 2010; US Forest Service, 2008-CER). Apache and Navajo Counties have been among the most depressed economies in the nation (see SocioEconomics Report). Due to recently rising market demands since 2009, local operators are now keeping pace with treating all acres offered, almost as fast as we can prepare them and make them available (Drury, 2011). Limiting the ability to continue cutting surplus tree volume growth that is far above the LTSYC level could have some degree of adverse impact upon communities in the local economic area.
- Situation 4: Multiple-resource objectives would be best met by correcting the serious imbalance of excessive forest wood volume growth rates, which would help to reduce the risk of uncharacteristic wildfires and other extreme or long-lasting disturbances. These uncharacteristic events are not consistent with the proposed plan desired conditions, which are focused on restoring our forested ecosystem to benefit watershed/soils stability, riparian and aquatics, wildlife and fish habitats, ground vegetation and herbaceous cover, range production, water and oxygen cycles, and recreation experiences, as well as economics and the human environment.

Alternative C, because it produces volumes above the LTSYC, would contribute to the reduction in overgrowth and offer a greater opportunity to maintain forest lands at a sustainable level for at least the first four decades. ASQ cutting departures above the LTSYC can be temporarily justified to correct the imbalance of excess net growth, provided the volumes cut decline over time to eventually level out at or below the LTSYC. This is the case for **Alternative C**. This declining volume trend came from the VDDT model runs for decades one through five and is based on treatment inputs for each alternative that are documented in Appendices C, B5, E1, and E2. A declining trend is logical when heavy restoration cuts are needed early to prevent excessive tree mortality from severe wildfires, competition, and insect/disease outbreaks. Once overgrowth levels have been reduced, then subsequent decades should produce volumes which taper down toward reaching desired conditions that are intended to promote a more sustainable forest. Because VDDT modeling was not done beyond 50 years, it is assumed that continued aggressive cutting levels beyond decade five would be needed to bring forested conditions closer to desired conditions and the LTSYC.

Non-declining even flow of harvest volume from one decade to the next is not expected for **alternative C** until desired conditions are met. By the fifth decade, **none of the alternatives**

would actually treat enough acres fast enough to fully reach desired conditions within the first five decades because the alternatives were realistically designed to reflect anticipated budgets and workforce capabilities. Because volumes were not modeled beyond the fifth decade, it is not possible to predict when their ASQ volumes might most closely meet the LTSYC. But in forestry theory, because **alternative C** produces a better-regulated forest the fastest it means that cutting departure above the LTSYC could diminish faster in the future than without such regulation.

Because **alternatives A and B** under-cut noticeably below the LTSYC, they would remain threatened by high mortality losses to uncharacteristic disturbance events. At some point **alternative C** would need to align with the LTSYC (i.e., regulated forest) to prevent over-cutting. VDDT modeling indicates that after 50 years of treatments the forested PNVTs would not fully reach desired vegetation conditions. Review of all VDDT model run vegetation outcomes and trends indicate that changes in management strategy would likely be needed following the planning period for **any alternative** (see the Vegetation Specialist Report).

For example, **unlike alternatives A and C**, modeling indicates that **alternative B's** restoration strategy would need to change after this planning period to steadily increase cutting treatments in decades two through five on closed canopy acres and shift to emphasizing low-severity prescribed fire, in order to sustain a non-declining even-flow of ASQ volumes. It is assumed that continued restoration treatments on suitable timberlands toward desired conditions beyond decade five would eventually increase ASQ levels closer to the LTSYC.

Initial modeling methodology used for **alternatives A and C** produced results compliant with the non-declining even-flow legal requirement by continuing the same treatment strategy each decade in the initial level of VDDT modeling. In the case of **alternative B**, however, the initial VDDT model runs which repeated the same treatment strategy in subsequent decades after this planning period originally produced ASQ volumes that consistently declined each decade, while staying below the LTSYC. Therefore, additional analysis at a more refined level of modeling revealed that the treatment strategy would need to change after the 15-year planning period for **alternative B** to produce the compliant results shown here in tables 11 and 12 (also see the “Methodology and Analysis Process” and “Relationship of Short-term Management to Long-term Productivity” sections).

Cuts under **alternative D** are not comparable because no suitable timberlands are present.

It is likely in some years that acres cut would not reach the *high* treatment objective level. So the next table compares the alternatives' *average* treatment objective cutting levels.

Table 12. Estimated ASQ volume and LTSYC in CCF of average objective harvest for decades one through five (rounded to nearest thousand).

Annual Average Objective Harvest Volumes in CCF	Alternative A Annually Cuts 10,041 average acres of 764,872 Suitable acres LTSYC = 184,000	Alternative B Annually Cuts 8,010 average acres of 596,743 Suitable acres LTSYC = 143,000	Alternative C Annually Cuts 17,541 average acres of 604,746 Suitable acres LTSYC = 145,000	Alternative D Annually Cuts only on Non-Suitable Lands LTSYC = 0
ASQ Decade 1	73,000	74,000	153,000	0
ASQ Decade 2	73,000	74,000	120,000	0

ASQ Decade 3	73,000	74,000	110,000	0
ASQ Decade 4	73,000	75,000	107,000	0
ASQ Decade 5	73,000	76,000	107,000	0

At the *average* cutting objective levels, only **alternative C**, during the first decade, would exceed (depart from) the LTSYC. Then it would drop below the LTSYC by decade 2 and stay below it thereafter. **Alternative B** maintains the same patterns as seen previously at the high level. When viewed at the *low* cutting objective levels, **none of the alternatives** exceeds the LTSYC as modeled.

Harvest volumes and departures were not modeled beyond decade 5, nor are they clearly predictable beyond that point. VDDT model trends become less reliable the farther they are projected into the future.

Cutting at the modeled levels is not expected to cause any programmatic irreversible or irretrievable commitment of timber-production resources. In fact, it is more likely that if cuts don't occur at these higher levels, then the risk of adverse environmental impacts from uncharacteristic wildfires and other severe disturbances will not be avoidable. Even with elevated cutting levels, there would remain a risk that uncharacteristic wildfire, insect/disease outbreaks, and/or other major disturbances could alter the excessive growth imbalance before management actions can.

Under **all alternatives**, the cutting and total treatment levels projected for the first five decades would not reduce the forest overgrowth backlog to sustainable levels. Therefore, there would still be a risk of massive tree mortality due to overgrown stand conditions, large uncharacteristic wildfires, and insect-disease outbreaks during the planning period. Should a majority of remaining forest cover acres become deforested in this case, the local wood products industry would likely collapse, move elsewhere, or shift into temporary salvage and reforestation employment. The lost revenues from severely decreased wood products industry, recreation, and other forest uses could have major impacts to the local society (see SocioEconomics Report in the planning record).

Total Wood Products

Trees cut from non-suitable lands can also provide wood and tree products for local markets. Table 13 displays criteria from the draft Plan where any amount of tree cutting can be an appropriate activity for meeting desired conditions, including lands not designated as suitable timberlands. This may involve removing just a single tree of any size, to implementing a clearcut, depending on the area management and/or project objectives.

Cuts from non-suitable lands may be a one-time entry, such as removing encroaching trees from grassland or a new energy corridor. Subsequent cuts may not be needed if desired conditions can be maintained with fire, or other methods. PNVTs with stump re-sprouting species (e.g., alligator juniper, oak species) may need additional cuts (or other tree control methods) that would produce less wood volume than the first entry. Nevertheless there is a rising demand from local markets for "non-industrial" tree species and sizes, such as juniper and piñon pine, whether they are thinned from woodland acres, or from grassland acres, or even from forested PNVT acres.

Table 13. Lands where tree cutting is generally appropriate to meet desired conditions (regardless of timber suitability), by area

Management Area	Tree Cutting ¹	
	Generally Appropriate	Generally Not appropriate
General Forest (includes all PNVTs present, see Other Areas)	X	
Community-Forest Intermix	X	
High Use Developed Recreation Area	X	
Energy Corridor	X	
Wild Horse Territory	X	
Wildlife Quiet Area	X	
Natural Landscape	X	
Recommended Research Natural Area		X
Research Natural Area		X
Primitive Area		X ²
Recommended Wilderness		X ³
Wilderness		X ²
Other Areas		
Areas with soil condition rating of unsuited/inherently unstable		X
Lands not cost efficient in meeting timber production objectives	X	
Grasslands, woodlands, interior chaparral, and riparian forests	X	
Communication sites	X	
Developed recreation and administrative sites	X	
Eligible wild and scenic river	X ⁴	
MSO (Mexican spotted owl) protected and restricted lands	X ⁵	

¹ Appropriate refers to areas that are accessible and operable for cutting with motorized or non-motorized equipment. Most areas are appropriate for non-motorized (e.g., handsaw, axe) tree cutting, regardless of access.

² Trees may be cut in the Primitive Area or Wilderness Management Areas with non-motorized equipment (e.g., axe, bucksaw) and primarily for trail maintenance (FSM 2323.13f, 2323.53, 2326).

³ Trees may be cut in the Recommended Wilderness Management Area with motorized (e.g., chainsaw) or non-motorized equipment for trail maintenance.

⁴ Tree cutting is not appropriate in sections classified as wild except where needed in association with a primitive recreation experience such as to clear trails (FSH 1909.12 Chapter 82.51).

⁵ Cut tree size generally restricted to diameter limits in current MSO Recovery Plan.


The number of total annual mechanical treatment (cutting) acres by alternative for all PNVTs regardless of timber suitability classification) was displayed back in table 5. Table 14 below compares estimated wood product volumes for the first decade of plan implementation by alternative. To simplify table labeling, “ASQ” is used here to represent industrial wood volumes cut only from suitable timberlands, while “Non-ASQ” is used to represent non-industrial wood volumes cut from both suitable and non-suitable timberlands. (Volumes are not rounded here, to display the raw estimates resulting from modeling and recent local wood volume sales.)

Table 14. Estimated ranges of annual wood product volumes potentially available to offer in decade 1, by alternative from all NFS lands (suitable timberlands and non-suitable)

PRODUCT CLASS	Alt A Average	Alternative B		Alternative C		Alternative D	
		High	Low	High	Low	High	Low
Cuts on SUITABLE Lands:							
ASQ Industrial Species¹ (Timber 9+” and Pulp 5-9”) in CCF	74,392	121,591	25,585	268,353	38,522	0	0
Fuelwood (5+” non-industrial conifer and hardwood species) in CCF , Non-ASQ	14,606	17,530	8,533	33,615	10,019	0	0
Biomass (0+” non-industrial sizes and species) in TONS , Non-ASQ	323,302	400,667	59,336	1,202,219	128,463	0	0
Cuts on NON-SUITABLE Lands:							
Non-ASQ Industrial Species (Timber 9+” and Pulp 5-9”) in CCF	5,780	17,804	2,959	31,192	3,402	48,403	6,065
Fuelwood² (5+”non-industrial conifer and hardwood species) in CCF , Non-ASQ	10,976	76,528	46,633	18,413	8,699	59,438	32,203
Biomass (0+”non-industrial sizes and species) in TONS , Non-ASQ	24,822	185,132	82,848	122,548	13,418	246,798	66,026
SUMMARY of TOTAL CUTS on ALL TREATED LANDS (ASQ and NON-ASQ COMBINED):							
Industrial Species¹ (Timber 9+” and Pulp 5-9”) in CCF	80,172	139,395	28,544	299,545	41,924	48,403	6,065
Fuelwood² (non-timber conifer and hardwood species) in CCF	25,582	94,058	55,166	52,028	18,718	59,438	32,203
Biomass (non-industrial sizes and species) in Tons: or ³ CONVERTED TO CCF:	348,124 or 99,464	585,799 or 167,371	142,184 or 40,624	1,324,767 or 378,505	141,881 or 40,537	246,798 or 70,514	66,026 or 18,865
GRAND TOTAL of ALL WOOD PRODUCTS, ALL in CCF :	205,218	400,824	124,334	730,078	101,179	178,355	57,133
AVERAGED GRAND TOTAL of ALL WOOD PRODUCTS, ALL in CCF :	Alt. A 205,218	Alt. B Average 262,579		Alt. C Average 415,629		Alt. D Average 117,744	
¹ Industrial species for all alternatives include different live trees modeled in VDDT for restoration cutting, plus additional constant volume sold in small sales and on TIM permits (miscellaneous live and dead small salvage sales, road and recreation site hazard trees, pulp and poles). ² Fuelwood for all alternatives is different live trees modeled for restoration cutting plus additional constant TIM permit sales for dead/down fuelwood sales, plus posts sold in TIM. ³ Conversion factor used: 3.5 tons = 1 CCF. Source: R3 Measurements Specialist, based on R3 weight scale study conducted locally.							

Converting biomass tons to CCF in the above table provides a comprehensible measure for fair comparison between alternatives when all three wood product categories are added together as one total CCF.

The total estimated annual average treatment objectives cutting volumes are graphed in the following figure (figure 2), with ASQ and non-ASQ volumes combined.



Alternative C would provide the highest average wood products volume for the first decade, followed by **alternatives B, A, and D**. However, the **action alternatives** would provide more average volume from non-suitable timberlands in the first decade than **alternative A**.

Under **all alternatives**, if plan desired conditions are met and maintained by the cutting practices used, the non-suitable timberland acres should provide long-term sustainable tree cover. However, these lands would not be subject to the ASQ volume or LTSYC controls.

Not included in the volume estimates are additional dead tree volumes from intentional thinning with moderate and/or high severity fire (both planned and unplanned ignitions). If salvage volumes of fire-killed trees were included, **alternatives A, B, and D** could possibly approach **alternative C** for total wood products available for at least the first few decades. However, this fire salvage would have to be harvested within approximately 3 to 4 years post-burn before decay agents destroy its wood fiber integrity.

In general, **Alternatives A, B, and C** would use a wide variety of mechanical treatment methods to meet multiple ecological needs. Because **Alternatives A and B** would emphasize the retention of more large and old trees than **Alternative C** would, use of certain cutting methods may be more limited under these two alternatives (see Appendices B1, B2, B5, and E2).

Alternative A would be reliant on tree cutting as the primary tool to thin the forest, with fire used mostly as a secondary, slash cleanup tool. This approach is slow and costly. Currently, it is uneconomical to move raw cut materials more than 70 miles. Raw wood values are so low that little cut volume is sold. Currently, the Forest Service pays local operators to cut and remove the volumes from the White Mountain Stewardship projects. Where it is uneconomical to move raw wood, a portion of the cut volume may be left and would be disposed of at additional cost to the government. As displayed in the previous table, **alternative A** would provide far less volume to support large, landscape-scale restoration efforts like the Four Forests Restoration Initiative (4FRI, see the cumulative effects section), than would the high and average treatment objective levels of **alternatives C or B**. Yet, **alternative A** would provide more volume if these two alternatives were to consistently be implemented at their low objective levels.

The **action alternatives** would rely on using fire as a primary tool to thin more of the forest (kill trees) than **alternative A**. Thus, less green wood and more dead and fire-charred wood available as a harvestable byproduct from these alternatives, provided enough woody material is left on-site for ecological needs such as soils stability, site productivity, and wildlife habitat.

Alternative B would fall in between the cutting levels of **alternatives C and D**, due to the blend of treatment methods and acreages proposed (see Appendices B1, B2, B5, C, and E2). It employs a tree cutting and burning strategy that restores more acres faster toward desired conditions than **alternative A**. It would not reach a regulated supply of sustainable timber as fast as **alternative C**, but it would do so faster than **alternative A** and would be more sustainable than **alternative D**. Less suitable timberland acreage enables more non-suitable lands to be treated with the fire-only method, in order to reduce the high costs associated with mechanized thinning, so that more acres can be treated annually overall by **alternative D**.

Alternative C would emphasize a mix of more cutting treatments designed for optimum commercial timber species volume production (maximized growth and harvest) on suitable timber acres, such that it should produce more total wood volumes than the levels that would be harvested in the other alternatives. **Alternative C's** high objective cutting level would produce the most total wood volume to support large, landscape-scale restoration efforts like 4FRI.

However, in the first decade this alternative's low objective cutting level would produce less total volume than either the high or low objective level of **alternative B**. This is because **alternative C** is focused on treating mostly suitable timberlands, while **alternative B** spends the first decade focusing on restoring grasslands and other non-suitable lands that can provide high volumes of non-ASQ wood products in addition to the ASQ volume.

Alternative D would provide the least wood volume for meeting social and economic desired conditions for local and regional markets and related jobs. It would produce the least wood product volumes, due to its emphasis on using fire as the primary treatment method, as well as a 16-inch diameter cap imposed on the few acres that are mechanically thinned or cut. After one cutting entry, those mechanical acres would be maintained by regular intervals of planned and unplanned ignitions thereafter. Long-term consequences of continuing understory burning beneath an unthinned overstory may result in failure to meet desired conditions. An over-abundance of large trees could eventually result with a generational gap lacking younger trees to replace older trees over time, as has been seen occur on acres of recent fuels reduction projects on the Apache-Sitgreaves NFs (Triepke et al., 2011; Abella et al., 2006; Youtz, 2010a; Drury, 2011). Dwarf mistletoe infection left in the unthinned overstory would also be difficult to control as it tends to intensify and spread to adjacent trees and the understories, causing mortality which interferes with the uneven-aged forest structure desired (see the Forest Health Report).

Not included in the above volume estimates are the additional dead tree volumes indirectly created by intentionally using moderate intensity prescribed fire as a tree-thinning tool (both planned and unplanned ignitions). Only volumes of live trees cut were modeled for this analysis, with incidental salvage volumes of wildfire-killed trees included. If salvage volumes of prescribed fire-killed trees were included, then Alternatives B and D might possibly approach Alternative C, for total wood products available to offer, at least in the first few decades. Estimates of this extra prescribed fire-created volume would require further analysis of the R3 FVS-FFE fire simulations.

As a major consequence of using fire to more quickly and cheaply kill trees of all sizes in **alternatives D, B, and A**, less green volume of higher market value wood would be available for harvest. Instead, lower market value wood would be offered from those acres, and it would have to be salvaged within about three years to still possess its solid fiber density and integrity that most markets need as a raw resource. Moreover, some markets cannot, or will not, use black-charred wood (e.g. pulpwood for paper manufacturing and wood heating stove pellets).

It is conceivable that intentional fire use to reduce treatment costs of moving toward vegetation desired conditions ultimately can create an extra cost when the resulting tree mortality volumes would need sale layout/preparation/administration (and possibly extra NEPA process) costs to offer the volume afterward for salvage. By thinning it mechanically first, with a low severity burn follow-up treatment, the green volume that could be provided for higher market price to meet social/economic desired conditions might turn out to be just as cost-effective as burning it at high and/or moderate severity and then salvaging the fire-kill.

Likewise, no long-term supply of wood volume could continue to come from grassland acres once they are restored by cutting and maintained by fire thereafter (basically a one-time harvest entry). **Alternatives B and D** would rely more heavily on fuelwood and biomass cut from grasslands than **alternative A**. **Alternative C** would rely the least on fuelwood and biomass cut from grasslands and other non-suitable timberlands.

Use of moderate and/or high severity fire combined with diameter caps as thinning tools used most heavily in **alternatives D and A**⁷ would not assure removal of encroached off-site tree species of seed-cone bearing size/age for restoration to the correct fire regimes, thus perpetuating a species mix that is not well adapted to the site (Appendix G and Triepke et al., 2011). These two management methods are less certain to selectively leave the tree sizes, species, and arrangements desired for a truly sustainable uneven-aged structure and predictable wood volume production in more challenging climate conditions. Failure to restore all forests and woodlands to their correct species composition, size distribution, and spatial arrangement can make them more vulnerable to climate shifts or other uncharacteristic disturbances. **Alternatives C and B** would utilize no 16-inch diameter caps. **Alternative A** would use fewer diameter caps than **alternative D**. Therefore **alternatives C and B** would provide more control over tree species composition on their mechanically treated acres.

Under all alternatives, there would be risks of using prescribed fire as follow: (1) some prescribed fires could continue to burn after field conditions change from the desired burn prescription; (2) some prescribed fires could escape and become wildfires; and (3) some planned ignition fires may not get accomplished due to narrow burning windows and/or smoke management constraints; (4) use of high and/or moderate severity burns may result in more acres needing reforestation efforts (e.g. Wilkins, Weimer-Durfee, and Wagon Draw resource benefit fires, per Richardson, 2012). All of these risks would be higher for **alternatives D and B**, over **alternatives C and A**, in this order.

Under **all alternatives**, restoration and maintenance of green tree thinning could be reduced if large salvage sales of fire-killed trees dominate the Apache-Sitgreaves NF's workload for the next 10-15 years. This could elevate the risk of losing even more acres to uncharacteristic disturbances. Under all alternatives, too much emphasis and time (NEPA, layout, administration) spent on efforts to salvage dead trees would detract from treating the overgrowth backlog that still threatens remaining green forest acres.

Forested/Overgrown Lands

The following table (table 15) displays the estimated percent of forested/overgrown lands that would be treated within the planning period. If continued at the same rate, the minimum number of years needed to treat all the forested/overgrown lands (approximately 673,437 acres) just once, is also shown. Cutting and burning treatments on suitable and non-suitable timberland acres are included. (For tree planting rates, see table 16.)

⁷ Alternative A (1987 forest plan) does not specify a 16-inch diameter cap. However, this diameter cap has been used as a treatment in recent and current vegetation management. See "Other Planing Efforts" section of this report.

Table 15. Percent of forested/overgrown acres treated by year 15, and average years required to treat¹ all acres in need of thinning with one entry, by alternative average objective levels (both suitable and non-suitable acres are included).

Alternative	Total Annual Thinning ¹ Treatments (acres) ²	Percent of Forested/Overgrown Lands Thinned Annually ²	Percent of Forested/Overgrown Lands Thinned by Year 15	Years Required to Thin All Acres of Forested/Overgrown Lands ²
A	16,182	2.4 %	36 %	41.6 years
B	20,037	3.0 %	45 %	33.6 years
C	30,220	4.5 %	67 %	22.3 years
D	28,914	4.3 %	64%	23.3 years

¹ Treatments include cutting and the use of fire, but not planting.

² These values are for years 1-15 in all alternatives. The treatment rates are held constant here for a consistent basis of comparison across all alternatives, even though additional modeling shows that alternative B would need to modify annual cutting rates on just the suitable acres in years 16-50 to provide non-declining ASQ.

Annual thinning treatment percentages in **all alternatives** would affect a minor amount (less than 3 to 5 percent) of the forested PNVTs each year, with nature managing the rest. Without fire as a thinning tool, treatment rates for all alternatives would be far slower. Restoration rates could be too slow and costly without incorporating fire treatments, in addition to cutting, as a thinning tool to increase treated acres. Those untreated areas would still have trees that continue to grow and die, are affected by insects and diseases, and possibly burned or affected by other disturbance processes.

According to analysis data displayed in tables 5 and 15, and Appendix E1, the small percentages of forested acres treated annually would take a number of decades just to implement the first treatment entry toward desired conditions. On most PNVTs desired conditions would need more than one entry to be achieved, especially when a cutting cycle of 30 years is recommended between cutting entries, and 3 or more distinct age classes need to be developed for uneven-aged structure.

According to table 15 above, **alternative A** would only treat about 36% of the forested/overgrown acres by year 15. The **action alternatives** would move more acres toward desired conditions faster, at treatment rates of about 45 to 67% of ground covered in the first 15 years. Only at these higher rates could subsequent entries manage to return back to the same acres again in about 20 to 30 years for follow-up treatments to maintain the benefits of the prior entry. (See the “Relationship of Short-term Management to Long-term Productivity” section for the modified cutting rate modeled on suitable timberlands for **alternative B** after year 15.)

The true challenge under any alternative will be to keep treating new acres each year, while at the same time returning to maintain those acres already previously treated. For an initial time, (varies by alternative), total annual treatment acres would have to increase exponentially to achieve both. This analysis reveals that **alternative C** would come closer to achieving this on suitable timberlands, and in the Community Forest Intermix Management Area than the other alternatives would.

Because the best we can do is only thin forestlands at a rate of 3.0-4.5% per year, it becomes critically important to strategically place those treatments in the right priority locations across the landscape.

This consideration makes **Alternative B** more advantageous over the other alternatives because of its emphasis to prioritize treatment locations inside the community wildfire protection plan (CWPP) areas and in priority watersheds which include large acreages of untreated pine and dry mixed conifer forests. These areas are located in Navajo and western Apache counties nearer to rail lines and centralized markets, with a greater success of being implemented as wood product transportation costs increase. **This alternative** also emphasizes more acres of fire treatments as a faster and more cost-effective restoration treatment that is less dictated by fluctuating wood markets. It also features treatment of grassland acres, which can function as natural fire-breaks when included as part of larger restoration projects on the landscape. This alternative could probably come close to meeting the needs of 4FRI markets and others, while also restoring the PNVTs most in jeopardy. By year 15, it would treat about 45% of the forested/overgrown acres.

Alternative C would emphasize treatments on suitable timberlands, the Community-Forest Intermix Management Area, and other lands that can contribute wood products. The suitable timberlands may or may not be near rail lines or centralized markets. **Alternatives A and D** would emphasize mechanical treatments around communities and in the Community-Forest Intermix Management Area (a subset of CWPPs), many of which have already been treated and now only require follow-up maintenance thinning that may produce less total wood volume in subsequent entries.

Alternative C would accomplish treatments the fastest at a rate 23 years to cover all acres in need of thinning, followed by **alternatives D, B, and A**, respectively. **Alternatives A, B, and C** would all use a mix of cutting on some acres with burning on other acres. **Alternatives C and D** treatment rates would permit more timely return entry intervals for required maintenance of restored desired conditions on the most acres.

In contrast, **alternative D** would accomplish treatments in just 24 years by using fire as the only tree thinning tool on many treated acres in need of thinning. Moreover, all cuts done under **alternative D** exclusively use diameter caps for large tree retention emphasis, which would stall progress toward or move those acres away from many desired conditions (Triepke, 2011; Abella et al., 2006). This is not evident in the VDDT model results for **alternative D** because the benefits of fire use on so many acres overshadow the negative cutting results.

This analysis shows that total treatment levels projected for the first 5 decades **under any alternative** never do get the forest overgrowth backlog down to sustainable levels; and so more massive tree mortality due to overstocked stand conditions, large uncharacteristic wildfires, and insect-disease outbreaks are inevitable to occur during this planning period, **regardless of the alternative chosen**.

Deforested/Early Development Lands

Once adequate quantities of green seed have been collected for each native tree species, then each alternative could begin planting activities on deforested lands. Cone collection may take about 3 to 10 years from present, depending on each species, but it would be the same constant for all alternatives.

For the estimated 37,695 acres (4% of all forested PNV lands) proposed for artificial reforestation the following planting table (table 16) shows a similar timeline for establishing plantations. Both suitable and non-suitable acres are included. It displays the estimated percent of deforested lands that would be treated within the planning period. If continued at the same rate, the minimum number of years needed to plant all deforested acres targeted for artificial

reforestation is also shown. Plantation success is assumed with the initial planting. If there is poor tree survival with replanting or additional fill-in planting required on some sites, then the timeframes shown would be longer.

Table 16. Percent of deforested acres planted by year 15, and average years required for each alternative to accomplish planting treatment one time, on deforested acres targeted for artificial reforestation (both suitable and non-suitable acres are included).

Alternative	Average Annual Planting Treatment Acres	Percent of Forest Planting Acres Treated Annually	Percent of Forested/Overgrown Lands Thinned by Year 15	Years Required to Plant Needed Forest Acres
A	880	2.3 %	35 %	42.8 years
B	1,623	4.3 %	65 %	23.2 years
C	2,066	5.5 %	83 %	18.2 years
D	413	1.1 %	17 %	91.3 years

Alternative C would plant the most acres, especially on suitable timberlands in order to return them into timber production as soon as possible. This rate is at the extreme high end of current workforce capabilities. **Alternative B** would plant at a rate consistent with current workforce capability, and it would focus on reforesting more of a mix of both suitable and unsuitable lands for ecological recovery emphasis, including some Mexican spotted owl habitat.

Alternative A would plant at the lower end of current workforce capacity, focusing primarily on sites near private lands and along highly visible roadways. **Alternative D** would emphasize letting natural processes dominate so that the vast majority of acres needing reforestation would be left for natural regeneration to occur on nature’s timeline. The few acres planted would be near private lands and in some Mexican spotted owl habitat identified for accelerated recovery.

At the planting rates modeled (see Forest Products Specialist Report in the plan set of documents), **alternative A** would plant an average of 880 acres a year, thereby treating all of the deforested acres proposed for artificial reforestation within 43 years. **Alternative B** would plant an average of 1,623 acres a year, thereby treating all of the deforested acres proposed for artificial reforestation within 23 years. **Alternative C** would plant an average of 2,066 acres a year, thereby treating all of the deforested acres proposed for artificial reforestation within 18 years. **Alternative D** would plant an average of 413 acres a year, thereby treating all of the deforested acres proposed for artificial reforestation within 91 years.

Under **all alternatives**, rates of natural conifer regeneration would be the same. Regeneration occurrence and survival would depend upon local site conditions and climate over time.

However, a portion of deforested acres may not return to a productive forested condition. A body of research in the southwestern ponderosa pine forest type has shown that, on average, approximately 50% of severely burned acres tend to convert to grass/forb/shrub/rock lands that can take a very long time to return to forest, possibly centuries (Roccaforte et al., 2012; ERI, 2011; Savage and Mast, 2005; Strom and Fule, 2007). Artificial reforestation is needed to reduce that percentage (Higgins, 2008; ERI, 2012b). Climax spruce-fir and upper elevation (wet) mixed conifer forests have been documented as also undergoing very long-term type conversions to grass/forb/shrubland following severe fire (Alexander 1974).

These environmental reactions can be highly variable, depending on factors such as: unfavorable soil, site and weather conditions for seed germination and/or successful seedling establishment; quick domination by prolific root-sprouting species such as oak, aspen, and other woody shrubs; stiff competition for bare soil, water and nutrients by thick grass cover; a lack of cone-bearing seed trees; size of created openings too large for wind-dispersed seed to reach the opening interior from seed trees on the perimeter; erratic seed production, topsoil loss due to water/wind erosion; substantial rodent, bird, insect consumption of tree seed (Jones, 1974; Alexander, 1974; White, 1985; Savage and Mast, 2005; Publick et al. 2012). Thick seedling/sapling cover (“hyper-dense forest”) that can become established may also be vulnerable to fire reburn destruction (Savage and Mast 2005), and to insect damage (see Forest Health Specialist Report), as well as some wildlife damage. If more workforce capacity and funding become available, more acres of possible grass/shrub site conversion lands could be planted for return to forest cover.

The existing condition has 21.5% of all forestland acres in an undesired deforested condition. VDDT modeling shows that by year 15, **Alternative A** would have improved to 6.6% of ground left in deforested condition, while **Alternative B** would have 5.7%, **Alternative C** leaves 6.5%, and **Alternative D** leaves 6.7%. These figures include natural and artificial reforestation combined within the model transitions, as the results of cutting, burning, planting and natural stage progressions. Even though **Alternative C** plants more acres annually, curiously **Alternative B** is indicated by VDDT as the better management strategy for reforestation overall. Under **all alternatives**, no tree planting could be done on forest lands expected to convert into grass/forb/shrub/rock cover, unless extra workforce capacity and/or funding is made available.

Under **all alternatives**, early development forest lands would need time to grow, with periodic pre-commercial thinning taking place (either mechanically or by prescribed fire) to maintain vigor and facilitate growth into larger size classes. Protection from excessive animal, insect/disease, and fire treatment damage would be necessary. Fire use could be the most cost-effective means of thinning small trees under 5 inches diameter, but has its limits as a silvicultural tool beyond that, especially on suitable timberlands.

Arno and Fiedler (2005, chapters 4 and 16) and Feidler et al., (2010) present various case studies as examples of combining prescribed fire with tree cutting as a complete and effective restoration treatment in western USA forests. They present strong rationale against using fire by itself as the only restoration tool.

Savage and Mast (2005) show a proposed model diagram from their research suggesting that frequent surface fires in the ponderosa pine type could be used to break up “hyper-dense” pine regeneration from suffering another future crown fire, by maintaining clumped pine patches with open spaces between the patches. This concept is presumed to also apply to other PNVTs in Fire Regime I, like dry mixed conifer.

The US Forest Service Southwestern Region used Forest Vegetation Simulator-Fire and Fuels Extension (FVS-FFE) modeling applied to regional FIA plots for each plan revision PNVT to calibrate the VDDT model structural state transitions (Weisz et al., 2012; and Appendix B4). Those results show that when low severity prescribed fire is applied as the only treatment to thin the seedling/sapling states (state *B* = open canopy, and state *F* = closed canopy) in the ponderosa pine and dry mixed conifer PNVTs, a variety of transitions can occur.

Ponderosa pine closed canopy seedling/sapling state *F* responds as follow: 25% of the treated plots transition into the desired open canopy seedling/sapling state *B*; while 42% stayed in the closed state *F*; while another 33% of plots lost enough seedling/sapling sized trees that they converted into larger size classes of trees also present (states *C*, *G*, and *D*).

Should prescribed fire be applied or allowed to creep into ponderosa pine open canopy seedling/sapling state *B* to keep it open, the following responses may result as predicted by the model runs: only 20% of plots stayed in the desired open state *B*; 40% moved into the grass/forb/shrub state *A*; while another 40% of plots lost enough seedling/sapling sized trees that they converted into larger size classes of trees also present (states *E* and *J*).

VDDT model acres of ponderosa pine converted from the seedling/sapling structural state to the grass/forb/shrub state may represent the desired canopy gaps between pine regeneration patches that are desired forest structure to prevent future crown fires. However, these results also suggest that if low severity fire is used to thin acres already regulated for uneven-aged management, fire results may be too unpredictable to maintain the proper balance of regulated age/size classes for long-term sustained yield. Similar plot transition results occurred in the Regional FVS-FFE model runs done to calibrate the VDDT models for the dry mixed conifer and piñon-juniper woodland PNVTs. Thus, using fire-only to thin the early development forest acres would need very careful application and control to successfully meet and maintain desired conditions on suitable timberlands in the frequent fire types. Careful long-term monitoring and adaptive management would be key to this effort.

FVS-FFE model results are more difficult to determine low severity fire effects upon just the seedling/sapling component of the wet mixed conifer and spruce-fir PNVTs, because their structural states combine the seedling/sapling size (<5" diameter) with the small size class (5-9" diameter). Given that these two PNVTs are composed of species that are less fire-adapted, it is likely that even more of the seedling/sapling sized trees would be killed by prescribed fire, with less success to maintain this youngest age class on the landscape. Thinning by cutting would be more preferable to ensure that the correct prescription for desired tree stocking per acre is met and maintained in each size class. Therefore, those alternatives which use the least amount of exclusive fire as a thinning tool on suitable timberlands would be preferred in the infrequent fire forested PNVTs. Waiting until the regeneration has reached merchantable size to thin mechanically could be a more cost-effective solution.

Where maintaining a strong aspen component on the ASNFs landscape is desired long-term, keeping all fire (all severity levels) off of aspen regeneration/early development acres would be imperative (Fairweather, 2008; Roccaforte, Nov. 2012; and see the Forest Health Specialist Report).

Relationship of Short-Term Management to Long-Term Productivity

Several short-to-long-term relationships have been covered throughout this report where pertinent to the topic. In keeping with the natural ranges of tree lifespans and forested ecosystem development, this analysis considers short-term to be within the 15-year planning period, and long-term is considered to be after that time out to 50 years and beyond. Indeed, what actions are implemented today can influence forest conditions for hundreds of years from now. Development of mature forest with a late-successional (old growth) component that was lost in stand replacement disturbances does not occur in the short-term, but the process can be initiated almost immediately. Likewise, conversion from even-aged forest to uneven-aged forest is not likely to fully occur on many acres within 15 years, yet the actions undertaken now will directly affect that outcome in decades hence.

The previous section above addressed the short-term need to return existing deforested PNVT acres to adequately-stocked tree cover in order to improve long-term timber production and ecosystem recovery. There is an additional risk of the possible need to increase reforestation

activities resulting from large acreages treated with moderate and/or high severity fire that may burn out of prescription, which could be higher for **alternatives D and B**, over **alternatives C and A**. Additional deforested acres caused by this management emphasis during this planning period would delay their long-term productivity of harvestable wood fiber, at a possibly higher total long-term restoration cost.

On those acres which are overgrown, short-term cutting departures above the LTSYC level are necessary to bring excessive tree stocking under control as soon as possible. This would enable the treated forest types to reach desired conditions faster than without treatments. It would also improve vigor of individual trees to help withstand or adapt to environmental stressors. In the meantime, such treatments will possibly help to avoid rapid tree losses, and severe losses of other key ecosystem components, to uncharacteristic disturbances expected to continue before the desired conditions are met across the greater landscape. Treatment projects should be located strategically across the landscape to help minimize spatial impacts of such undesired disturbances. As a result, key components like large/old trees, critical wildlife habitats, native species, and soil productivity, would benefit long-term.

In this LTSYC departure context, only **alternative C** would provide such benefits the fastest during the short-term (15-year planning period). **Alternatives B and A** respectively do not provide these benefits to as many forested PNVN acres within the same time frame (refer back to tables 11 and 15 discussion in those sections). **Alternative D** would initially treat (thin) almost as many acres as **alternative C** would within the first 15 years. But because it has no suitable timberlands designated, cutting levels compared to LTSYC and ASQ would remain unknown and untracked over time; so long-term productivity would not necessarily be measured based on net growth compared to controlled cutting levels. Long-term sustainability of uneven-aged desired forested conditions under **alternative D** would certainly not be achieved after year 15, due to the primary uses of fire and large tree retention on a large-scale basis.

In order to sustain a non-declining even flow of ASQ volumes on suitable timberlands in **alternative B**, additional VDDT modeling revealed that the restoration strategy for decades two through five would need to do the following: increase treatment acreages in closed canopy transition vegetation states in the ponderosa pine and dry mixed conifer PNVN; and shift to using low-severity prescribed fire as a maintenance tool for thinning just the seedling/sapling sizes.

These modeling shifts represent adaptive management that is predictable because as more acres are restored to desired open-canopy in these two PNVNs, cuts in each transition state would produce less volume per acre; thus the need to cut more acres overall to sustain the same total volume yields. Likewise, using moderate-high-severity fire as a thinning tool would predictably reduce measurable volume available for ASQ harvest. Thinning only seedlings/saplings that have very little measurable wood volume by using only low-severity fire would not impact available ASQ volume.

These shifts in management methodology could begin after the planning period. It is assumed that continued restoration treatments toward desired conditions beyond decade five would eventually bring **alternative B's** ASQ levels up closer to the LTSYC, provided uncharacteristic disturbances don't occur first to drastically alter the trends shown in this analysis.

The long-term need to keep **alternative B** from producing a declining ASQ yield in decades 2-5 required modeling constraints applied to the ponderosa pine and dry mixed conifer PNVNs (also discussed previously in the LTSYC section of this report), which revealed the following possible effects or consequences of continuing to implement this alternative after this planning period:

- The need to cut increasingly more suitable timberland acres in subsequent decades after the planning period, by especially targeting the remaining acres of closed-canopy structural states. This would exclude the MSO PAC acres because they are not classified as suitable timberlands. The need to cut increasingly more suitable timberland acres makes sense because as more acres are restored to desired open-canopy, cuts in each state will produce less volume per-acre, thus the need to cut more acres overall to maintain same total volume. This also makes sense when considering that in **alternative B** sometime after decade 2 many non-suitable acres (such as grasslands) should theoretically be finished with restoration cutting, and those annual treatment acres could be shifted into additional cuts on suitable timberland acres.
- Budget and workforce levels would need to increase accordingly to mechanically cut more acres each subsequent decade.
- Treatments in the wet mixed conifer type would not need to make such shifts, because desired conditions for this PNVNT do not emphasize as much open-canopy forest.
- Increased acres of cutting in decades 2-5 decreases the re-entry intervals on all suitable timberland acres (all three suitable PNVNTs combined) from approximately 43 years in decade 1, to 31 years by end of decade 5. This new re-entry interval result is comparable to the Southwestern Region's ideal 30-yr. re-entry cycle that they modeled for LTSYC.
- The additional cutting acres treated in decades 2-5 for **alternative B** would still not be as many as those cut in **alternative C**.
- Using a mix of low plus moderate and high severity fire in decade 1 is needed for cost-effective restoration and is realistic to expect with intended use of unplanned ignition fires. Conversion to using all low-severity fire in decades 2-5 on suitable timberlands for ecosystem maintenance is another effect of needing to keep the ASQ volumes up at a non-declining level. This makes sense because low-severity fire only thins (kills) seedling/sapling sized trees which do not contain harvest volume, while moderate-high-severity fires kill larger trees that would reduce available harvest volume.
- If needed, use of moderate and high-severity prescribed fire could still continue to be used on non-suitable timberlands in decades 2-5 to economically thin trees larger than the sapling size for restoration purposes. If leadership wished to continue using moderate and/or high-severity fire as a management tool on all forested lands after this planning period for **alternative B**, the consequence to avoid violating the non-declining ASQ legal requirement would be a need to remove all forested acres from the suitable timberland classification at that time.

Climate Change Considerations

Currently a major strategy to insure that at least a subset of managed forests will not be affected extensively by global change is to provide as wide a range of diversity as possible. Under traditional silvicultural practices, diversity can be provided by a mix of timber stand structural conditions ranging from even-aged monocultures at various developmental stages, to multiage species mixtures; and by applying treatments at a variety of spatial and temporal scales. But silvicultural concepts and practices need to shift toward accepting a tradeoff between traditional industrial plantation and thinning schemes for efficient timber production versus managing forestlands as complex, adaptive ecosystems. As future variability and uncertainty increase, as has been predicted under global change, the increased emphasis on adaptation will become more

important where providing various ecosystem goods and services is most important (Puettmann, 2011). Conversely where efficient production of industrial timber is most emphasized, there may be less certainty of the flexibility needed for forests to adapt to changing environmental stressors.

Forest Plan classification of lands according to potential natural vegetation types (PNVTs) and successional status under the **action alternatives** would more easily enable managers to view forest and woodlands as the complex, dynamic, and potentially adaptive ecosystems that they are (US Forest Service, Dec. 2008-ESR). The current Plan (**alternative A**) only classifies vegetation according to the cover types visibly present at this point in time, with the implication that the cover type present is the one that management should continue to maintain there.

Species-specific tree regeneration needs are increasingly less likely to be met due to loss of mature (seed producing) trees across increasingly large areas where high-severity wildfires and bark beetle-induced mortality have occurred, coupled with ongoing and projected increases in drought stress due to climate change (Williams, et al., 2012). Long-term forest structural and compositional changes and type conversions from forests to shrublands or grasslands are expected to be an increasing risk, especially where tree planting efforts are not well designed to account for changed site conditions and climate.

Tree species currently preferred as “industrial” for timber and other traditional uses may not be as readily available in the next 50+ years or so, as it may become less prudent or less possible to plant them and expect them to mature and reproduce successfully. The early recovery role of native vegetative sprouters (like aspen, oaks, and alligator juniper) on deforested sites is already becoming more important in future forest species composition mixtures. Accordingly, the target tree densities in thinning prescriptions would need adjustment to facilitate success of the more hardy species. The recovery role of native nitrogen-fixing species (like New Mexico locust and other leguminous shrub/herbaceous species for example) may need consideration on nutrient-depleted sites; yet if soil nitrogen levels increase on other sites because of higher atmospheric nitrogen deposition, then the role of nitrogen fixation may be re-evaluated when species mixtures are considered (Puettmann, 2011). Under **all alternatives**, selection of hardier native conifers (like Chihuahua pine and junipers) and non-coniferous tree species for reforestation efforts would be wise to capitalize on genetic diversity most adaptable to a wide range of possible future conditions.

With the diversity of native tree species that exist here today, it is expected that more than one tree species are capable of surviving climate shifts in any direction, provided management efforts enable enough healthy sub-populations to remain with a broad gene pool within the local range of each species. Thus, various wood and tree products should remain available for society’s use during this planning period. Beyond this planning period greater utilization of non-traditional woody species might be expected. Under **all alternatives**, continued monitoring and adjustments to the acres classified as suitable timberlands will be needed.

Under a projected persisting warmer/dryer climate fewer tall, straight-boled trees would be able to grow, for milling large or high-grade solid wood boards. Such boards would become higher priced, as their raw source might be somewhat rare. However, society has already shifted into commonly using oriented strand board, pressed particle board, and finger-joined trim/mouldings in the construction industry now, so that solid wood boards are in less demand.

Climate change and its impacts on forests would likely affect market incentives for investment in biomass technology and wood-conservation techniques. The market for wood products in the U.S. is highly dependent on the acreage, location, and species composition of forests; supplies of wood; technological change in production and use of wood; availability of wood substitutes;

demand for wood products; and international competition. Rising atmospheric CO₂ would increase forest productivity and carbon storage in forests if sufficient water and nutrients were available. Any increased carbon storage would be in live trees. However, in the Southwest and Apache-Sitgreaves NFs, as discussed above, overall production may be limited by a decrease in available water. While increases in wildfire may decrease some available wood supply, treatment of wildland-urban interface and restoration of the fire-adapted ecosystems on the Apache-Sitgreaves NFs and elsewhere may increase the overall availability of small-diameter timber and related wood products (Joyce et al., 2001).

All alternatives could promote a future sustainable availability of various wood products, by moving forested and woodland PNVTs toward desired conditions, which should make these lands more resilient in responding to climate change. By implementing treatments that can reduce losses to drought, insect/disease outbreaks, and severe wildfires, the alternatives would rank in this order from fastest to slowest restoration rates: **alternative C**, followed by **alternatives D, B, and A**.

Multiple socioeconomic impacts often follow drought and severe insect outbreaks. Timber production, manufacturing, and markets may not be able to take advantage of vast numbers of killed trees, and beetle-killed timber has several disadvantages from a manufacturing perspective. In, addition, when insect outbreaks occur, the public often perceives this as an increased fire risk and as detrimental to the aesthetics of montane areas (Ryan et al. 2008). These factors could drive future public policy. Furthermore, wood supplies would vary by forest and woodland type (Sprigg et al. 2000; Joyce et al. 2001).

As increasing tree mortality rates are already underway in relation with these very same climate-related factors, wood markets may be asked to take more dead and black-charred wood than their enterprises can utilize. **Alternative D**, followed by **alternatives B, C, and A** would create more intentionally fire-killed volume in addition to dead trees already being offered for salvage.


Salvaging and converting biomass into boards, fuelwood, and other wood products (as a byproduct of forest restoration) could help reduce carbon loss from fire. Another consideration may be to use biomass for bioenergy production. Bioenergy production can be carbon neutral and could replace the fossil fuels in generators. Mobile generation facilities could provide power to schools, hospitals, and command centers in the event of an emergency.

If new markets for forest biomass to generate heat and electricity in place of fossil fuels should develop locally or regionally, then traditional “non-industrial” wood species and sizes could become more of an “industrial” demand. Indeed, this trend is already somewhat underway. Alternatives offering the most dependable supply of wood volume from both suitable and non-suitable timberlands would provide the most flexibility to meet changing market demands, in this order from greatest to least: **alternative C, B, A, and then D**.

Cumulative Environmental Consequences

The cumulative effects area for this analysis of forest products is the White Mountains-San Francisco Peaks-Mogollon Rim M313A Ecoregion Section and the seven subsections within the ecoregion (see map 1). In addition to the Apache-Sitgreaves NFs, this ecoregion includes: most of the Coconino NF, portions of the Prescott and Tonto NFs, the southern end of the Kaibab NF, and all of the Gila NF and portions of the Cibola NF in New Mexico. Non-Forest Service land ownerships in this ecoregion include: BLM lands; Arizona and New Mexico state lands; Fort Apache and San Carlos Apache Indian Reservations in Arizona; other tribal lands in New Mexico; and private lands.

Map 1. ASNFs in context to the land areas of the White Mountains-San Francisco Peaks-Mogollon Rim M313A ecoregion



Past, present, and foreseeable forest and woodland management actions on Federal and tribal lands which could contribute to cumulative effects are fire suppression and the lack of thinning trees less than 9 inch diameter that have resulted in an overabundance of small trees with no market value. A similar situation exists on state and private lands.

National forests and State, tribal, and private lands have not been able to institute long-term uneven-aged management practices designed to provide sustainable levels of wood products because adequate markets to purchase small diameter trees have not existed on a consistent basis. Management emphasis has focused on short-term fuels reduction at a cost to the land owner.


Wood volumes cut from State and private lands are less likely to impact the total market situation, as their treatments are smaller, widely scattered across the ecoregion, and less likely to provide long-term wood volumes. Tribes typically utilize their cut volumes in their own industries, although they may supply some to local markets.

However, the White Mountain Apache tribe (Fort Apache Indian Reservation) currently has no active timber cutting program because they no longer have any markets, and their industrial mill is sitting idle (Drury, 2011 and May 2012). Therefore, the bulk of products available to markets come from Federal lands.

Future forest/woodland management strategies across all other national forests within the ecoregion are expected to be similar to those proposed for the Apache-Sitgreaves NFs. They are revising their land management plans or intend to revise their plans in the near future. The other national forests and the Apache-Sitgreaves NFs would use the same desired conditions for the forested and woodland PNVTs, with uneven-aged silviculture and the return of fire and other natural disturbances to their natural roles.

The largest foreseeable action is the Four Forests Restoration Initiative (4FRI), for which the contract was recently awarded. This project includes 2.4 million acres on four national forests (Apache-Sitgreaves, Coconino, Kaibab, and Tonto) in northern Arizona, and it will focus on restoring the ponderosa pine and dry mixed conifer forested PNVTs (see map 2). Several of its landscape scale projects are planned within the White Mountains-San Francisco Peaks-Mogollon Rim Ecoregion. The management actions associated with many future 4FRI projects may be implemented in this planning period. This initiative seeks to develop sustainable markets for wood products as the result of restoring and maintaining desired conditions, which are similar across all four national forests involved.

Map 2. Four Forests Restoration Initiative project scale in Northern Arizona



Logging/thinning industrial operations may need to expand considerably to utilize all the material potentially to be offered annually across the ecoregion (ERI, 2012a; Hunt, 2012). If that does happen, and demand rises more than expected, then it may be possible to treat more acres annually with cuts. In that case, the ASQ might need to be re-evaluated to accommodate a greater early departure above the LTSYC, for the opportunity to reach desired conditions faster. However, if national and local economies/budgets are unable to overcome rising transportation costs and logging/thinning costs, then a glut of wood products may result. This could jeopardize the Apache-Sitgreaves NFs ability to meet maximum treatment acreage objective each year. Should that happen, then return cutting cycles on suitable timberlands would be missed and uneven-aged management strategies would suffer delayed attainment of desired conditions as a negative consequence. More use of controlled fire than originally planned for the chosen alternative could be needed as a primary tree thinning method on suitable timberlands. That would provide its' own consequences of increased smoke outputs, as well as less management control over tree thinning to create/maintain an even progression of age/size classes for long-term sustained yield.

The 4FRI project is likely to become the major instrument to implement **alternatives A, B, or C** treatments on the ponderosa pine and dry mixed conifer forest lands. It has the potential to become the principal market for the majority of logging operations in north and central Arizona during the planning period. Treatments and harvest volumes on the Gila NF would not be included in 4FRI.

Alternative D can provide wood volumes during the planning period, but would fall short of contributing to sustainable markets. It treats so many more acres exclusively with fire, and cuts on several PNVTs for which 4FRI has no focus, thereby limiting the amount of harvestable green wood volumes to support the markets dependent upon the 4FRI collaboration.

At its high objective level, **alternative C** may be moresuited to supply the high volume that 4FRI is expected to demand because the **other alternatives'** cutting volumes may not be high enough. However, **alternative B's** high and low treatment objectives would provide more wood volume to 4FRI than **alternative C's** low objective would.

Alternative A is not expected to provide enough volume for 4FRI because this alternative emphasizes mechanical treatments around communities. Currently, most areas around communities have already been thinned and now only require follow-up maintenance thinning during this planning period. As a result, harvesting may shift to other national forests for more volume, which could pull operators and contractors away from completing restoration work elsewhere on the Apache-Sitgreaves NFs. A Forest Service strategy for 4FRI is to discourage Federal payment for tree cutting and removal services, in favor of returning to selling the cut trees at minimal to fair sale values. This shift in strategy could eliminate some logging/thinning companies from fully participating, and/or reduce the number of sales sold as transportation fuel prices rise. Fewer cutting treatment acres could result, and they would likely be located closer to established markets, and/or to lands nearest to railroad lines.

The proposed 2012 Wallow West Fuel Reduction and Forest Recovery Project proposes to cut standing dead, dying, and hazard trees across 15,402 acres inside the Wallow fire burn perimeter in the next 3 years. It is covered under the existing forest plan. This project would also reforest up to 4,000 acres of tree plantations in the next 10 years. It would only perform these treatments on moderate and high severity burn acres. It would not reduce overgrowth or fuel loadings on the

lightly burned and unburned acres. The fire salvage volumes it cuts will not apply to the ASQ limit of **any alternative**. They will provide additional employment and wood volume to local markets while the White Mountain Stewardship Project concludes, and the first 4-FRI projects are begun on the Coconino and Kaibab NFs during the next 3 years. Green tree restoration thinning for this area would have to be proposed under the new forest plan.

See the Socioeconomic Resources section for additional cumulative environmental consequences.

Unavoidable Adverse Environmental Consequences

The proposed plan provides a programmatic framework that guides site-specific actions but does not authorize, fund, or carryout any project or activity. Therefore, decisions made in the proposed plan do not cause unavoidable adverse environmental consequences. The application of standards and guidelines during future project and activity decision-making would provide resource protection measures and would limit the extent and duration of any adverse environmental impacts.

Irreversible and Irretrievable Commitment of Resources

Irreversible commitments of resources are those that cannot be regained, such as the extinction of a species or the removal of mined ore. Irretrievable commitments are those lost for a period, but could be regained, such as the temporary loss of timber productivity in forested areas kept clear for use as a power line rights-of-way or road.

Because the proposed plan does not directly authorize or mandate any site-specific project or activity (including ground-disturbing actions), none of the alternatives cause an irreversible or irretrievable commitment of resources. Future project-level decisions under any of the alternatives may result in potential irreversible or irretrievable commitments of resources, which would be disclosed accordingly with the project site-specific analysis.

Adaptive Management

The desired conditions, management approaches, standards and guidelines, and **action alternatives** for the proposed plan provide the adaptive framework to help deal with impacts from climate change. They focus on creating and/or maintaining resilient and redundant resource conditions to provide reasonable assurance of the ability to adapt to a changing climate. Each alternative emphasizes some level of return to desired forest/woodland structure, function, species composition, health, fire regime, and other natural disturbance patterns that evolved here during millennia in which variable climate patterns were present.

A consequence of shifting into using more fire as the tree-thinning tool of choice across all action alternatives, would be the critical silvicultural timing needed for repeat fires to occur at least once every 15 years (or sooner), as a tree stocking control method. Cool ground fire often acts as a good seed bed preparation that increases the abundance and survival of natural conifer regeneration. New pine, juniper, and Douglas-fir regeneration that become established on average to highly productive sites can grow rapidly enough into the large sapling/pulp/small pole-size to reach height and/or bark thickness that are less susceptible to cool ground fire (Boehning, 1982-2012; Youtz, 2010b; Jones, 1974; Laughlin et.al, 2011; Roccaforte, Nov. 2012). If we expect fire

to keep the forest thinned out in place of cutting excess trees, then it must occur before those trees are too large for fire to do its intended job. Missing regular fire cycles for any reason (limited burning condition windows, low budgets, workforce shortages, restrictions on smoke production, etc.) jeopardizes reaching and maintaining the desired conditions, so adaptive management would be needed in that case. This consequence would be the least under **alternative C**; and the greatest under **alternative D**, which relies more heavily on using fire across the most treatment acres as the only tool for tree thinning.

If project-level decisions should choose to implement moderate severity fire as the only tree stocking control method on suitable timber lands, then the timberland suitability classification and the ASQ would need to be re-evaluated. The traditional long-term sustained yield concept of “cut = net growth” would no longer be applicable, unless adjusted to account for “cut + burn = net growth”. If the burning program fails to keep up with the needed schedule, then excess trees will need to be thinned by cutting instead, which could require plan amendments to modify the management objectives for any action alternative selected.

During re-evaluation every 10 years, recalculation of suitable timberland acres and ASQ may need to consider shifts of tree species ranges across the landscape onto different areas, and tree growth rate changes in response to shifts in climate. The LTSY may need to be recalculated as well. Timber growth and yield models and landscape level vegetation models would have to be recalibrated for these new calculations. New markets to utilize more abundant species would likely develop. If certain tree species or sizes become rare and in higher demand, they may bring high enough market prices to offset currently cost-prohibitive cable or helicopter harvesting operations.

Another component of the proposed plan is the chapter on monitoring strategy. Information from monitoring items listed there will enable changes needed in the plan for adaptive management.

Other Planning Efforts

Several other planning efforts were previously described in the Cumulative Effects section.

Although the proposed plan (**alternative B**) emphasizes addressing the needs of communities at risk of catastrophic fire, it is not entirely consistent with the Community Wildfire Protection Plans (CWPPs) for Apache, Navajo, Coconino and Greenlee counties, which were published in 2004 and 2005. These CWPPs include a generic prescription to “thin from 40 to 60 BA, with a 16-inch diameter cap” on Federal lands, which may not move project areas toward the land management plan’s desired conditions. As designed, **alternative C** would be inconsistent with such a prescription, while **alternative D** would adopt the 16-inch diameter cap for nearly all cuts. **Alternative A** may continue to use this prescription on many treatment acres. Under **all alternatives**, this generic prescription would not automatically be proposed or adopted at the project level. However, the CWPP prescription will need to be considered as an alternative analyzed in detail under any project (**regardless of plan alternative**) proposed under the Healthy Forests Restoration Act of 2003. Even though **Alternatives B and C** were designed to have no 16” diameter cap cuts, they would still implement the 24” diameter cap required on Restricted habitat acres, and the 9” diameter cap required on Protected habitat acres, for compliance with the current Mexican Spotted Owl Recovery Plan (USDI-FWS, 1995).

No other potential conflicts are evident from this analysis between the proposed action and the objectives of Federal, regional, State, local, or Tribal land use plans, policies, and controls for the area concerned.

References

The following information sources were reviewed and/or consulted for this analysis.

- Abella, Scott R., Peter Z. Fule, and W. Wallace Covington. December 2006. Diameter caps for thinning Southwestern ponderosa pine forests: Viewpoints, effects, and tradeoffs. *Journal of Forestry*, 104:407-414. Society of American Foresters.
- Arno, Steven F. and Carl E. Fiedler, 2005. *Mimicking Nature's Fire: Restoring Fire-Prone Forests in the West*. Island Press, Washington DC. ISBN 1-55963-143-0. 242 pp. [www.islandpress.org]
- Alexander, Robert R. 1974. *Silviculture of subalpine forests in the central and southern Rocky Mountains: The status of our knowledge*. USDA Forest Service Research Paper RM-121, 88p. Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO 80521
- Boehning, Monica. 1982 to 2012. Professional observations during work and field visits across the Apache-Sitgreaves National Forests, while employed in various forestry and silviculture positions, including: timber compartment/stand delineations and field mapping; stand examinations and diagnosis; treatment prescription development; timber marking and cruising; cutting unit boundary layout; area determination; old growth surveys; wildfire detection, prevention patrols, and suppression; prescribed burning; insect/disease identification and monitoring; thinning contract preparation and administration; wildlife habitat analysis; NEPA analysis; pre-treatment and post-treatment monitoring comparisons; reforestation field-assessments; hosting academic researchers; coordinating with adjacent landowners and other natural resource management agency personnel; and familiarity/interpretation of the current forest land management plan.
- Covington, W. Wallace, Peter Z. Fule, Margaret M. Moore, Stephen C. Hart, Thomas E. Kolb, Joy N. Mast, Stephen S. Sackett, and Michael R. Wagner. 1997. Restoring ecosystem health in ponderosa pine forests of the southwest. In *Journal of Forestry*: Vol.95, No.4, April, 1997: pp.23-29. Society of American Foresters, Bethesda, MD.
- Cram Douglas S., Terrell T. Baker, and Jon C. Boren. February 2006. Wildland fire effects in silviculturally treated vs. untreated stands of New Mexico and Arizona. Research Paper RMRS-RP-55. USDA Forest Service, Rocky Mountain Research Station, Fort Collins, CO. 28 pp.
- Drury, Jerry. April 2011. Several in-office professional conversations with the Apache-Sitgreaves Natl. Forest Timber Program Manager/Timber Contracting Officer. Forest Supervisor's Office, Springerville, AZ; and past BIA Forester with the Fort Apache Indian Reservation.

- Drury, Jerry. February 14, 2012. Professional correspondence - email regarding current White Mountain Stewardship trend in annual acreage treatment under existing forest plan.
- Drury, Jerry. May 2, 2012. Professional in-office communication with the Apache-Sitgreaves Natl. Forests Natural Resources Staff Officer regarding fire salvage, product values, local wood market demands and limitations.
- ERI, 2007. Case study of a community stewardship success: The White Mountain Stewardship Contract. Ecological Restoration Institute. Northern Arizona University, Flagstaff, AZ. www.eri.nau.edu.
- ERI. March 2011. Fact Sheet: Post-wildfire fuels and regeneration dynamics – What to expect following severe wildfires in ponderosa pine forests. Ecological Restoration Institute. Northern Arizona University, Flagstaff, AZ. www.eri.nau.edu.
- ERI, 2012a. Workforce needs of the Four Forest Restoration Initiative Project: An analysis. Ecological Restoration Institute. Northern Arizona University, Flagstaff, AZ. www.eri.nau.edu.
- ERI. 2012b. White paper: Reforestation following severe fires. Ecological Restoration Institute. Northern Arizona University, Flagstaff, AZ. www.eri.nau.edu.
- Fairweather, Mary Lou. June 26, 2008. Insect and Disease Activity in Hall Ranch WUI, Springerville RD. File Code 3420, Project Report by Forest Pathologist, USFS SW Region Forest Health, AZ Zone Office, Flagstaff.
- Feidler, Carl E., Kerry L. Metlen, and Erich K. Dodson. 2010. Restoration treatment effects on stand structure, tree growth, and fire hazard in a ponderosa pine/douglas-fir forest in Montana. *Forest Science* 56(1) 2010, the Society of American Foresters.
- Friederici, Peter G. 2006a. ERI Fact Sheet: Forest Restoration Treatments and Fire Behavior. Ecological Restoration Institute, Northern Arizona University, Flagstaff, AZ.
- Friederici, Peter G. 2006b. ERI Fact Sheet: Restoring the Ecological and Social Integrity of Western Forests. Ecological Restoration Institute, Northern Arizona University, Flagstaff, AZ.
- Fule, Peter Z., Allison E. Cocke, Thomas A. Heinlein, and W. Wallace Covington. 2004. Effects of an intense prescribed forest fire: Is it ecological restoration? *Restoration Ecology* Vol. 12 No. 2, pp. 220-230. June 2004.
- Higgins, Bruce J. June 2008. White paper: A comparative analysis of forest recovery after uncharacteristic fire in ponderosa pine forest. USDA Forest Service, Kaibab National Forest. Williams, AZ. 17pp.
- Hunt, Scott. April 27, 2012. Conference to focus on forest economy. New article by the Arizona State Forester. Published in White Mountain Independent Newspaper, Show Low, AZ.

- Jones, John R. 1974. Silviculture of southwestern mixed conifers and aspen: The status of our knowledge. USDA Forest Service Research Paper RM-122, 44p. Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO 80521
- Joyce, L., J. Aber, S. McNulty, V. Dale, A. Hansen, L. Irland, R. Neilson, and K. Skog. (2001). Potential consequences of climate variability and change for the forests of the United States. Chapter 19, Pp. 489-522. In: Climate change impacts on the United States: The potential consequences of climate variability and change. National Assessment Synthesis Team (ed.). A Report for the US Global Change Research Program. Cambridge University Press. New York, NY.
- Korb, Julie E., Peter Z. Fule, Michael T. Stoddard. 2012. Forest restoration in a surface fire-dependent ecosystem: An example from a mixed conifer forest, southwestern Colorado, USA. *Forest Ecology and Management* 269 (2012) 10-18.
- Laing, L., Ambos, N., Subirge, T., McDonald, C., Nelson, C., and Robbie, W. (1987). Terrestrial Ecosystem Survey of the Apache-Sitgreaves National Forests. U.S. Forest Service, Southwestern Region, Albuquerque, NM. Pp 453.
- Laughlin, D.L., J.P. Roccaforte, and P.Z. Fulé. 2011. Effects of a second-entry pre-scribed fire in a mixed conifer forest. *Western North American Naturalist* 71(4):557-562.
- Logan Simpson Design, Inc. (2004a). Community Wildfire Protection Plan for At-Risk Communities of the Sitgreaves National Forest in Apache, Coconino, and Navajo Counties. Tempe, Arizona.
- Logan Simpson Design, Inc. (2004b). Community Wildfire Protection Plan for At-Risk Communities of the Apache National Forest in Apache County. Tempe, Arizona.
- Logan Simpson Design, Inc. (2005). Greenlee County Community Wildfire Protection Plan for At-Risk Communities of the Apache National Forest in Greenlee County. Tempe, Arizona.
- Lynch, Ann M., John A. Anhold, Joel D. McMillin, Steve M. Dudley, Roberta A. Fitzgibbon, Mary Lou Fairweather. February 2010. Forest insect and disease activity on the Apache-Sitgreaves N.F., and Fort Apache Indian Reservation, 1918-2009: Report for the Apache-Sitgreaves N.F./Regional Analysis Team. USDA Forest Service Rocky Mountain Research Station, Tucson AZ; and Arizona Zone Office Forest Health Protection, USFS, Flagstaff, AZ.
- O'Brien, Renee A. Jan. 2002. Arizona's Forest Resources, 1999. Resource Bulletin RMRS-RB-2. USDA-FS Rocky Mountain Research Station, Ogden, UT.
- Puhlick, Joshua J., Daniel C. Laughlin, Margaret M. Moore. 2012. Factors influencing ponderosa pine regeneration in the southwestern USA. *Forest Ecology and Management* 264(2012) 10-19. Elsevier B.V. [www.elsevier.com/locate/foreco]
- Puettmann, Klaus J. 2011. Silvicultural challenges and options in the context of global change: "simple" fixes and opportunities for new management approaches. *Journal of Forestry*, September 2011 issue. Society of American Foresters.

- Richardson, Gayle. 2012. Professional communications with the Black Mesa Ranger District Silviculturist (and former Alpine/Springerville Districts Forester), regarding reforestation success in general on different soils, and post-fire reforestation needs on the Apache-Sitgreaves Natl. Forests, including post-burn reforestation needs reported in the USFS FACTS Database.
- Roccaforte, John Paul. November 2012. ERI Fact Sheet: Assessing Restoration Objectives Following a Second-entry Prescribed Fire in an Unharvested Mixed Conifer Forest. Ecological Restoration Institute, Northern Arizona University, Flagstaff, AZ.
- Roccaforte, John P., Peter Z. Fule, W. Walker Chancellor, and Daniel C. Laughlin. 2012. Woody debris and tree regeneration dynamics following severe wildfires in Arizona ponderosa pine forests. *Can. J. For. Res.* 42: 593-604. NRC Research Press. www.nrcresearchpress.com
- Rogers, Paul. Sept. 2003. Forest Resources of the Apache-Sitgreaves National Forest. USDA-FS Rocky Mountain Research Station, Ogden, UT.
- Rogers, Paul C. (ed.) June 12, 2008. Summary and Abstracts from Sudden Aspen Decline (SAD) Meeting. Compilation of case studies presented at the Ft. Collins, CO meeting of SAD researchers held Feb. 12-13, 2008.
- Rogers, Paul C. Oct. 24, 2011. Letter to USFS Southwest Regional Forester summarizing post-Wallow Fire aspen conditions and regeneration monitoring field visit (under USFS Region 3 Forest Health Cooperative Agreement No. 11-PA-11031600-080); by Director of Western Aspen Alliance and professor at Utah State University Wildland Resources Department, Logan Utah. (<http://www.western-aspen-alliance.org/>)
- Ryan, M., S. Archer, R. Birdsey, C. Dahm, L. Heath, J. Hicke, D. Hollinger, T. Huxman, G. Okin, R. Oren, J. Randerson, and W. Schlesinger. (2008). Land resources. Chapter 3, Pp. 75-121. In: The effects of climate change on agriculture, land resources, water resources, and biodiversity. A Report by the U.S. Climate Change Science Program and the Subcommittee on Global Change Research, Washington, DC.
- Savage, Melissa and Joy Nystrom Mast. 2005. How resilient are southwestern ponderosa pine forests after crown fires? *Can. J. For. Res.* 35: 967-977. NRC Research Press.
- Shafer, John. circa 1993. White paper by the Forest Silviculturist entitled "Apache-Sitgreaves National Forests Timber Program Re-assessment – Ecosystem Management"
- Shepperd, W.D. and M.L. Fairweather. 1994. Impact of Large Ungulates in Restoration of Aspen Communities in a Southwestern Ponderosa Pine Ecosystem. In: Covington W.S.; DeBano L.F. (eds.), Sustainable Ecological Approach to Land Management. Gen. Tech. Rep. RM-247. Fort Collins, CO: USDA FS, Rocky Mountain Forest and Range Experiment Station: pp. 344–347.
- Sitko, S. and S. Hurteau. 2010. Evaluating the Impacts of Forest Treatments: The First Five Years of the White Mountain Stewardship Project. The Nature Conservancy, Phoenix AZ.

- Sprigg, W. A., T. Hinkley, et al. (2000). Preparing for a Changing Climate: The Potential Consequences of Climate Variability and Change: Southwest. A Report of the Southwest Regional Assessment Group. University of Arizona. The Institute for the Study of Planet Earth, Tucson, AZ, US Global Change Research Program: 66.
- Schwilk, Dylan, W., Jon E. Keeley, Eric E. Knapp, James McIver, John D. Bailey, Christopher J. Fettig, Carl E. Fiedler, Richy J. Harrod, Jason J. Moghaddas, Kenneth W. Outcalt, Carl N. Skinner, Scott L. Stephens, Thomas A. Waldrop, Daniel A. Yaussy, and Andrew Youngblood. 2009. The national Fire and Fire Surrogate study: effects of fuel reduction methods on forest vegetation and fuels. *Ecological Applications*, 19(2), 2009, pp.285-304. Ecological Society of America.
- Stephens, Scott L., Jason J. Moghaddas, Carl Edminster, Carl E. Fiedler, Sally Haase, Michael Harrington, Jon E. Keeley, Eric E. Knapp, James D. McIver, Kerry Metlen, Carl N. Skinner, and Andrew Youngblood. 2009. Fire treatment effects on vegetation structure, fuels, and potential fire severity in western U.S. forests. *Ecological Applications*, 19(2), 2009, pp.305-320. Ecological Society of America.
- Strom, Barbara A. and Peter Z. Fule. 2007. Pre-wildfire fuel treatments affect long-term ponderosa pine forest dynamics. *International Journal of Wildland Fire*, 2007, **16**, 128-138.
- Triepke, F.J., B. J. Higgins, R. N. Weisz, J. A. Youtz, and T. Nicolet. November 2011. Diameter caps and forest restoration - Evaluation of a 16-inch cut limit on achieving desired conditions. USDA Forest Service Forestry Report FR-R3-16-3. Southwestern Region, Regional Office, Albuquerque, NM. 31 pp.
- U.S. Department of Interior, Fish and Wildlife Service (USDI-FWS). (1995). Recovery Plan for the Mexican Spotted Owl (*Strix occidentalis lucida*). Vol. 1. Albuquerque, NM. 172 pp.
- U.S. Forest Service. (July, 2008). Forest plan revision resource evaluations: Apache-Sitgreaves National Forests. U.S. Department of Agriculture, Forest Service Southwestern Region, Springerville, AZ.
- U.S. Forest Service. (August, 2008). Assessment of Vegetation Diversity and Risks to Ecological Sustainability: Vegetation specialist's report. Apache-Sitgreaves National Forests. U.S. Department of Agriculture, Forest Service Southwestern Region, Springerville, AZ.
- U.S. Forest Service. (December, 2008). Comprehensive evaluation report (CER): Apache-Sitgreaves National Forests. U.S. Department of Agriculture, Forest Service Southwestern Region, Springerville, AZ.
- U.S. Forest Service. (December 2008). Ecological sustainability report (ESR): Apache-Sitgreaves National Forests. U.S. Department of Agriculture, Forest Service Southwestern Region, Springerville, AZ.
- U.S. Forest Service. (January 2009). Apache-Sitgreaves National Forests Economic and social sustainability assessment. Apache-Sitgreaves National Forests. U.S. Department of Agriculture, Forest Service Southwestern Region, Albuquerque, NM.

- Weisz, Reuben, D. Vandendriesche, M. Mouer, M. Boehning, L. Wadleigh, J. Triepke, M. White, C. Nelson, J. Palmer, J. Youtz, B. Higgins, T. Nicolet, P. Bostwick, D. Mindar, M. Pitts, M. Manthei, W. Robie. 2012. White Paper O: Process overview of using FVS to create VDDT models. USDA Forest Service, Southwestern Region, Regional Office, Albuquerque, NM. Interoffice publication.
- White, Alan S. 1985. Presettlement regeneration patterns in a southwestern ponderosa pine stand. *Ecology*: 66(2), pp. 589-594. Ecological Society of America.
- Williams, A. Park, Craig D. Allen, Alison K. Macalady, Daniel Griffin, Connie A. Woodhouse, David M. Meko, Thomas W. Swetnam, Sara A Rauscher, Richard Seager, Henri D. Grissino-Mayer, Jeffrey S. Dean, Edward R. Cook, Chandana Gangodagamage, Michael Cai, and Nate G. McDowell. September 2012. Temperature as a potent driver of regional forest drought stress and tree mortality. *Nature Climate Change*, Macmillan Publishers Limited. 6pp. (www.nature.com/natureclimatechange)
- Youtz, James. 2010a. Poster Presentation entitled: "Post-Treatment Comparison of R3 Desired Condition (Eagar South) Versus Thinning from Below Diameter Limit (Mineral WUI) Apache-Sitgreaves National Forest." Electronic copy of 22 x 34 inch color landscape orientation poster when plotted on paper. Shows photos and bar graphs of Mineral WUI 2005 pre-treatment and 2009 post-treatment plot data comparison of diameter limit cut treatment. Prepared by the Regional Silviculturist and GIS Specialists, dated 03/15/2010. USDA Forest Service, Southwestern Region, Albuquerque, NM.
- Youtz, James A. 2010b. Professional communication during June 4, 2010 USFS Region-3 tele-video-conference with local fire researchers, fuels, and forestry practitioners, about using fire as the only tree thinning tool. Written notes recorded by M.Boehning, in attendance.
- Youtz, James A, and Don Vandendriesche. February, 2010. Professional conversations (teleconference calls) with the Regional Silviculturist and USFS FMSC Biometrician, re: FVS and VDDT modeling work done by the USFS Region 3 FVS Timber Calculations Working Group for plan revision, including LTSYC.
- Youtz, James A, and Don Vandendriesche. September, 2012 (revised version of March 2011 document with same title). White paper entitled: National Forest Planning and Sustained Yield of the Timber Resource Long-Term Sustained-Yield Calculations for Forest Land and Resource Management Planning. USDA Forest Service, Southwestern Region, Albuquerque, NM, and Washington Office Forest Management Service Center. 39 pp.

Credits

Major supporting contributors to this report are:

- Nancy Loving, Forest GIS Specialist
- Chris Nelson, Forest Soils Scientist (timber suitability)
- Mitchel White, Forest Ecologist (VDDT model cutting volumes, some climate change info)
- Deborah MacIvor, Forest Engineer (road cost calculations in economic efficiency study)
- Ray Rugg, Apache Zone Timber Staff (logging operability mapping)
- James Youtz, Southwestern Region Silviculturist (advice, review, and edits)
- Jerry Drury, Apache-Sitgreaves Timber Program Mngr./Natural Resources Staff Officer (review and edits)

Appendices - Forest Products

Due to the size and nature of these 18 appendices, they are available in the Planning Record Set of Documents as separate electronic files. They are listed below as an index.

Appendix A1: 1992 WO amendment to FSH 2409.13, zero code. (5 page Word doc.)

Appendix A2: Timber Suitability Analysis Procedures and Alternative Tables (12 page Word doc.) By C. Nelson, N. Loving, M.Boehning, M.Davalos. Finalized August, 2012.

Appendix A3: Suitable Timberland Logging Cost Efficiency Analysis (Excel Spreadsheet with “Read Me” tab and 12 worksheet tabs). By M. Boehning and D.MacIvor, March 2012.

Appendix A3Roads: Road Maintenance, Reconstruction and New Construction Costs Developed for the Suitable Timber Lands Economic Assessment Worksheet. By Forest Engineer D. MacIvor, March 2012. (9 page .pdf text doc.)

Appendix A4: Suitable Timberland Maps, for Alternatives A, B, & C (6 .jpg image maps by N.Loving, May 2012)

Appendix B1: Surrogate Methods of Prescribed Cut and/or Burn Adopted by Region-3 for Using the Forest Vegetation Simulator (FVS) Model to Calibrate the VDDT Model. By M. Boehning/Region-3, April 2011 through Oct. 2012. (9 page Word doc.)

Appendix B2: Silvicultural Rationale for Different Timber Management Approaches and Cutting Mixes by Each ASNFs Alternative for Modeling in VDDT. By M. Boehning, February, 2012. (4 page Word doc.)

Appendix B3: Example PNVT Percentages of Surplus and Deficit Structural States, Differences between Current and Desired Conditions. By M. White and USFS Region-3 (2 page Word doc./diagram)

Appendix B4: USFS Region-3 FVS Model Prescribed Treatment Resulting State Transitions for Calibrating VDDT. By Region-3 Plan Revision Timber Working Group. Assembled for display by M. Boehning, 5/6/12. (Excel spreadsheet with a “Read-Me” tab and 5 large worksheet tables.)

Appendix B5: Apache-Sitgreaves NFs Silviculture Cutting and Planting Prescriptions to Model in VDDT for Suitable and Non-suitable Timberlands, by Alternative and PNVT. Master File-folder, with 4 sub-folders inside (1 folder for each alternative) = total of 31 separate electronic files.

Appendix C: ASNFs Master Acreage Tables for Treatments to Model in VDDT on Suitable, Non-Suitable, and Fire-Only Lands, by PNVT and Alternative. By C.Nelson, 2/10/12. (2 spreadsheets)

Appendix D1: Sample live green tree raw cutting volumes table from VDDT. By M.White, Sept. 2012. (2 large tables, using Alternative B, PPF PNVT, Suitable lands, Decade 1, High and Low options as examples)

Appendix D2: Cutting Volume Spreadsheets of all wood products from VDDT, by PNVT, product type, and Alternative. By M.Boehning, 3/21/12 through 9/26/12. (large Excel spreadsheet with 5 worksheet tables)

Appendix E1: Acreages of treatment methods modeled in VDDT, by alternative. By ASNFs Plan Revision Team, M.Boehning and M. Davalos. Updated 5/1/12. (1 large Excel spreadsheet)

Appendix E2: Acres by treatment type used to model the low and high annual treatment objectives in VDDT. By ASNFs Plan Revision Team and M. Davalos, May 2012. (4 page Word table)

Appendix F: Youtz, James A, and Don Vandendriesche. September 2012 (revised version of March 2011 document with same title). White paper entitled: National Forest Planning and Sustained Yield of the Timber Resource Long-Term Sustained-Yield Calculations for Forest Land and Resource Management Planning. USDA Forest Service, Southwestern Region, Albuquerque, NM, and Washington Office Forest Management Service Center. 39 pp.

Appendix G: Sample USFS Region-3 FVS model results of pre-cut, harvest, and post-cut tree species composition changes for the Free Thinning, Diameter Cap, and Group Selection cutting methods used in VDDT (using the Dry Mixed Conifer PNVT, veg. structural state I as one example.) FVS Model runs done by Region-3 in 2010-2011. Results of 3 simulations assembled for display by M.Boehning in May, 2012. (16 page Word doc. with color emphasis)

Appendix H: USFS Forest Inventory Data Online (FIDO) website queries of ASNFs years 1985/1999-2008/2009 FIA plot data showing mortality rates of large trees due to causes other than fire, insects, disease or animals (Website homepage screen capture + 4 large tables, double-sided). Queries run by M. Boehning on 5/17/2011. (10 page Word doc)