

SALT CREEK WATERSHED ANALYSIS

APPENDIX F: Wet and Special Habitat Types by Subwatershed

Habitat Feature	Acres	Subwatershed
MB, MW, MS	75.8	20 3
MB, MW, MM	12.2	20 2, 20 3
MB, MW, MS, MM	90.4	20 2, 20 3
MP	0.7	20 2
MP, WS	62.1	20 2
MS	68.2	20 1, 20 2, 20 3
MS, MW	2.1	20 3
MS, WP, UC, RC, GD, RT, HD, RO	4.5	20 1
MS, WR, MP, HD, MW	8.5	20 1
MW	151.7	20 2, 20 3
MW, HD	2.2	20 1
MW, MB	32.5	20 3
MW, MM	255.3	20 2, 20 3
MW, MM, SA	6.1	20 2
MW, MS	9.7	20 1, 20 2
MW, MS, MM	7.6	20 2
MW, MS, MM, MB	70.6	20 2
WH, MS	59.9	20 1
WP	147.9	20 1, 20 2, 20 3
WP, MM	0.4	20 3
WP, MS	32.0	20 1, 20 2
WP, MS, MP, HD, MM, MW	15.7	20 1
WP, WT, MM	0.7	20 2
WS	120.7	20 2
WT	102.8	20 2
WT, RO	6.4	20 1
WT, RT, ST, MD, SA	10.6	20 2
WT, WS	174.3	20 2
WX	477.0	20 1, 20 2, 20 3

HD = Hardwood inclusion
 MB = Bog
 MD = Dry meadow
 MM = Mesic meadow
 MS = Sedge meadow
 MP = Swamp
 MW = Wet meadow
 RC = Cliff
 RO = Rock outcrop
 RT = Talus
 GD = Dry rock garden
 SA = Sitka alder
 ST = Vine maple (talus)
 UC = Cave
 WH = Headwater
 WP = Pond
 WS = Seep
 WT = Small stream
 WX = Open water

SALT CREEK WATERSHED ANALYSIS

APPENDIX G: Herpetile Habitat

Species	Life History Stage	Habitat	Elevation Range (ft)	Prey	Predators
Tailed Frog (<i>Ascaphus truei</i>)	Adult & Larvae	In fast flowing permanent stream; adults in/near cold, clear streams. Headwaters	0-6562	Adults: snails, ticks spiders, mites, insects. Larvae: diatoms, conifer pollen, algae, small insects	Pacific Giant Salamander
Western Toad (<i>Bufo boreas</i>)	Adult & Larvae	Large lakes, small ponds, shallow marshes; adults: forested areas possibly away from water	0-11812+	Flying insects, crayfish, sowbugs, earthworms	Birds, garter snakes, aquatic insects
Pacific Chorus Frog (<i>Pseudacris Regilla</i>)	Adult & Larvae	Variety: coastal sloughs, old-growth, deserts; adults quite terrestrial	0-9843+	Beetles, flies, ants, spiders, isopods, various insects	Garter snakes, bullfrogs, birds, mammals. Larvae: aquatic insects, salamanders
Red-Legged Frog (<i>Rana aurora</i>)	Adult & Larvae	Permanent bodies of quiet water, ponds, pools, reservoirs, springs, lakes, marshes; adults hang out on land near water's edge	<2789	Many insect species, arachnids, mollusks	Snakes, raccoons, herons, NW Salamander, bullfrogs, Roughskin Newts, cutthroat trout, owls, hawks, ducks, skunks, minks, cats. Larvae: giant water bugs, Dytiscid beetles, Odonate nymphs...
Foothills Yellow-Legged Frog (<i>Rana boylei</i>)	Adult & Larvae	Near streams & rivers w/rocky or gravelly substrate	<5906	Insects, snails	Garter snakes, fish, birds, mammals. Eggs: Roughskin Newts
Cascades Frog (<i>Rana cascadae</i>)	Adult & Larvae	Montane meadows, marshes, ponds, relatively small water bodies, along creeks, lakeshore alcoves	>2625	Aquatic insects	salamanders, fish, snakes, birds
Western Spotted Frog (<i>Rana pretiosa</i>)	Adult & Larvae	Marshes, near edges of ponds and lakes, colder waters	?-9843	Insect species, arachnids, mollusks	Garter snakes
Western Aquatic Garter Snake (<i>Thamnophis couchii</i>)	Adults & Neonates	Permanent streams w/rocky substrate, stream margins	0-8006	Fish, amphibians	Otters, herons, hawks, osprey, Steller's jay
Western Pond Turtle (<i>Clemmys mammorata</i>)	Adults & Juveniles	Marshes, sloughs, lakes, ponds, slow portions of creeks/rivers.	0-6004	Various aquatic inverts/verts., algae, cattail/tule roots	Adults/Eggs: Otters, raccoons, coyotes, foxes. Juveniles: Bullfrogs, bass
Northwestern Salamander (<i>Ambystoma gracile</i>)	Terrestrial (Adult) Aquatic (Adult/Larvae)	Coniferous forests, inland valleys, subalpine areas Ponds, lakes, slow parts of semi-permanent streams	<10171	Larvae: aquatic inverts	Trout, aquatic beetle larvae
Roughskin Newt (<i>Taricha Granulosa</i>)	Terrestrial (Adult) Aquatic (Adult/Larvae)	Farmland, grassland, uplands, forests. Ephemeral/permanent ponds/lakes/streams w/slow areas with veg.	0-9187	Small inverts. & verts, amphib eggs & larvae Zooplankton, aquatic insects	Few predators due to skin toxicity, some predation by trout
Pacific Giant Salamander (<i>Dicamptodon tenebrosus</i>)	Terrestrial (Adult) Aquatic (Adult/Larvae)	Moist conifer forest, in/near streams, talus roadcuts. High gradient streams w/coarse substrate	0-7087	Terrestrial invertebrates & vertebrates. Aquatic inverts. & vertebrates	Fish, weasels, shrews, other Pacific Giant Salamanders, Western Aquatic Garter snakes.
Clouded Salamander (<i>Aneides ferreus</i>)	Terrestrial (Adult/Larvae)	Rock faces (talus), 2 different age classes of large downed logs, often Douglas fir	0-5578	Anthopods, ants	--
Oregon Slender Salamander (<i>Batrachoseps wrighti</i>)	Terrestrial (Adult/Larvae)	Mature fir forests, rotting logs/stumps, wood fragments, substrate temp 51-570 Fahrenheit	49-4397	--	--

SALT CREEK WATERSHED ANALYSIS

Herpetile Habitat CONTINUED

Species	Life History Stage	Habitat	Elevation Range (ft)	Prey	Predator
Ensatina (Ensatina eschscholtzii)	Terrestrial (Adult/Larvae)	Conifer/deciduous forests of diff. ages, under surface debris, decaying logs, small mammal burrows	-	Invertebrates	Garter Steller's jay
Dunn's Salamander (Plethodon dunni)	Terrestrial (Adult/Larvae)	Semi-aquatic, associated w/rocks, moss covered rubble and seeps. Under rocks/logs, moist talus	0-3281 absent in most Willamette Valley	-	-
Western Red-Backed Sal. (Plethodon vehiculum)	Terrestrial (Adult/Larvae)	Coniferous forests, soft shale/sandstone outcrops, decaying logs, bark piles. Damp soils rather than wet situations	-	-	-
Long-Toed Salamander (Ambystoma macrodactylum)	Terrestrial (Adult) Aquatic (Adult/Larvae)	Lowland forest, disturbed pastures, high elevation lakes/ponds. Lakes/ponds, temporary water sources	0-8120	Larvae: variety of invertebrates, copepods, fairy shrimp, young Chorus Frog tadpoles, smaller NW Sal. larvae	Introduced fish species

*Adapted from Blaustein et al., 1995; except for Long-Toed Salamander which is from Leonard et al., 1993.

NOTE: Shading indicates species known to occur in the Salt Creek watershed

Amphibian Distribution and Sensitivity in the Westslope Cascades Province

Species	Occurrence In Province	ODFW Status	ONH Status	BLM Status	USFS Status	USFWS Status
Northwestern Salamander	throughout					
Long-Toed Salamander	throughout					
Roughskin Newt	throughout					
Cope's Giant Salamander	few localities	Pr-S/u	2	AS		
Pacific Giant Salamander	throughout					
Cascade Torrent Salamander	few localities	Pr-S/v	4	TS		
Clouded Salamander	throughout	S/u	3			
Oregon Slender Salamander	few localities	S/u	1			
Ensatina	throughout					
Dunn's Salamander	throughout					
Larch Mountain Salamander	few localities	S/v	3	ROD	ROD	C2
W. Red-Backed Salamander	few localities					
Western Toad	few localities	S/v				
Pacific Chorus Frog	throughout				S	
Tailed Frog	throughout	Pr-S/v	4	AS		
Red-Legged Frog	throughout	S/u	4	TS	S	C2
Cascades Frog	throughout	Pr-S/v	3	AS		
Spotted Frog	few localities	Pr-S/c	2	BS		C1
Foothill Yellow-Legged Frog	few localities	Pr-S/v	4	TS		C2
Bullfrog	few localities					

Key to the Different Status Codes:

(from Corkran and Thoms, 1995)

ODFW (Oregon Department of Fish and Wildlife)

OHN (Oregon Natural Heritage Database)

Pr = Protected

c = Critical

1 = Threatened throughout range 3 = Review

S = Sensitive

v = Vulnerable

2 = Threatened in Oregon only

p = Peripheral, naturally rare u = undetermined status

BLM (Bureau of Land Management) and USFS R6

USFWS (U.S. Fish & Wildlife Service)

TS = Tracking Species

AS = Assessment Species

C1 = Candidate, sufficient information

S = USFS Sensitive Species

BS = BLM Sensitive Species

C2 = Candidate, insufficient information

ROD = Record of Decision for Amendments to USFS and BLM Planning Documents within Range of the Northern Spotted Owl, April 1994; survey & manage species

APPENDIX H: Aquatic Habitat and Species Lists

Streams surveyed in the Salt Creek Watershed

Stream	Subwatershed	Miles surveyed	Reference Survey Year	Current Survey Year	Method
Salt Creek	20 1, 20 2, 20 3	18.9, 27.5, 19.93, 5.3	1937	1964, 1990, 1996	?, ODFW, Reg, Reg
Tumble	20 1	spot check @ mouth		1964	ODFW
Sage	20 1	spot check @ mouth		1964	ODFW
Pepper	20 1	spot check @ mouth		1964	ODFW
Sugar	20 1	spot check @ mouth		1964	ODFW
Gobel	20 1	0.1		1964	ODFW
Basin	20 1	spot check @ mouth		1964	ODFW
Fin Roberts	20 1	spot check @ mouth		1964	ODFW
Warner	20 1	mouth, 1.9		1964, 1996	ODFW, Reg
McCredie	20 1	mouth, 1.33		1964, 1992	ODFW, Reg
T20e 36.7	20 1	at least 0.4		1992	Reg
T20e 38.6	20 1	N/A		1992	Reg
T20f 41.3	20 1	at least 0.16		1992	Reg
T20f 41.5	20 1	0.8		1992	Reg
T20f 43.3	20 1	0.2		1992	Reg
T20f 45.3	20 2	0.1		1992	Reg
T20f 46.3	20 2	1.1		1992	Reg
T20w 11.8	20 2	1.3		1992	Reg
Eagle	20 2	0.9, 3.5, 5.3	1937	1964, 1992	?, ODFW, Reg, Reg
Coyote	20 2	mouth, 1.4		1964, 1992	ODFW, Reg
S. Fk. Salt	20 2	3.9, 7.3		1964, 1996	ODFW, Reg
Fall	20 2	4.3		1996	Reg
Diamond	20 2	2.5		1996	Reg
Deer	20 3	2.05		1996	Reg

NOTE: For Mainstem Salt Ck., apparently 32 miles were surveyed by the Oregon State game commission in 1970 - no data are available

Observed Pool Values and Objectives for Mainstem Salt Creek (1937, 1964, 1990, 1996 Data)

Stream Name	Reach No.	Reach Length (miles)	Gradient (%)	Average Width (feet)	Residual Depth (feet)	Width/Depth Ratio	% Area in Pools	Observed Pools/mile	Min. Obj. PACFISH	Min. Obj. Forest Plan
Salt Creek 1937	Station A-B	5.7	1.97	52*	---	---	---	15.3	26	14
	Station B-C	5.5	1.47	77*	---	---	---	12.2	23	12
	Station C-D	1.8	2.47	33*	---	---	---	20.6	35	34
	Station D-E	1.9	13	29*	---	---	---	7.4	45	14
	Station E-F	4	5.77	35*	---	---	---	20	40	30

* Widths are not averages for 1937 but rather assumed to be single measurements at each station (A - F)

Station A = confluence of Salt Creek with Middle Fork of Willamette River

Station B = 2nd bridge above confluence

Station C = Eagle Creek confluence with Salt Creek

Station D = South Fork Salt Creek confluence with Salt Creek

Station E = 1.5 miles upstream of UPRR trestle

Station F = Salt Creek Falls (1 mile from Abernathy)

SALT CREEK WATERSHED ANALYSIS

Aquatic Habitat and Species Lists CONTINUED

Observed Pool Values and Objectives for Mainstem Salt Creek Continued

Stream Name	Reach No.	Reach Length (miles)	Gradient (%)	Average Width (ft)	Residual Depth (ft)	Width/Depth Ratio	% Area in Pools	Observed Pools/Mile	Minimum Objective PACFISH	Minimum Objective Forest Plan
Salt Creek 1964	1	5	1 to 6	40	----	----	----	16**	35	----
	2	5.5	1 to 6	20	----	----	----	11**	56	----
	3	6.9	4 to >6	20	----	----	----	11**	56	----
	4	5	4 to >6	15	----	----	----	11**	70	----
	5	3.5	4 to 6	30	----	----	----	8**	45	36
	6	1.6	4 to 6	9	----	----	----	16**	115	119
Salt Creek 1990 (lower only)	1	6.9	2	50.1	4.6	8.5	2	2.7	26	21
	2	8.4	4	43.3	3.7	10.7	8	7.5	35	23
	3	2.9	5	32.3	3.5	7	6	7.1	35	32
	4	1.7	5	35.4	2.7	9.7	3	6.2	40	30
	5			A 5 th reach was never delineated						
Salt Creek 1996 (Upper only)	6	0.7	2.7	26	2.8	32.5	36.6	10.9	47	41
	7	1.1	3.2	23.8	2.8	----	23.7	13.9	50	45
	8	0.56	0.8	18.9	2.9	11.4	71.3	21.3	56	46
	9	1.14	0.6	17.6	2.1	42.9	44.1	11.4	61	50
	10	1.76	4.8	21.1	2	----	19.9	5.3	53	50

** Calculated from pools per yardage data (size of pool not noted, only depths ≥ 3-4 ft & 4-5 ft)

Observed Pool Values and Objectives for Streams Located in Subwatershed 20 1

Stream Name	Reach No.	Reach Length (miles)	Gradient (%)	Average Width (ft)	Residual Depth (ft)	Width/Depth Ratio	% Area in Pools	Observed Pools/Mile	Minimum Objective PACFISH	Minimum Objective Forest Plan
Tumble 1964	Mouth	----	>6	3	----	----	----	----	184	352
Sage 1964	Mouth	----	>6	3	----	----	----	----	184	352
Pepper 1964	Mouth	----	>6	2	----	----	----	----	184	528
Sugar 1964	Mouth	----	>6	2	----	----	----	----	184	528
Gobel 1964	Station 1	0.1	>6	6	----	----	----	----	164	181
Basin 1964	Mouth	----	>6	5	----	----	----	----	184	211
Fin Roberts 1964	Mouth	----	>6	DRY	----	----	----	----	----	----
McCredie 1964	Mouth	----	>6	3	----	----	----	----	184	352
McCredie 1992	1	0.25	6	6.3	1.9	----	1.7	----	138	141
	2	0.11	7	5.2	1.6	----	6.6	----	122	125
	3	0.1	7	8.1	----	3.2	0	----	130	132
	4	0.12	7	8.4	----	----	0	----	122	125
	5	0.15	6	10.8	----	----	0	----	90	98
	6	0.14	10	5.2	----	6	0	----	184	240
	7	0.15	11.3	3.8	----	2	0	----	184	280
	8	0.25	30	5.9	----	10.9	0	----	184	190
T20e 36.7 1992	1	0.4	2	7	----	3.2	0	0	145	151
	2	----	----	----	----	----	----	----	----	----
	3	----	----	----	----	----	----	----	----	----
	4	----	----	----	----	----	----	----	----	----
	5	----	----	----	----	----	----	----	----	----
T20e38.6 1992	1	----	----	----	----	----	----	----	----	----

SALT CREEK WATERSHED ANALYSIS

Aquatic Habitat and Species Lists CONTINUED

Observed Pool Values and Objectives for Streams Located in Subwatershed 20 1

Stream Name	Reach No.	Reach Length (miles)	Gradient (%)	Average Width (ft)	Residual Depth (ft)	Width/Depth Ratio	% Area in Pools	Observed Pools/ Mile	Minimum Objective PACFISH	Minimum Objective Forest Plan
T20f 41.3 1992	1	0.16	1	2	—	—	—	—	184	>350
	2	—	14	3.5	—	—	—	—	184	308
	3	—	8	3	—	—	—	—	184	352
	4	—	6	3.5	—	—	—	—	184	308
T20f 41.5 1992	1	0.1	4	—	—	—	—	—	—	—
	2	0.3	9	—	—	—	—	—	—	—
	3	0.4	16	—	—	—	—	—	—	—
T20f 43.3 1992	1	0.2	5	12	—	—	—	91	94	
Warner 1964	Station 1	mouth	>6	2	—	—	—	—	184	352
Warner 1996	1	0.5	14	8.1	1.1	22.4	20	3.9	130	132
	2	0.5	14	7.6	1.2	21.1	26.4	0	138	141
	3	0.9	21	5.8	0.8	11.36	—	168.96*	184	190

* 1996 Reach 3 modified Level II survey resulted in elevated pools/mile because every pool is considered

Observed Pool Values and Objectives for South Fork Salt Creek, Subwatershed 20 2 (1964, 1996 data)

Stream Name	Reach No.	Reach Length (miles)	Gradient (%)	Average Width (ft)	Residual Depth (ft)	Width/Depth Ratio	% Area in Pools	Observed Pools/ Mile	Minimum Objective PACFISH	Minimum Objective Forest Plan
S.Fk. Salt Creek 1964	1	0 - 0.9	>6	8	—	—	—	—	130	132
	2	0.9 - 2.9	<4	9	—	—	—	21.4*	115	119
	3	2.9 - 3.9	>6	6	—	—	—	107.3*	138	141
S.Fk. Salt Creek 1996	1	0.8	9	17.4	2.3	35.6	25.9	11.5	61	61
	2	1	1	private land—denied access						
	3	1.1	3	9.5	1.8	23.9	66.7	5.4	106	112
	4	0.1	42	Poor access—not surveyed						
	5	1	13	11.2	1.5	20	—	114.97**	90	96
	6	0.6	12	10.7	1.1	19	—	11.81**	90	98
	7	1.1	3	6.3	1.3	25	—	90.2**	153	170
	8	1	11	7.5	0.9	19	—	184.8**	145	151
	9	0.6	8	4.3	0.5	18	—	188.1**	184	230

* Calculated from pools per yardage data

** 1996 modified Level II survey resulted in elevated pools/mile because every pool is considered

SALT CREEK WATERSHED ANALYSIS

Aquatic Habitat and Species Lists CONTINUED

Observed Pool Values and Objectives for Streams Located in Subwatershed 20 2

Stream Name	Reach No.	Reach Length (miles)	Gradient (%)	Average Width (feet)	Residual Depth (feet)	Width /Depth Ratio	% Area in Pools	Observed Pools/mile	Min.Obj. PACFISH	M. For. Plan
T20f 45.3 1992	1	0.1	7	3.6	1.8	---	4.7	21	184	310
T20f 46.3 1992	1	0.36	6	6.3	---	---	0	0	153	170
	2	0.24	5	4.6	---	---	0	0	184	240
	3	---	---	---	---	---	---	---	---	---
	4	---	15	---	---	---	---	---	---	---
T20w 11.8 1992	1	0.6	3	---	---	---	0	0	---	---
	2	0.7	4	---	---	---	---	---	---	---
Eagle 1937 Eagle 1964	Station A-B	0.9	---	---	---	---	---	---	---	---
	Station 1	Mouth	>6	25	---	---	---	---	47	40
	Station 2	at 2 miles	>6	15	---	---	---	---	70	81
	Station 3	at 3.5 miles	>6	10	---	---	---	---	96	106
Eagle 1992	1	0.8	2	26.9	3.15	5	50.8	16.3	45	41
	2	0.3	5	22.5	3.2	6.73	16.9	16.7	50	48
	3	1	12	25.8	2.7	8.89	9	24	47	40
	4	0.3	3	19.2	2	---	32.9	3.3	56	56
	5	0.3	4	20.7	2	5.33	0	3.3	56	50
	6	0.4	3	14.7	2.7	---	2.1	5	70	72
	7	0.7	5	12.4	2.6	5.63	6.8	0	91	94
	8	1	3	7.3	4.3	5.17	0.9	2	145	151
	9	0.3	9	6.3	1.3	---	89.9	3.3	164	181
	10	0.2	12	3.9	---	---	---	---	184	---
Coyote 1964	Station 1	mouth	>6	3	---	---	---	---	184	---
Coyote 1992	1	0.6	3	1.8	0.4	---	2.3	3.3	184	528
	2	0.8	20	3	---	---	---	---	184	352
Fall 1996	1	0.9	16	17.6	2.3	28.3	33.1	10.6	61	58
	2	0.6	6	12.5	1.6	13.3	20.6	4.7	91	94
	3	0.5	16	10.5	1.9	9.4	16.6	11.6	93	102
	4	1.1	10	7.2	---	---	Poor access—not surveyed		145	151
	5	0.9	12	11.1	1.3	5.8	---	89.76*	90	96
	6	0.3	2	12.8	3.6	7.5	---	32.27*	91	94
Diamond 1996	1	0.8	4	10.6	1.4	16.4	29.7	58	122	125
	2	0.9	7	9.9	1.9	13.7	17.4	45	102	110
	3	0.8	7	9.2	1.3	12.2	21.4	2.5	106	112

* 1996 modified Level II survey resulted in elevated pools/mile because every pool is considered

SALT CREEK WATERSHED ANALYSIS

Aquatic Habitat and Species Lists CONTINUED

Observed Pool Values and Objectives for Deer Creek, Subwatershed 20 3 (1996 data)

Stream Name	Reach No.	Reach Length (miles)	Gradient (%)	Average Width (feet)	Residual Depth (feet)	Width /Depth Ratio	% Area in Pools	Observed Pools/mile	Min.Obj. PACFISH	Min. Obj. Forest Plan
Deer 1996	1	0.7	6	12.2	1.5	11.3	32.4	2.8	91	94
	2	1.2	10	9.4	1.8	---	8.4	---	106	112
	3	0.2	2	10.3	1.1	5.11	---	112.2*	93	102

* 1996 modified Level II survey resulted in elevated pools/mile because every pool is considered

Instream Large Woody Material (LWM) for Mainstem Salt Creek (1990, 1996 Stream Survey Data) *

Stream Name	Reach No.	Reach Length (miles)	No. Small LWM/mile	No. Medium LWM/mile	No. Large LWM/mile	No. Med+Large LWM per mile	Minimum LWD/mile PACFISH	Minimum LWD/mile Forest Plan
Salt Creek 1990 (lower)	1**	6.9	14	8	4	12	80	105
	2**	8.4	17	16	15	31	80	105
	3	2.9	30	14	3	17	80	105
	4	1.7	14	14	1	15	80	105
	5	Not Surveyed						
Salt Creek 1996 (Upper)	6	0.7	36.6	16.6	5	21.6	80	105
	7	1.1	27.3	0.9	2.7	3.6	80	105
	8	0.56	110	38.3	20	58.3	80	105
	9	1.14	120.7	17.7	0.77	18.5	80	105
	10	1.76	77.6	12.9	7.6	20.5	80	105

* 1937 and 1964 surveys lack detailed LWM data

** Additional large and medium woody material from habitat structures

Instream Large Woody Material (LWM) for Streams in Subwatershed 20 1 (1992 Stream Survey Data)

Stream Name	Reach No.	Reach Length (miles)	No. Small LWM/mile	No. Medium LWM/mile	No. Large LWM/mile	No. Med+Large LWM per mile	Minimum LWD/mile PACFISH	Minimum LWD/mile Forest Plan
McCredie 1992	1	0.25	55.5	27.7	19.8	47.5	80	105
	2	0.11	18.9	37.9	75.8	113.7	80	105
	3	0.1	38.7	-----	-----	-----	80	105
	4	0.12	16.2	24.3	8.1	32.4	80	105
	5	0.15	62.7	48.7	13.9	62.6	80	105
	6	0.14	24	48.1	8	56.1	80	105
	7	0.15	78.4	47.1	15.7	62.8	80	105
	8	0.25	91.7	116.7	66.7	183.4	80	105
T20 e 36.7 1992	1	0.4	-----	-----	-----	-----	80	105
	2	-----	-----	-----	-----	-----	80	105
	3	-----	-----	-----	-----	-----	80	105
	4	-----	-----	-----	-----	-----	80	105
	5	-----	-----	-----	-----	-----	80	105
T20 f 41.3	1	0.16	76	151.9	227.9	-----	80	105
	2	-----	-----	-----	-----	-----	80	105
	3	-----	-----	-----	-----	-----	80	105
	4	-----	-----	-----	-----	-----	80	105

NOTE: Small LWM is 12 inch dbh and 25 feet in length Medium LWM is 24 in. dbh and 50 feet in length Large LWM is 36 inch dbh and 50 feet in length

SALT CREEK WATERSHED ANALYSIS

Aquatic Habitat and Species Lists CONTINUED

Instream Large Woody Material (LWM) for Warner Creek, Subwatershed 20 1 (1996 Stream Survey Data)

Stream Name	Reach No.	Reach Length (miles)	No. Small LWM/mile	No. Medium LWM/mile	No. Large LWM/mile	No. Med+Large LWM per mile	Minimum LWD/mile PACFISH	Min. LWD/mile Forest Plan
Warner 1996	1	0.5	128	60	10	70	80	105
	2	0.5	120	66	28	94	80	105
	3	0.9	107	61	13	74	80	105

Instream Large Woody Material (LWM) for S.Fk.Salt Creek, Subwatershed 20 2 (1996 Stream Survey Data)

Stream Name	Reach No.	Reach Length (miles)	No. Small LWM/mile	No. Medium LWM/mile	No. Large LWM/mile	No. Med+Large LWM per mile	Minimum LWD/mile PACFISH	Minimum LWD/mile Forest Plan	
SFk Salt Ck 1996	1	0.8	63.75	26.25	13.75	40	80	105	
	2	1	Denied access - not surveyed					80	105
	3	1.1	48.2	16.4	6.4	22.8	80	105	
	4	0.1	Not surveyed					80	105
	5	1	53	14	3	17	80	105	
	6	0.6	17	3	0	3	80	105	
	7	1.1	35	23	3	26	80	105	
	8	1	20	16	1	17	80	105	
	9	0.6	48	12	3	15	80	105	

Instream Large Woody Material (LWM) for Streams in Subwatershed 20 2 (1992 Stream Survey Data)

Stream Name	Reach No.	Reach Length (miles)	No. Small LWM/mile	No. Medium LWM/mile	No. Large LWM/mile	No. Med+Large LWM per mile	Minimum LWD/mile PACFISH	Minimum LWD/mile Forest Plan
T20 f 45.3 1992	1	0.1	175.3	133.2	49.1	182.3	80	1
T20 f 46.3 1992	1	0.36	87.9	76.9	41.2	118.1	80	105
	2	0.24	72.8	44.5	12.1	56.6	80	105
	3	----	----	----	----	----	80	105
	4	----	----	----	----	----	80	105
T20 w 11.8 1992	1	0.6	9.4	4	0	4	80	105
	2	0.7	----	----	----	----	80	105
Eagle 1992	1	0.8	971.8*	722.3*	302*	1024.3*	80	105
	2	0.3	312.5*	312.5*	0	312.5*	80	105
	3	1	46*	43.2*	21.2*	64.4*	80	105
	4	0.3	0*	0	0	0	80	105
	5	0.3	271.3*	162.8*	54.3*	217.1*	80	105
	6	0.4	43.7*	21.8*	21.8*	43.6*	80	105
	7	0.7	82.1*	83.7*	40.2*	123.9*	80	105
	8	1	36.1*	22.6*	12*	34.6*	80	105
	9	0.3	9139*	5415.4*	3046.2*	8461.6*	80	105
	10	0.2	21120*	31680*	31680*	63360*	80	105
Coyote 1992	1	0.6	51.7	6	2	8	80	105
	2	0.8	----	----	----	----	80	105

* 1992 Eagle Creek LWM data are suspect

SALT CREEK WATERSHED ANALYSIS

Aquatic Habitat and Species Lists CONTINUED

Instream Large Woody Material (LWM) for Streams in Subwatershed 20 2 (1996 Stream Survey Data)

Stream Name	Reach No.	Reach Length (miles)	No. Small LWM/mile	No. Medium LWM/mile	No. Large LWM/mile	No. Med+Large LWM per mile	Minimum LWD/mile PACFISH	Minimum LWD/mile Forest Plan	
Fall 1996	1	0.9	27.8	27.8	2.2	28.9	80	105	
	2	0.6	21.7	5	0	5	80	105	
	3	0.5	18	8	0	8	80	105	
	4	1.1	No Access - not surveyed					80	105
	5	0.9	69	4	0	4	80	105	
	6	0.3	76	0	0	0	80	105	
Diamond 1996	1	0.8	35	13.8	0	13.8	80	105	
	2	0.9	30	7.8	4.4	12.2	80	105	
	3	0.8	26.3	5	1.25	6.25	80	105	

Instream Large Woody Material (LWM) for Deer Creek, Subwatershed 20 3 (1996 Stream Survey Data)

Stream Name	Reach No.	Reach Length (miles)	No. Small LWM/mile	No. Medium LWM/mile	No. Large LWM/mile	No. Med+Large LWM per mile	Minimum LWD/mile PACFISH	Minimum LWD/mile Forest Plan
Deer 1996	1	0.7	22.8	15.7	0	15.7	80	105
	2	1.2	33	13	1	21	80	105
	3	0.2	40	0	0	0	80	105

Physical Information for Mainstem Salt Creek (1990, 1996 Stream Survey Data)

Stream	Reach No.	Reach Length (miles)	Valley Segment Type	Dominant Substrate	Subdominant Substrate	Channel Stability Rating	# Failures (Average Size)	# Wood Debris Jams	Rosgen Channel Type		
Salt Creek 1990 (lower)	1	6.9	M2	CO	SB	--	--	--	B		
	2	8.4	M2	CO	SB	--	--	--	B		
	3	2.9	M2	CO	SB	--	--	--	B		
	4	1.7	M2	SB	CO	--	--	--	G		
	5	No Survey									
				Wetted Channel Substrates							
				% SA	% GR	% CO	% BO	% BR			
Salt Creek 1996 (upper)	6	0.7	V1	9	31	28	11	22	--	--	B
	7	1.1	M1	14	21	20	19	25			B
	8	0.6	F6	44	37	17	1	0			DA
	9	1.1	F3	31	38	24	6	2			C
	10	1.8	V2	7	27	31	21	13			A

SALT CREEK WATERSHED ANALYSIS

Aquatic Habitat and Species Lists CONTINUED

Physical Information for Streams Located in Subwatershed 20 1

Stream	Reach No.	Reach Length (miles)	Valley Segment Type	Dominant Substrate	Subdominant Substrate	# Failures (Average Size)	# Failures (Average Size)	# Wood Debris Jams	Ro. Channel Type			
Tumble, 1964	Sta 1	Mouth	--	BO	GR	--	--	--				
Sage, 1964	Sta 1	Mouth	--	BO	GR	--	--	--				
Pepper, 1964	Sta 1	Mouth	--	BO	GR	--	--	--				
Sugar, 1964	Sta 1	Mouth	--	BO	GR	--	--	--				
Gobel, 1964	Sta 1	0.1	--	BO	GR	--	--	--				
Basin, 1964	Sta 1	Mouth	--	BO	CO	--	--	--				
Fin Roberts 1964	Sta 1	Mouth	--	BO	CO	--	--	--				
McCredie, 1964	Sta 1	Mouth	--	BO	CO	--	--	--				
McCredie 1992	1	0.25	F4	CO	SB	FAIR	Some MW	--				
	2	0.11	U2	CO	SB	FAIR	Some MW	--				
	3	0.1	U2	SB	CO	FAIR	Some MW	--				
	4	0.12	U2	SB/CO	CO	FAIR	Some MW	--				
	5	0.15	U2	GR	CO	FAIR	small failures	1				
	6	0.14	V4	CO	GR	FAIR	--	--				
	7	0.15	H1	CO	GR	FAIR	--	--				
	8	0.25	H2	CO	GR/CO	N/A	--	--				
T20e 36.7	1	0.4	val code 1	SA	GR	--	--	--				
	2	--	--	--	--	--	--	--				
	3	--	--	--	--	--	--	--				
	4	--	--	--	--	--	--	--				
	5	--	--	--	--	--	--	--				
T20f 41.3	1	0.16	val code 1	SA	GR	FAIR	--	0				
	2	--	V4	--	--	FAIR	--	2				
	3	--	V4	--	--	FAIR	--	5				
	4	--	H2	--	--	FAIR	0	0				
T20f 41.5 1992	1	0.1	val code 1	SA	GR	--	--	--				
	2	0.3	M2	GR	SA	--	--	--				
	3	0.4	H2	SA	GR	--	--	--				
T20f 43.3 1992	1	0.2	M2	CO	SB	--	--	10				
Warner, 1964	Sta 1	Mouth	--	BO	GR							
				Wetted Channel Substrates								
				% SA	% GR	% CO	% BO	% BR				
Warner, 1996	1	0.5	V2	16	26	31	25	0.8	--	14 (22X4)	6	
	2	0.5	V2	16	31	31	22	0	--	2 (6X1)	3	
	3	0.9	H2			D	SD		--	18 (3.8% of Reach)	20+	Aa+

D = dominant, SD = subdominant

SALT CREEK WATERSHED ANALYSIS

Aquatic Habitat and Species Lists CONTINUED

Physical Information for South Fork Salt Creek, Subwatershed 20 2 (1964 and 1996 Data)

Stream	Reach No.	Reach Length (miles)	Valley Segment Type	Wetted Channel Substrates					Channel Stability Rating	# Failures (Average Size)	# Wood Debris Jams	Rosgen Channel Type
				% SA	% GR	% CO	% BO	% BR				
S.Fk. Salt 1964	Sta 1	0.9	--	CO/BR		BR			--	--	--	--
	Sta 2	2.0	--	GR		SA			--	--	4	--
	Sta 3	1.0	--	BR		BO			--	--	1	--
				Wetted Channel Substrates								
				% SA	% GR	% CO	% BO	% BR				
S. Fk. Salt	1	0.8	U2	21	18	36	25	0.2	FAIR	10 (50x10)	5	B
	2	1.0	U1	23	43	28	6	0.5	--	--	--	C
	3	1.1	U1	--	--	--	--	--	FAIR	6 (27x4)	15	B
	4	0.1	V3							No Survey		Aa+
	5	1.0	V2			SD		D	FAIR	6 (23X11)	25	Aa+
	6	0.6	U2		D		SD		FAIR	5 (18X5)	25	B
	7	1.1	U1		D	SD			POOR	14 (20X4)	20	C
	8	1.0	V2		D	SD			FAIR	17 (19X5)	15	A
	9	0.6	F6		D		SD		GOOD	2 (7X4)	25	B

D = dominant, SD = subdominant

Physical Information for Streams Located in Subwatershed 20 2

Stream Name	Reach No.	Reach Length (miles)	Valley Segment Type	Dominant Substrate	Subdominant Substrate	Channel Stability Rating	# Failures (Average Size)	# Wood Debris Jams	Rosgen Channel Type
T20f 45.3 1992	1	0.1	M2	SA	CO	--	--	1	
T20f 46.3 1992	1	0.36	H1	SA	GR/SB	--	--	--	
	2	0.24	U1	SA	GR	--	--	--	
	3	--	H2	SA	SB	--	--	--	
	4	--	H2	SA	SA	--	--	--	
T20w 11.8 1992	1	0.6	H1	GR	CO	--	--	--	
	2	0.7	H1	GR	SA	--	--	--	
Eagle 1937	Sta A-	0.9	--	large CO	small CO	--	--	--	
Eagle 1964	Sta 1	Mouth	--	CO/BO	GR	--	--	--	
	Sta 2	@ 2 mi	--	CO/BO	GR	--	--	--	
	Sta 3	@ 3.5 mi	--	CO/BO	GR	--	--	--	
Eagle 1992	1	0.8	F2	CO	GR	--	--	--	
	2	0.3	V4	SB	CO	--	--	1	
	3	1.0	V2	SB	GR	--	1 (50x100)	2	
	4	0.3	V4	GR	SB	--	--	--	
	5	0.3	M2	CO	GR	--	--	--	
	6	0.4	V4	CO	GR	--	--	3	
	7	0.7	V4	CO	GR	--	--	blowdown	
	8	1.0	V4	GR	SB	--	4 (20x30)	3	
	9	0.3	M2	SA	GR	--	--	1	
	10	0.2	H2	GR	SA	--	--	blowdown	
Coyote 1964	Sta 1	Mouth	--	BO/CO	GR	--	--	--	
Coyote 1992	1	0.6	F6	SA	GR	--	--	--	
	2	0.8	H1	CO	SB	--	1 (10x4)	--	

SALT CREEK WATERSHED ANALYSIS

Aquatic Habitat and Species Lists CONTINUED

Physical Information for Streams Located in Subwatershed 20 2

Stream Name	Reach No.	Reach Length (miles)	Valley Segment Type	Wetted Channel Substrates					Channel Stability Rating	# Failures (Average Size)	# Wood Debris Jams	Rosgen Channel Type	
				% SA	% GR	% CO	% BO	% BR					
Fall 1996	1	0.9	V2	3	35	41	12	0.9	--	3 (28 ft)	5	Aa+	
	2	0.6	V1	19	30	43	13	13	--	1 (15 ft)	1	A	
	3	0.5	U2	19	17	25	25	32	--	1 (15 ft)	--	Aa+	
	4	1.1	U1							No Survey		A	
	5	0.9	V2					D	SD	--	4 (0.7% of Reach)	1	Aa+
	6	0.3	F5	D	SD					--	--	--	B
Diamond 1996	1	0.8	F5	10	35	28	15	12	--	(105 ft)	--	B	
	2	0.9	V1	11	27	22	10	30	--	(195 ft)	--	A	
	3	0.8	U2	8	24	24	22	22	--	(165 ft)	--	Aa+	

D = dominant, SD = subdominant

Physical Information for Deer Creek, Subwatershed 20 3

Stream Name	Reach No.	Reach Length (miles)	Valley Segment Type	Wetted Channel Substrates					Channel Stability Rating	# Failures (Average Size)	# Wood Debris Jams	Rosgen Channel Type
				% SA	% GR	% CO	% BO	% BR				
Deer 1996	1	0.7	M1	6	26	32	11	25	FAIR	7 (15x10)	3	A
	2	1.2	V2	3	18	23	12	44	FAIR	5 (20x5)	3	Aa+
	3	0.2	H1		SD	D			FAIR	3 (6x1)	0	F

D = dominant, SD = subdominant

Biological Information About Species Located in the Riparian Area of Mainstem Salt Creek (1937, 1964, 1980, 1987 Data)

Stream & Location	Survey Date	Reach No.	Seral Stage Inner	% Inner Riparian	Inner Riparian Species	Seral Stage Outer	% Outer Riparian	Outer Riparian Species	Water Temp (F) (Time)	Fish Species	Amphib. Species
Salt Creek 20 1,20 2	8/28/37	Sta A-B	---	---	HA, HX, maple	---	---	CX	51 (1000)	CH, ONXX	---
		Sta B-C	---	---	HA, HX, maple	---	---	CX	50 (1100)	CH, ONXX	---
		Sta C-D	---	---	HA, HX, maple	---	---	CX	50 (1230)	CH, ONXX	---
		Sta D-E	SS	---	HA, HX, maple	---	---	CX	50 (1445)	CH, ONXX	---
		Sta E-F	SS	---	HA, HX, maple	---	---	CX	52 (1630)	CH, ONXX	---
Salt Creek 20 1- 20 3	8/64	1	---	---	HA, HX, maple	---	---	CX	56 (1115)	RB, CT	---
		2	---	---	HA, HX, maple	---	---	CX	58 (1440)	RB, CT	---
		3	---	---	HA, HX, maple	---	---	CX	52 (1100)	RB, CT	---
		4	---	---	HV, CX	---	---	CX	52 (1110)	CT	---
		5	---	---	HX, CX	---	---	CX	51 (1000)	CT	---
		6	---	---	HA	---	---	CX	50 (1000)	CT	---
Salt Creek 20 3	8/20/80	6	---	---	HA, HV, CD	---	---	---	50 (1400)	---	---
	8/20/80	7	---	---	---	---	---	---	63 (1600)	RB	---
	8/21/80	?	---	---	---	---	---	---	59 (1100)	RB, BT	---
	8/21/80	10	---	---	GF, CD	---	---	---	61 (1345)	RB, BT	---
	8/21/80	10	---	---	GF, HA	---	---	---	50 (1600)	RB, BT	---
	9/17/80	11	---	---	GF, HW	---	---	---	46 (1200)	BT	---
Salt Creek 20 2	8/3/87	wicopee	---	---	HX, CX	---	---	---	62 (1100)	RB,CT,SC, SD	---
	8/3/87	wicopee	---	---	HX,CX	---	---	---	62 (1400)	RB,CT,SC	---

SALT CREEK WATERSHED ANALYSIS

Aquatic Habitat and Species Lists CONTINUED

Biological Information About Species Located in the Riparian Area of Mainstem Salt Creek
(1990, 1996 Data)

Stream & Location	Survey Date	Reach No.	Seral Stage Inner	% Inner Riparian	Inner Riparian Species	Seral Stage Outer	% Outer Riparian	Outer Riparian Species	Water Temp (F) (Time)	Fish Species	Amphib. Species
Salt Creek 20 1, 20 2 (Lower)	7/9/90	1	LT	100	CC, HA	LT	100	CD, CC	58	CT, RB, BT, SC, MW, LM	
		2	LT	48	CD, HA	LT	62	CD, CC	53	BT	---
		3	SP	58	HA, CD	LT	58	CD, CC	51	---	---
		4	SP	90	HA, HV	LT	50	CD, CC	55	---	---
	8/10/90	2	---	---	---	---	---	---	50	CT, RB, M W, SC	---
	8/10/90	2	---	---	---	---	---	---	---	CT, RB, M W	---
	8/10/90	2	---	---	---	---	---	---	---	CT, RB, M W	---
Salt Creek 20 3	8/9/96	6	SS	100	HA, HW, HX	LT	100	CD, CC	50	---	---
		7	SS	100	HA, HW, HX	LT	100	CD, CC	55 (1330)	CT, RB, BT	---
		8	SS	100	HA, HX	LT	67	CD, CF, CH	56 (1530)	BT	---
		9	SS	100	HA, HW, HX	LT	100	CF, CH, CW	54	---	---
		10	SS	75	HA, HX, CW, CX	LT	100	CD, CH, CF	3	---	Unk frog

Biological Information About Species Located in the Riparian Area of Streams in Subwatershed 20 1

Stream & Location	Survey Date	Reach No.	Seral Stage Inner	% Inner Riparian	Inner Riparian Species	Seral Stage Outer	% Outer Riparian	Outer Riparian Species	Water Temp (F) (Time)	Fish Species	Amphib. Species
Tumble	9/9/64	-----	-----	-----	-----	-----	-----	CX	-----	-----	-----
Sage	9/9/64	-----	-----	-----	-----	-----	-----	CX	49 (1230)	-----	-----
Pepper	9/9/64	-----	-----	-----	-----	-----	-----	CX	49 (1230)	-----	-----
Sugar	9/9/64	-----	-----	-----	-----	-----	-----	CX	49 (1230)	-----	-----
Gobel	9/9/64	-----	-----	-----	-----	-----	-----	CX	49 (1300)	-----	-----
Basin	9/9/64	-----	-----	-----	-----	-----	-----	CX	47 (1300)	-----	-----
Fin Roberts	9/9/64	-----	-----	-----	-----	-----	-----	CX	---	-----	-----
McCredie	9/9/64	-----	-----	-----	-----	-----	-----	CX	49 (1300)	-----	-----
McCredie	4/21/92	1	-----	-----	-----	-----	-----	-----	47 (1100)	-----	-----
		2	-----	-----	-----	-----	-----	-----	-----	-----	-----
		3	SP	100	CC, CH	-----	-----	-----	-----	-----	-----
		4	-----	-----	-----	-----	-----	-----	43 (1400)	-----	-----
		5	-----	-----	-----	-----	-----	-----	47 (1230)	-----	-----
		6	SS	100	CC, HW	-----	-----	-----	-----	-----	-----
		7	SS	100	CH, CC	-----	-----	-----	44 (0915)	-----	-----
		8	SP	100	CH, CD	-----	-----	-----	44 (1000)	-----	-----
T20e 36.7	5/5/92	1	LT	99	CC, HV	-----	-----	-----	50 (0900)	---	unk frog
		DATA NOT AVAILABLE FOR REACHES 2 -5									
T20e 38.6	5/6/92	1	-----	-----	-----	-----	-----	-----	-----	-----	-----
T20f 41.3	5/12/92	1	SS	-----	HW	-----	-----	-----	-----	CT	---
		2	-----	-----	-----	-----	-----	-----	-----	-----	-----
		3	GF	-----	-----	-----	-----	-----	-----	-----	-----
		4	GF	-----	-----	-----	-----	-----	-----	-----	-----

SALT CREEK WATERSHED ANALYSIS

Aquatic Habitat and Species Lists CONTINUED

Biological Information About Species Located in the Riparian Area of Streams in Subwatershed 20 1

Stream & Location	Survey Date	Reach No.	Seral Stage Inner	% Inner Riparian	Inner Riparian Species	Seral Stage Outer	% Outer Riparian	Outer Riparian Species	Water Temp (F) (Time)	Fish Species	Amphib. Species
T20f 41.5	5/13/92			NO DATA FOR REACHES 1-3							
T20f 43.3	5/20/92	1	NV	---	---	---	---	---	---	0	---
Warner	1964	Sta 1	----	----	----	----	----	CX	48 (1300)	----	----
Warner	7/3/96	1	SP	47	HA, HV	ST	73	CC, CF	55	CT	DITE
		2	ST	67	CF, CH	ST	67	CC, CH	56	ONXX	----
		3	LT	67	CC, CF	LT	57	CC, CF	50	----	DITE, ASTR, RAAU

Biological Information About Species Located in the Riparian Area of S. Fk. Salt Creek (1964, 1970, 1995, 1996 Stream Survey Data)

Stream & Location	Survey Date	Reach No.	Seral Stage Inner	% Inner Riparian	Inner Riparian Species	Seral Stage Outer	% Outer Riparian	Outer Riparian Species	Water Temp (F) (Time)	Fish Species	Amphib. Species	
S. Fk. Salt	1964	Sta 1	----	-----	HX, CX	----	-----	---	54 (1300)	CT, SC	-----	
		Sta 2	----	-----	HV, HW, HX	----	-----	---	56 (1653)	CT	---	
		Sta 3	----	-----	CX	----	-----	---	46 (1345)	----	---	
S. Fk. Salt		1	----	-----	----	----	-----	---	57	CT	---	
S. Fk. Salt	8,9/85	1-2	----	-----	----	----	-----	---	49.5	CT, UNK, SC	---	
		5	----	-----	----	----	-----	---	---	---	---	
		7	----	-----	----	----	-----	---	56	---	DIT RAC ASTR	
		9	----	-----	----	----	-----	---	---	---	RACA, ASTR	
S. Fk. Salt	8/15/96	1	SS,LT	86	HA, HW, CC, CF	LT	57	CF, CC	55	---	---	
		2						DENIED ACCESS - NOT SURVEYED				
		3	SS	91	HA, HV, HW, HD	SP	73	CF, CF	68	CT	ASTR	
		4						NOT SURVEYED				
		5	ST	57	CC, CF	ST	57	CC, CF	47	0	ASTR, RACA DITE, TAGR	
		6	SP	94	HA, HW	ST	57	CF, CF	---	---	ASTR	
		7	SS,LT	76	HA, HV, CF, CH	LT	67	CF, CH	43	---	RACA, ASTR	
		8	SS	80	HW, HW	LT	52	CF, CH	48.5	---	RACA, ASTR	
		9	ST	80	CF, CH	ST	80	CF, CH	45	---	RACA, ASTR	

SALT CREEK WATERSHED ANALYSIS

Aquatic Habitat and Species Lists CONTINUED

Biological Information About Species Located in the Riparian Area of Streams in Subwatershed 20 2

Stream & Location	Survey Date	Reach No.	Seral Stage Inner	% Inner Riparian	Inner Riparian Species	Seral Stage Outer	% Outer Riparian	Outer Riparian Species	Water Temp (F) (Time)	Fish Species	Amphib. Species	
T20f 45.3	5/20/92	1	----	----	----	----	----	----	----	----	----	
T20f 46.3	1992	1	----	----	----	2nd grow	----	----	44 (1000)	----	RAAU	
		2	----	----	----	2nd grow	----	----	----	----	----	
		3	----	----	----	LB3, RB3	----	----	----	----	----	
		4	----	----	----	LB3, RB3	----	----	----	----	----	
T20w 11.8	8/25/92	1	ST	100	CD, HA	----	----	----	47 (1100)	----	----	
		2	----	----	----	----	----	----	----	----	----	
Eagle	1937	Sta A-B	----	----	----	----	----	----	----	----	----	
Eagle	1964	Sta 1	----	----	----	----	----	CX	42 (1430)	RB, SC	----	
		Sta 2	----	----	----	----	----	CX	42 (1500)	0	----	
		Sta 3	----	----	----	----	----	----	----	----	----	
Eagle	5/28/92	1	SP	100	HA, CC	----	----	----	48	----	----	
		2	ST	100	CH, CC	----	----	----	47 (1200)	----	----	
		3	LT	100	CC, CS	----	----	----	44 (1140)	----	----	
		4	----	----	----	LB4, RB2	----	----	----	----	----	
		5	MT	100	CD, CC	----	----	----	43 (1100)	----	----	
		6	----	----	----	LB6, RB6	----	----	47 (1150)	----	----	
		7	ST	100	HV, CH	----	----	----	----	----	----	
		8	LT	100	CD, HW	----	----	----	52 (1520)	----	----	
		9	----	----	----	LB1, RB1	----	----	----	----	----	RAAU
		10	----	----	----	LB1, RB6	----	----	----	----	----	----
Coyote	1964	----	----	----	----	----	----	CX	52 (1330)	----	----	
Coyote	8/4/92	1	ST	100	CD, HA	----	----	----	55 (1000)	unk. species	----	
		2	----	----	----	Reprod	----	----	----	----	----	
Fall	1995	1	----	----	----	----	----	----	44 (1300)	CT, unk	ASTR	
		2	----	----	----	----	----	----	----	BT	----	
		3	----	----	----	----	----	----	----	----	----	
		4	----	----	----	----	----	----	----	----	----	
		5	----	----	----	----	----	----	----	----	----	
Fall	1996	1	SS	83	HA, HB, CF	LT	100	CF, CH	46	----	ASTR, RACA	
		2	SS	50	HA, CF	LT	67	CF, CH	48	BT	----	
		3	SS	84	HA, CH	LT	100	CF, CH	43	----	----	
		4	----	----	----	----	NO ACCESS - NOT SURVEYED					
		5	SS	100	HA, CH	LT	100	CF,CM	39	0	----	
		6	----	----	----	LT	100	CF,CM	40	0	----	
Diamond	1995	1	----	----	----	----	----	----	46 (1130)	CT, BT	----	
Diamond	1996	1	SS	100	HA,HW, HV,CH	LT	94	CH, CF	64	UNK	RACA	
		2	SS	90	HA,CH,CF	LT	100	CH, CF	60	----	----	
		3	SS	100	HA,CH,CF	LT	100	CF, CH	55	----	----	

SALT CREEK WATERSHED ANALYSIS

Aquatic Habitat and Species Lists CONTINUED

Biological Information About Species Located in the Riparian Area of Deer Creek , Subwatershed 20 3
(1995, 1996 Stream Survey Data)

Stream & Location	Survey Date	Reach No.	Seral Stage Inner	% Inner Riparian	Inner Riparian Species	Seral Stage Outer	% Outer Riparian	Outer Riparian Species	Water Temp (F) (Time)	Fish Species	Amphib. Species
Deer	9/95	1	-----	-----	-----	-----	-----	-----	42	0	
		2	-----	-----	-----	-----	-----	-----	42	BT	----
		3	-----	-----	-----	-----	-----	-----	-----	0	----
Deer	9/96	1	SS	100	HA, HX	LT	73	CH, CF, CS	48	0	RACA, ASTR
		2	SS	100	HX, HX	LT	91	CF, CH	42	BT	RACA, ASTR
		3	-----	-----	----	SP	100	CF, CH	39	-----	-----

- BT = Brook trout (*Salvelinus fontinalis*)
- CH = Chinook salmon (*Oncorhynchus tshawytscha*)
- CT = Cutthroat trout (*Oncorhynchus clarki*)
- LM = Lamprey (*Lampetra spp.*)
- MW = Mountain whitefish (*Prosopium williamsoni*)
- ONXX = Unknown salmonid species
- RB = Rainbow trout (*Oncorhynchus mykiss*)
- SC = Sculpin species (*Cottus spp.*)
- SD = Speckled dace (*Rhinichthys osculus*)
- UNK = Unknown fish species

Wolman Pebble Counts (1996)

Salt Creek Wolman Pebble Count (20 1)

Reach Number	D50 (mm)	D84 (mm)	Dominant Substrate	Subdominant Substrate
6	70	148		
7	117	567		
8	6	35		
9	34	86		
10	131	355		

Warner Creek Wolman Pebble Count (20 1)

Reach Number	D50 (mm)	D84 (mm)
1	57	201
2	45	159
3	--	--

Fall Creek Wolman Pebble Count (20 2)

Reach Number	D50 (mm)	D84 (mm)
1	35	119
2	238	3399
3	117	3370
4	--	--
5	Modified	Survey
6	No	Wolman

Deer Creek Wolman Pebble Count (20 3)

Reach Number	D50 (mm)	D84 (mm)
1	94	561
2	265	1834.25

Diamond Creek Wolman Pebble Count (20 2)

Reach Number	D50 (mm)	D84 (mm)
1	45	254
2	64	334
3	Modified Survey	No Wolman

APPENDIX I: High lakes Fish Stocking Records

Betty Lake

Year	Species	Number
1994	RB	7242
1995	RB	1920
1996	RB	4000

Lower Betty Lake

Year	Species	Number
1994	RB	578
1996	RB	4462

Birthday Lake

Year	Species	Number
1993	BT	411
1995	BT	480

Boo Boo Lake

Year	Species	Number
1993	CT	303
1995	RB	300
1995	CT	600

LeMay Lake

Year	Species	Number
1993	RB	788
1995	CT	469
1995	RB	1260
1996	RB	413

Lorin Lake

Year	Species	Number
1993	RB	788
1995	CT	1125
1995	RB	800

Lucas Lake

Year	Species	Number
1993	CT	505
1993	RB	506
1996	RB	319
1996	CT	500

Horsefly Lake

Year	Species	Number
1993	BT	294
1995	BT	360
1996	RB	207
1996	BT	300

Howkum Lake

Year	Species	Number
1995	CT	563
1996	RB	207
1996	CT	400

Lower Island Lake

Year	Species	Number
1993	CT	606
1995	CT	750

Upper Island Lake

Year	Species	Number
1993	RB	788
1995	RB	720

JoAnn Lake

Year	Species	Number
1993	CT	808
1995	CT	1313

Verde Lake

Year	Species	Number
1993	CT	303
1995	CT	375

Vivian Lake


Year	Species	Number
1993	BT	1528
1995	BT	1680

BT = Brook trout (*Salvelinus fontinalis*)CT = Cutthroat trout (*Oncorhynchus clarki*)RB = Rainbow trout (*Oncorhynchus mykiss*)

APPENDIX J: Aggregate Recovery Percentages by Drainage

Drainage	Code	Weighted Recovery Rate	Drainage	Code	Weight Recovery Rate
20	1.01	44.64	20	1.36	68.53
20	1.02	86.35	20	2.01	80.91
20	1.03	89.40	20	2.02	76.87
20	1.04	74.25	20	2.03	85.71
20	1.05	91.77	20	2.04	50.28
20	1.06	85.38	20	2.05	83.04
20	1.07	81.23	20	2.06	70.42
20	1.08	82.24	20	2.07	74.38
20	1.09	84.52	20	2.08	76.05
20	1.10	81.48	20	2.09	88.85
20	1.11	84.09	20	2.10	94.45
20	1.12	79.32	20	2.11	67.44
20	1.13	84.77	20	2.12	73.45
20	1.14	78.12	20	2.13	49.19
20	1.15	80.61	20	2.14	68.03
20	1.16	79.82	20	2.15	78.56
20	1.17	74.57	20	2.16	78.03
20	1.18	75.12	20	2.17	73.01
20	1.19	72.46	20	2.18	76.85
20	1.20	73.83	20	2.19	75.74
20	1.21	87.36	20	2.20	55.46
20	1.22	71.28	20	2.21	80.62
20	1.23	87.10	20	2.22	71.14
20	1.24	79.53	20	2.23	86.16
20	1.25	74.70	20	2.24	87.86
20	1.26	75.09	20	2.25	75.82
20	1.27	81.82	20	2.26	82.37
20	1.28	86.29	20	2.27	84.87
20	1.29	80.05	20	3.01	78.80
20	1.30	79.35	20	3.02	83.17
20	1.31	84.00	20	3.03	78.68
20	1.32	76.97	20	3.04	94.70
20	1.33	88.57	20	3.05	85.25
20	1.34	80.10	20	3.06	62.72
20	1.35	65.41	20	3.07	82.13

Salt Creek Watershed Analysis Appendix A Drainages



Appendix K: Literature Cited

- Agee, J. K., 1981. "Fire effects on Pacific Northwest forests: flora, fuels, and fauna." From: Conference Proceedings, Northwest Forest Fire Council 1981, Portland Oregon, November 23-24, pp. 54-66.
- Agee, J. K., and R. Flewelling. 1983. "A fire cycle model based on climate for the Olympic Mountains, Washington." *Fire For. Meteorol. Conf.* 7:32-37.
- Barry, J. 1993. Wellfield Hydrologic Assessment Report. Prepared for the City of Oakridge. CH2M Hill Consulting Engineers. On File: Willamette National Forest, Oakridge Ranger District..
- Behnke, R. J. 1992. Native trout of western North America. American Fisheries Society, Bethesda, Maryland. 275 pp.
- Beschta R. L., and R. L. Taylor. 1988. "Stream temperature increases and land use in a forested Oregon watershed." *Water Resources Bulletin* 24(1):19-25.
- Bisson, P.A., R.E. Bilby, M.D. Bryant, C.A. Dolloff, G.B. Grette, R.A. House, M.L. Murphy, K.V. Koski, and J.R. Sedell. 1987. Large woody debris in forested streams in the Pacific Northwest: past, present, and future. In Salo, E.O. and T.W. Cundy (editors). *Streamside Management: Forestry and Fishery Interactions*. Contribution No. 57. Seattle, Washington: University of Washington, Institute of Forest Resources. pp:143-190.
- Bjornn, T.C. and D. W. Reiser. 1991. "Habitat requirements of salmon in streams." Pages 83-138 in *Influences of forest and rangeland management on salmonid fishes and their habitats*. W.R. Meehan, editor. American Fisheries Society Special Publication. Bethesda, Maryland. 751 pp.
- Black, G.L., D.D. Blackwell and J.L. Steele. 1983. Heat flow in the Oregon Cascades in Priest, G.R. and B.F. Vogt (editors). *Geology and geothermal resources of the central Oregon Cascade Range*. State of Oregon, Dept. of Geology and Mineral Industries. Special Paper 15. pp: 69-76.
- Blaustein, A.R., J.J. Beatty, D. H. Olson, and R.M. Storm. 1995. The biology of amphibians and reptiles in old-growth forests in the Pacific Northwest. USDA Forest Service. Pacific Northwest Research Station. General Technical Report PNW-GTR-337. March 1995. 98 pp.
- Cassirer, E.F., A. Breault, P. Clarkson, D.L. Genter, R.I. Goudie, B. Hunt, S.C. Latta, G.H. Mittlehauser, M. McCullough, G. Schirato, and R.L. Wallen. 1993. Status of harlequin ducks (*Histrionicus histrionicus*). Report of the Harlequin Duck Working Group.
- Christner, J., and Harr, R. D. 1982. "Peak streamflow from the transient snow zone, western Cascades, Oregon." In *Western Snow Conference*, pp. 27-38. Reno.
- Corkran, C. C. and C. Thoms. 1995. Field manual of Oregon and Washington amphibians. 128 pp. Northwest Ecological Research Institute.
- Duncan, J.R. and P.A. Hayward. 1994. Review of technical knowledge: great gray owls. In *Flammulated, boreal, and great gray owls in the United States: a technical conservation assessment*. General Technical Report RM-253. Forest Service, Rocky Mountain Forest and Range Experimental Station, Ft. Collins, Co.
- Fahnestock, G. R., and J. K. Agee. 1983. "Biomass consumption and smoke production by prehistoric and modern forest fires in western Washington." *Journal of Forestry* 81:653-657.
- Foltz, R. B. 1991. "Sediment production comparison between the use of normal tire pressure and central tire pressure on forest roads." Paper presented at Forestry and Environment Engineering Solutions, American Society of Agricultural Engineers. June 5-6, New Orleans, Louisiana.

SALT CREEK WATERSHED ANALYSIS

- Franklin, J. F., and C.T. Dyrness. 1988. Natural vegetation of Oregon and Washington. Oregon State University Press.
- Goggans, R. 1986. Habitat use by Flammulated owls in northeastern Oregon. Unpubl. MS. Thesis. Oregon State University, Corvallis, OR.
- Groves, C.R. 1988. Distribution of the wolverine in Idaho as determined by mail questionnaire. Northwest Science. 62:181-185.
- Guenther, K. and T.E. Kucera. 1978. Wildlife of the Pacific Northwest: Occurrence and distribution by habitat, BLM district, and national forest. USDA Forest Service, Pacific Northwest Region. 128 pp.
- Harr, R.D. 1981. "Some characteristics and consequences of snowmelt during rainfall in western Oregon." Journal of Hydrology 53:277-304.
- Harr R. D. 1983. "Potential for augmenting water yield through forest practices in western Washington and western Oregon." Water Resources Bulletin 19(3):383-393.
- Harr R. D., A. Levno, and R. Mersereau. 1982. "Streamflow changes after logging 130-year-old Douglas-fir in two small watersheds." Water Resources Research 18(3):637-644.
- Harris, D.D., L.L. Hubbard and L.E. Hubbard. 1979. Magnitude and frequency of floods in western Oregon. U.S. Dept. of the Interior, U.S. Geological Survey. Open File Report 79-553.
- Hemstrom, M. A., and J. F. Franklin. 1982. "Fire and other disturbances of the forests in Mount Rainier National Park." Quat. Res. 18:32-51.
- Hemstrom, M. A., S. E. Logan, and W. Pavlat. 1987. Plant association and management guide. USDA Forest Service, Pacific Northwest Region, Willamette National Forest. R6-Ecol-257-B-86.
- Hornocker, M.G. and H.S. Hash. 1981. Ecology of the wolverine in northwestern Montana. Canadian Journal of Zoology 59.
- Huff, M. H., and J. K. Agee. 1980. "Characteristics of large lightning fires in the Olympic Mountains, Washington." Fire For. Meteorol. Conf. 6:117-123.
- Hutchinson, J. M., K. E. Thompson, and J. D. Fortune, Jr. 1966. Basin investigations - Upper Willamette basin. Oregon State Game Commission, Portland Oregon. 40pp.
- Johnson, Al. 1997 Hydrologist, Oakridge Ranger District. Personal Communication
- Johnson D. R., 1917. Spawning ground studies of Willamette River spring CH salmon. Oregon Fish Commission.
- Jones, J. A. and G. E. Grant. 1996. "Peak flow responses to clearcutting and roads in small and large basins, western Cascades, Oregon." Water Resources Research Volume 32.
- Leverton, Mark 1996. Geologist, Oakridge Ranger District. Personal Communication
- Liss, W. J., 1991. "Ecological effects of stocked fish on naturally fishless high mountain lakes of the North Cascades National Park. Final Report of The U. S. National Park Service." Oregon State University. Corvallis, Oregon.
- Mannan, R. W., R. N. Conner, B. Marcot, and J. M. Peck. 1994. "Managing forestlands for wildlife." Pages 689-721 in Research and management techniques for wildlife and habitats. T.A. Bookhout, editor. Fifth edition. The Wildlife Society, Bethesda, MD.

SALT CREEK WATERSHED ANALYSIS

- MacDonald, L. H., W. W. Smart, and R. C. Wissmar. 1991. Monitoring guidelines to evaluate effects of forestry activities on streams in the Pacific Northwest and Alaska. U.S. Environmental Protection Agency Region 10. Seattle, WA. EPA/910/9-91-001.
- McIntosh, B. A., S. E. Clark, and J. R. Sedell. 1992. Summary report for Bureau of Fisheries stream habitat surveys - Willamette River Basin 1934-1942. USDA Forest Service, Oregon State University. Corvallis, Oregon. 476 pp.
- McIntosh, B., S. Clark, J. Sedell, B. Gresswell. 1996. Compilation of stream habitat surveys - Middle Fork of the Willamette Basin. Aquatic Inventory Project, Research and Development Section, Oregon Department of Fish and Wildlife, Corvallis, Oregon.
- Megahan, W.F. 1992. A cross section of procedures used to assess cumulative watershed effects on forest lands in the U.S. in Neary et al. National Hydrology Workshop Proceedings. RM GTR-279
- Mellen, K., M. Huff and R. Hagedstedt. 1994. Habscapes Reference Manual and User's Guide
- Miller, Glen. 1993. Personal communication. Oregon Department of Agriculture, Lowell, Oregon.
- Moring, J. R. 1975. The Alsea Watershed Study: Effects of logging on the aquatic resources of three headwater streams of the Alsea River, Oregon Part II - Changes in environmental conditions. Fishery Research Report Number 9. Oregon Department of Fish and Wildlife. Corvallis, Oregon.
- Nussbaum, R.A., E.D. Brodie and R.M. Storm. 1983. Amphibians and Reptiles of the Pacific Northwest. University Press of Idaho, A Division of Idaho Research Foundation, Inc.
- Oregon Department of Agriculture. 1995. Noxious weed policy and classification system.
- Oregon Department of Fish and Wildlife. 1996. Personal Communication. Mark Wade, Fish Biologist. Springfield, Oregon.
- Pickford, S. D., G. Fahnestock, and R. Ottmar. 1980. "Weather, fuels, and lightning fires in Olympic National Park." Northwest Sci. 54:92-105
- Potts, D. F., D. L. Peterson, H.R. Zuuring. 1989. Estimating postfire water production in the Pacific Northwest. USDA Forest Service. Pacific Southwest Forest and Range Experiment Station. Research Paper PSW-197.
- Rosgen, D.L. 1994. A classification of natural rivers. *Catena* 22:169-199.
- Schroer, G. 1994. Roosevelt elk selection of temperate rain forest seral stages in Western Washington. *Northwest Science*, vol 67, no. 1 pp. 23-29
- Sherrod, Geologic map of a part of the Cascade Range between latitudes 43(-44), central Oregon. U.S. Dept. of the Interior, U.S. Geological Survey.
- Swanson, Nikki. 1997. Fisheries Biologist, Oakridge Ranger District. Personal Communication.
- Taylor, Ronald J. 1990. Northwest weeds, the ugly and beautiful villains of fields, gardens, and roadsides. 177 pgs. Mountain Press Publishing Company.
- Thomas, J. W., and D. E. Towell. 1982. Elk of North America: ecology and management. Stackpole Books, Harrisburg, PA. 698 pp.
- USDA Forest Service. 1990. Land and resource management plan. Willamette National Forest. USDA Forest Service, Pacific Northwest Region.

SALT CREEK WATERSHED ANALYSIS

- USDA Forest Service, Pacific Northwest Region. 1993. A first approximation of ecosystem health on National Forest system lands.
- USDA Forest Service. 1993a. Integrated weed and management plan. Willamette National Forest, Eugene, Oregon.
- USDA Forest Service. 1994a. Soil resource inventory data dictionary. Willamette National Forest.
- USDA Forest Service. 1994b. Soil resource inventory, Willamette National Forest
- USDA Forest Service. 1994c. Willamette National Forest special forest products environmental assessment. Pacific Northwest Region.
- USDA Forest Service. 1995. Willamette National Forest wildlife habitat data base, 4/28/95
- USDA et al. 1993. Forest ecosystem management: An ecological, economic, and social assessment. Report of the Forest Ecosystem Management Assessment Team.
- USDA Forest Service, USDI Bureau of Land Management. 1993. Conservation strategy, *Frasera umpquaensis*. USFS Pacific Northwest Region, Umpqua, Rogue River, Willamette, and Siskiyou National Forests; USBLM, Medford and Eugene Districts.
- USDA Forest Service, USDI Bureau of Land Management. 1994a. Record of decision for amendments to Forest Service and Bureau of Land Management planning documents within the range of the northern spotted owl - Standards and guidelines for management of habitat for late-successional and old-growth forest related species within the range of the northern spotted owl.
- USDA Forest Service, USDI Bureau of Land Management. 1994b. Final supplemental environmental impact statement on management of habitat for late-successional and old-growth forest related species within the range of the northern spotted owl, Volume I. February, 1994.
- USDA Forest Service, USDI Bureau of Land Management. 1994c. Environmental assessment for the implementation of interior strategies for managing anadromous fish-producing watersheds in eastern Oregon and Washington, Idaho, and portions of California.
- USDA Forest Service/USDI Bureau of Land Management. 1995. Ecosystem analysis at the watershed scale, version 2.2 (The revised federal guide for watershed analysis). United States Department of Agricultural and United States Department of the Interior, Portland Oregon, August 28, 1995.
- USFWS, 1982. The Pacific Coast American peregrine falcon recovery plan. USDI Fish and Wildlife Service. Portland, OR. 87 pp.
- Wade, M. A. and J. S. Ziller. 1995. Upper Willamette fish district stock status review; April 11, 1995. Oregon Department of Fish and Wildlife. Springfield, Oregon.
- Waananen, A.O., D.D. Harris and R.C. Williams. 1971. Floods of December 1964 and January 1965 in the far western states. Part 1. Description. U.S. Dept. of the Interior, U.S. Geological Survey Water-Supply Paper 1866-A.
- Welcher, K. A., M. Beilharz, K. Beyer, D. T. Bates. 1991. Water Quality and Quantity at U.S. Geological Survey Stations, Willamette National Forest. Pacific Northwest Region. On file: U.S.D.A. Forest Service, Willamette National Forest Supervisor's Office, Eugene, OR.
- Wemple, B. C. 1994. "Hydrologic integration of forest roads with stream networks in two basins, western Cascades, Oregon." M.S. Thesis, Oregon State University. Corvallis, Oregon.
- Willamette Basin Task Force, 1969. The Willamette Basin, A Comprehensive Study of Water and Related Land Resources. Pacific Northwest Basin Commission.

SALT CREEK WATERSHED ANALYSIS

Woller, N.M. and G.L. Black. 1983. Geology of the Waldo Lake-Swift Creek area, Lane and Klamath counties, Oregon in Priest, G.R. and B.F. Vogt (editors). Geology and geothermal resources of the central Oregon Cascade Range. State of Oregon, Dept. of Geology and Mineral Industries. Special Paper 15. pp: 57-68.

Wolman, M.G. 1954. A method of sampling coarse river-bed material. Trans. Amer. Geophys. Union 35(6):951-956.

Wydoski, R.S. and R.R. Whitney. 1979. Inland Fishes of Washington. University of Washington Press.

Appendix L: Glossary

Many of the definitions in this glossary are referenced to the following sources. The sources are identified by a number in parentheses following the definition. This number corresponds to the list below. Some other terms will be referenced to Forest Service Manuals (FSM), Forest Service Handbooks (FSH), or other sources which are too numerous to list. Finally, many other definitions are not referenced, but are those in general use on the Forest.

Source List

- 1) CFR 219 National Forest Management Act Regulations.
- 2) Regional Guide for the Pacific Northwest Region, 1984.
- 3) SAF Dictionary of Forestry Terms, 1971.
- 4) The Random House College Dictionary, Revised Edition, 1975.
- 5) Webster's New International Dictionary, 1957.
- 6) Wildland Planning Glossary, 1976.
- 7) Webster's Third New International Dictionary, 1981.
- 8) Wildlife Habitats in Managed Forests, The Blue Mountains of Oregon and Washington, 1979.
- 9) A Glossary of Terms Used in Range Management.
- 10) Forest Service Manual or Forest Service Handbook.

- A -

Age class - An interval, usually 10 to 20 years, into which the age ranges of vegetation are divided for classification or use. (3)

Aggregate Recovery Percent (ARP) - Measure of the vegetative condition related to its ability to intercept rain, snow, and wind and its ability to modify snow accumulation and melting.

Airshed - A geographic area that, because of topography, meteorology, and climate, shares the same air. (2)

Alluvium, Alluvial - Sediments deposited by water.

Anadromous Fish - Those species of fish that mature in the sea and migrate into streams to spawn. Salmon, steelhead, and searun cutthroat trout are examples.

Andesite - A moderately hard light colored rock produced by volcanic eruption.

Appropriated Funds - Moneys authorized by an act of Congress which permit Federal agencies to incur obligations and to make payments out of the US Treasury for specified purposes.

Aquifer - Underground strata containing water.

Aquatic ecosystems - Stream channels, lakes, marshes or ponds, and the plant and animal communities they support.

Artifact - An object made or modified by humans. (4)

Available forest land - Land which has not been legislatively or administratively withdrawn by the Secretary of Agriculture or Forest Service Chief from timber production.

- B -

Background - In visual management terminology, refers to the visible terrain beyond the foreground and middleground where individual trees are not visible, but are blended into the total fabric of the stand. Also a portion of a view beyond three to five miles from the observer, and as far as the eye can detect objects. (6)

Bald Eagle Management Area (BEMA) - An area allocated by the Willamette National Forest Plan to be managed for the benefit of American Bald Eagles.

Basaltic - A hard generally dark and dense rock type produced by volcanic eruption.

Base Flow - The portion of a stream or river flow attributable to ground water interception, usually a very constant amount.

Bedload - The coarse sediment moved by a stream or river which moves along the bed of the stream.

Beneficial uses - In water use law the reasonable use of water for a purpose consistent with the laws and best interest of the people and the state.

Best Management Practices - A practice or combination of practices that is determined by a State (or designated areawide planning agency) after problem assessment, examination of alternative practices, and appropriate public participation, to be the most effective, practicable (including technological, economic, and institutional considerations) means of preventing or reducing the amount of pollution generated by nonpoint sources to a level compatible with water quality goals (Federal Register, Volume 40, No. 230 dated 11/28/75).

Big game - Large mammals hunted for sport. On the National Forest these include animals such as deer, elk, antelope, and bear. (8)

Big Game Emphasis Area (BGEA) - An area of land designated by the Willamette National Forest Plan with prescriptions for specific habitat qualities for deer and elk.

Big game summer range - A range, usually at higher elevation, used by deer and elk during the summer. Summer ranges are usually much more extensive than winter ranges. (8)

Big game winter range - A range, usually at lower elevation, used by migratory deer and elk during the winter months; usually more clearly defined and smaller than summer ranges. (8)

Biological diversity - Terms used in the Forest Plan to provide goals and direction for evaluating the significance of old growth stands, minimizing fragmentation of existing old growth forests, and maintaining many of the structural components of unmanaged stands in managed stands.

Board foot (BF) - The amount of wood equivalent to a piece of wood one foot by one foot by one inch thick. (3)

Broadcast Burn - Allowing a prescribed fire to burn over a designated area within well-defined boundaries for reduction of fuel hazard or as a silvicultural treatment, or both.

Browse - Twigs, leaves, and young shoots of trees and shrubs on which animals feed; in particular, those shrubs which are used by big game animals for food. (6)

Browse Enhancement - The act of cutting down brush or hardwood vegetation when it is too tall, decadent, or low in nutritional value to increase its future value to browsing animals, usually big game. This cutback allows the vegetation to resprout and become more available and or higher quality.

- C -

Canopy - The more-or-less continuous cover of branches and foliage formed collectively by the crown of adjacent trees and other woody growth. (3)

Cavity - The hollow excavated in trees by birds or other natural phenomena; used for roosting and reproduction by many birds and mammals. (2)

Char - A group of fish in the Salmonid family - in this watershed, brook trout and bull trout.

Clearcutting - The cutting method that describes the silviculture system in which the old crop is cleared over a considerable area at one time. Regeneration then occurs from (a) natural seeding from adjacent stands, (b) seed contained in the slash or logging debris, (c) advance growth, or (d) planting or direct seeding. An even-aged forest usually results. (3)

Climax - The culminating stage in plant succession for a given site where the vegetation has reached a highly stable condition. (6)

Collurturn - Material (soil and rock) that has been deposited through gravity (as opposed to water).

Commercial Forest Land - Land that is producing, or is capable of producing, crops of industrial wood and (1) has not been withdrawn by Congress, the Secretary of Agriculture, or the Chief of the Forest Service; (2) land where existing technology and knowledge is available to ensure timber production without irreversible damage to soil productivity or watershed conditions; and (3) land where existing technology and knowledge, as reflected in current research and experience, provides reasonable assurance that adequate restocking can be obtained within 5 years after final harvesting. See also "Tentatively Suitable Forest Land."

Commercial thinning - Any type of tree thinning that produces merchantable material at least equal in value to the direct costs of harvesting. (3)

Compaction - The packing together of soil particles by forces exerted at the soil surface, resulting in increased soil density.

Conk - The woody fruiting body of fungal species, that usually grow on dead or live tree stems.

Connectivity - A measure of the extent to which conditions among late-successional and/or old growth (LS/OG) areas provide habitat for breeding, feeding, dispersal, and movement of LS/OG associated wildlife and fish species.

Corridor - A linear strip of land identified for the present or future location of transportation or utility rights-of-way within its boundaries. (1)

Course sediment - Sands, gravels, cobbles, boulders.

Cultural resource - The remains of sites, structures, or objects used by humans in the past--historic or prehistoric. (2)

Cumulative effects or impacts - Cumulative effect or impact is the impact on the environment which results from the incremental impact of the action when added to other past, present, and reasonably foreseeable future actions, regardless of what agency (federal or nonfederal) or person undertakes such other actions. Cumulative impacts can result from individually minor but collectively significant actions taking place over a period of time. (40 CFR 1508.7 - these regulations use effects and impacts synonymously.)

- D -

Debris Torrent - A large debris slide that is charged with water and confined to a steep stream channel. Debris torrents may travel several thousand feet, but are generally shallow as opposed to deep-seated mass movement.

Deep-seated Mass Movement - The downhill movement of deep soils and weathered bedrock, usually under saturated conditions. Such events usually do not move as far as a debris torrents do.

Designated Area (Air Quality) - Those areas delineated in the Oregon and Washington Smoke Management Plans as principal population centers of air quality concern.

Developed recreation - Recreation that requires facilities that, in turn, result in concentrated use of an area. Examples of developed recreation areas are campgrounds and ski areas; facilities in these areas might include roads, parking lots, picnic tables, toilets, drinking water, ski lifts, and buildings. (2)

Diameter at breast height (d.b.h.) - The diameter of a tree measured 4 feet 6 inches above the ground. (6)

Dispersed recreation - A general term referring to recreation use outside developed recreation sites; this includes activities such as scenic driving, hiking, backpacking, hunting, fishing, snowmobiling, horseback riding, cross-country skiing, and recreation in primitive environments. (2)

Diversity - The distribution and abundance of different plant and animal communities and species within the area covered by a land and resource management plan. (2) (1)

Douglas-Fir Type - An association of tree species in which Douglas-fir is recognized as one of the principal seral species.

Duff - Organic matter in various stages of decomposition on the floor of the forest. (4)

- E -

Edge - An area where plant communities meet or where successional stages or vegetation conditions within the plant communities come together. (2)

Effects - Environmental changes resulting from a proposed action. Included are direct effects, which are caused by the action and occur at the same time and place, and indirect effects, which are caused by the action and are later in time or further removed in distance, but which are still reasonably foreseeable. Indirect effects may include growth-inducing effects and other effects related to induced changes in the pattern of land use, population density, or growth rate, and related effects on air and water and other natural systems, including ecosystems. Effects and impacts as used in this document are synonymous. Effects include ecological (such as the effects on natural resources and on the components, structures, and functioning of affected ecosystems), aesthetic quality, historic, cultural, economic, social, or healthy effects, whether direct, indirect, or cumulative. Effects may also include those resulting from actions that may have both beneficial and detrimental effects, even if on balance the agency believes that the effects will be beneficial. (40 CFR 1508.8, 2)

Ejecta - Material expelled forcibly from an erupting volcano, as opposed to lava flows.

Endangered species - Any species of animal or plant that is in danger of extinction throughout all or a significant portion of its range. Plant or animal species identified by the Secretary of the Interior as endangered in accordance with the 1973 Endangered Species Act.(6)

Environmental Analysis - A comprehensive evaluation of alternative actions and their predictable short- and long-term environmental effects, which include physical, biological, economic, social, and environmental design factors and their interactions. (2)

SALT CREEK WATERSHED ANALYSIS

Environmental Assessment - The concise public document required by the regulations for implementing the procedural requirements of the National Environmental Policy Act. (40 CFR 1508.9, 2)

Environmental Impact Statement (EIS) - A statement of the environmental effects of a proposed action and alternatives to it. It is required for major federal actions under Section 102 of the National Environmental Policy Act (NEPA), and released to the public and other agencies for comment and review. It is a formal document that must follow the requirements of NEPA, the Council on Environmental Quality (CEQ) guidelines, and directives of the agency responsible for the project proposal. (6)

Ephemeral draw - A drainage way which conveys surface water for short periods of time in direct response to snowmelt or rainfall runoff.

Even-aged stands - Stands in which all trees are of about the same age. (A spread of 10 to 20 years is generally considered one age class.) Cutting methods producing even-aged stands are clearcut, shelterwood, or seed tree systems.

- F -

Fire management - All activities required for protection of resources from fire and for the use of fire to meet land management goals and objectives. (6)

Fire return interval - The length of time between major, landscape level, stand replacement fire occurrences within a watershed or other large landscape. This term does not apply to a given acre and does not indicate the maximum age that forests attain in the area. It is simply an indication of the periodicity of large fires in the watershed.

Fire rotation - The time period between stand-replacing fire events on a given acre, stand, or site. While this figure may be most accurately used as an average of the periods between stand replacing fires, it is most frequently used to refer to the time between the last two events since dates of all fires which have affected a given site are usually not known.

Fisheries habitats - Streams, lakes, and reservoirs that support fish populations.

Flood plain - The lowland and relatively flat area adjoining inland waters, including, at a minimum, that area subject to a one percent or greater chance of flooding in any given year. (2)

Floristic - Relating to flowering plants.

Forage - All browse and nonwoody plants that are available to livestock or game animals and used for grazing or harvested for feeding. (6)

Foreground - A term used in visual management to describe the portions of a view between the observer and up to 1/4 to 1/2 mile distant. (6)

Forest system roads - Roads that are part of the Forest development transportation system, which includes all existing and planned roads as well as other special and terminal facilities designated as Forest development transportation facilities. (See arterial roads, collector roads, and local roads.)

Fuel management - The practice of planning and executing the treatment or control of living or dead vegetative material in accordance with fire management direction. (10)

Fuel treatment - The rearrangement or disposal of natural or activity fuels (generated by management activity, such as slash left from logging) to reduce fire hazard. Fuels are defined as both living and dead vegetative materials consumable by fire.

Fuels - Combustible wildland vegetative materials. While usually applied to above ground living and dead surface vegetation, this definition also includes roots and organic soils such as peat. (10)

SALT CREEK WATERSHED ANALYSIS

- G -

Geomorphic - The formation of geologic and topographic features.

Glaciation - Erosion and deposition of soil and rocks by movement of glacial ice.

Guilds, Guilding - Classes of wildlife relating to their habits and environment.

Group selection cutting - See Uneven-aged silvicultural systems.

- H -

Habitat Effectiveness Indices - A numerical quantification of various big-game habitat qualities.

Headwaters - The upper tributaries of a river. (4)

Herpetile - Any amphibian or reptile.

Hiding cover - Vegetation that will hide 90 percent of an adult deer or elk from the view of a human at a distance of 200 feet or less. The distance at which the animal is essentially hidden is called a "sight distance."

Historic site - Site associated with the history, tradition, or cultural heritage of national, state, or local interest, and of enough significance to merit preservation or restoration. (6)

Hydrology - The scientific study of the properties distribution and effects of water in the atmosphere, on the earth's surface, and in soil and rocks.

- I -

ID Team - See Interdisciplinary team.

Impacts - See Effects.

Indicator species - See Management indicator species.

Infrastructure - The collection of facilities (roads, campgrounds, structures, transportation corridors, power transmission lines, antenna) constructed to facilitate administration of land.

Interdisciplinary Team - A team of specialists talking and debating from the viewpoint of all relevant resources throughout scoping, developing alternatives, estimating effects, and comparing alternatives as a part of the NEPA environmental analysis process.

Interior habitat - Forest habitat that is not affected by adjacent non-forest or young forest. Forest habitat with no edge effects.

Intermittent Stream - A stream that runs water in most months, but does not run water during the dry season during most years.

Issue - A point, matter, or question of public discussion or interest to be addressed or decided through the planning process. (2)

- J,K -

Key Watershed - A Watershed containing populations of species at risk, containing potential habitat, or especially high quality water as designated by the Northwest Forest Plan.

- L -

Landing - Any place where round timber is assembled for further transport, commonly with a change of method. (3)

Lands Not Suited (Unsuitable) for Timber Production - Includes lands that: 1) are not forest land as defined in CFR 219.3; 2) are likely, given current technology, to suffer irreversible resource damage to soils productivity, or watershed conditions; 3) cannot be adequately restocked as provided in 36 CFR 219.27(c) (3); or, 4) have been withdrawn from timber production by an Act of Congress, the Secretary of Agriculture, or the Chief of the Forest Service. In addition, Forest lands other than those that have been identified as not suited for timber production shall be reviewed and assessed prior to formulation of alternatives to determine the costs and benefits of a range of management intensities for timber production. (1)

Landtype - A portion of the Forest mapped in the Soil Resource Inventory that has a defined arrangement of specific landforms that reacts to management activities in generally predictable ways. Landtypes range from 60 to 600 acres in size.

Large woody material (LWM) - Fallen large trees in streams or on the ground in terrestrial environments.

Late Successional - A vegetation type, usually forest, that is mature or old. Also old-growth.

Late Successional Reserve (LSR) - An area set aside from harvest and road building for species requiring late-succession habitat or interior habitat. An allocation specified for implementing the Northwest Forest Plan.

Lichens - Any of a large group of plants consisting of symbiotic fungi and algae.

Lithology, Lithologic - relating to rocks.

Low Flow - Minimum stream flows in summer or fall.

- M -

Management Area - An area with similar management objectives and a common management prescription. (1) (10)

Management direction - A statement of multiple use and other goals and objectives, and the associated management prescriptions, and standards and guidelines for attaining them. (1)

Management indicator species - A species selected because its welfare is presumed to be an indicator of the welfare of other species using the same habitat. A species whose condition can be used to assess the impacts of management actions on a particular area. (8)

Mass movement - A general term for any of the variety of processes by which large masses of earth material are moved downslope by gravitational forces - either slowly or quickly. (6) See Debris torrent and Deep-seated mass movement.

Mass wasting - Mass movement.

Matrix - That land outside of various reserves which is to be managed for timber production, among other objectives; designated by the Northwest Forest Plan.

Mature timber - Trees that have attained full development, particularly height, and are in full seed production. (3)

SALT CREEK WATERSHED ANALYSIS

Maximum modification - See Visual quality objective.

Mesic - Moist, referring to a soil or site.

Middleground - A term used in visual management to describe the portions of a view extending from the foreground zone out to 3 to 5 miles from the observer. (6)

Mineral soil - Weathered rock materials usually containing less than 20 percent organic matter. (6)

Mitigation - Mitigation includes: (a) avoiding the impact altogether by not taking a certain action or parts of an action; (b) minimizing impacts by limiting the degree or magnitude of the action and its implementation; (c) rectifying the impact by repairing, rehabilitating, or restoring the affected environment; (d) reducing or elimination the impact over time by preservation and maintenance operations during the life of the action; and, (e) compensating for the impact by replacing or providing substitute resources or environments. (40 CFR Part 1508.20)

Modification - See Visual quality objective.

Monitoring and evaluation - The periodic evaluation of Forest Plan management practices on a sample basis to determine how well objectives have been met.

Morphometry - Measurement of the morphology or form, as in lake bottom shapes.

Municipal Watershed - A watershed which provides water for human consumption, where Forest Service management could have a significant effect on the quality of water at the intake point, and that provides water utilized by a community or any other water system that regularly serves: 1) at least 25 people on at least 60 days in a year, or 2) at least 15 service connections. In addition to cities, this includes campgrounds, residential developments, and restaurants. (10)

- N -

National Environmental Policy Act (NEPA) of 1969 - An Act to declare a National policy which will encourage productive and enjoyable harmony between humankind and the environment, to promote efforts which will prevent or eliminate damage to the environment and biosphere and stimulate the health and welfare of humanity, to enrich the understanding of the ecological systems and natural resources important to the Nation, and to establish a Council on Environmental Quality. (The Principal Laws Relating to Forest Service Activities, Agriculture Handbook No. 453, USDA, Forest Service, 359 pp.)

National Forest Management Act (NFMA) - A law passed in 1976 as an amendment to the Forest and Rangeland Renewable Resources Planning Act, requiring the preparation of Regional Guides and Forest Plans and the preparation of regulations to guide that development.

Natural regeneration - Reforestation of a site by natural seeding from the surrounding trees. Natural regeneration may or may not be preceded by site preparation.

Nephelometric Turbidity Unit - A relative quantification of water turbidity.

Nonpoint source pollution - Pollution whose source is general rather than specific in location. It is widely used in reference to agricultural and related pollutants-- for example, production of sediments by logging operations, agricultural pesticide applications, or automobile exhaust pollution. (6)

Northwest Forest Plan - Final supplemental environmental impact statement on management of habitat for late-successional and old-growth forest related species within the range of the northern spotted owl, Volume I, February 1994. Sometimes called the "President's Plan."

SALT CREEK WATERSHED ANALYSIS

Noxious weeds - Undesirable, usually non-native, plant species that are unwholesome to the range or to animals or compete with native plants. (6)

- O -

Objective - A concise, time-specific statement of measurable planned results that respond to pre-established goals. An objective forms the basis for further planning to define the precise steps to be taken and the resources to be used in achieving identified goals. (1)

Occasion (recreation) - one person involved in recreation activity.

Old-growth stand (old growth) - Any stand of trees 10 acres or greater generally containing the following characteristics: contain mature and overmature trees in the overstory and are well into the mature growth stage; 2) will usually contain a multilayered canopy and trees of several age classes; 3) standing dead trees and down material are present; and 4) evidences of man's activities may be present, but do not significantly alter the other characteristics and would be a subordinate factor in a description of such a stand. (2)

Oligotrophic - Referring to bodies of water, a condition of very low levels of dissolved or suspended nutrients.

Optimal cover - Habitat for deer and elk which has tree overstory and understory, shrub and herbaceous layers; the overstory canopy generally exceeding 70% crown closure and dominant trees generally exceed 21 inches d.b.h.; provides snow intercept, thermal cover, and forage.

Overstory - That portion of the trees, in a Forest or in a stand of more than one story, forming the upper or uppermost canopy. (3)

- P -

PAOT - Persons at one time.

Partial retention - See Visual quality objective.

Particulates - Small particles suspended in the air and generally considered pollutants. (5)

Perennial stream - A stream that flows year round.

Peak flow - The highest amount of stream or river flow accruing in a year or from a single storm event.

Pests - Any animal or plant that, during some portion of its life cycle, inhibits the establishment or growth of some other species of plant or animal favored by man.

Phonology - The science dealing with the influence of climate on the recurrence of such annual phenomena of animal and plant life as bird migrations, budding, etc. (4)

Planning area - The area of the National Forest System covered by a Regional guide or forest plan. (1)

PM10 emissions - Particulate Matter smaller than 10 micrometers in size. A criteria pollutant comprised of airborne solid and liquid particles that are 10 micrometers or smaller in size. Because of its small size, PM10 readily lodges in the lungs, thus increasing respiratory and cardiac diseases in humans and other organisms.

Prehistoric site - An area which contains important evidence and remains of the life and activities of early societies which did not record their history.

SALT CREEK WATERSHED ANALYSIS

Prescribed fire - A wildland fire burning under specified conditions which will accomplish certain planned objectives. The fire may result from either planned or unplanned ignitions. Proposals for use of unplanned ignitions for this purpose must be approved by the Regional Forester. (2)

Prescription - A written direction for harvest activities and regeneration methods.

Pre-commercial Thinning - See Young Stand Density Management.

Primary cavity excavators - Wildlife species that excavate cavities in snags.

Primary productivity - the portion of biological activity and production attributed to plant life.

Pruning - Removing of limbs from the lower portion of a tree.

Public Involvement - A Forest Service process designed to broaden the information base upon which agency decisions are made by (1) informing the public about Forest Service activities, plan, and decisions, and (2) encouraging public understanding about and participation in the planning processes which lead to final decision making. (10)

Pumice - A light, frothy volcanic rock formed by explosive eruptions.

Pyroclastic - Rock formed in volcanic eruptions that is composed of broken fragments.

- Q -

Quark - Smallest subatomic particle known to man.

- R -

Raptors - Predatory birds, such as falcons, hawks, eagles, or owls.

Reforestation - The natural or artificial restocking of an area with forest trees. (2)

Regeneration - The renewal of a tree crop, whether by natural or artificial means. Also, the young crop itself, which is commonly referred to as reproduction. (2)

Rehabilitation - Action taken to restore, protect, or enhance site productivity, water quality, or other resource values over a period of time .

Release - The cutting of competing and unwanted vegetation to free conifers for growth.

Residual stand - The trees remaining standing after some activity such as selection cutting or an occurrence such as fire or windthrow. (2)

Retention - See Visual quality objective.

Riparian - Pertaining to areas of land directly influenced by water or influencing water. Riparian areas usually have visible vegetative or physical characteristics reflecting this water influence. Stream sides, lake borders, or marshes are typical riparian areas. (3)

Riparian Reserve - A protected area along streams and wetlands.

Roadless Area - Areas studied during the Roadless Area Review and Evaluation process (RARE II) which are roadless and at least 5,000 acres in size.

SALT CREEK WATERSHED ANALYSIS

Rotation - Planned number of years between the formation of a generation of trees and its final harvest at a specified stage of maturity. Appropriate for even-aged management only. (6)

Rotational failure - A general term for a mass movement landform and a process characterized by a slope in which shearing takes place on a well defined, curved shear surface, concave upward, producing a backward rotation in the displaced mass. The landform may be single, successive (repeated up- and down-slope), or multiple (as the number of slide components increases).

RVD - Recreation Visitor Day; one RVD equals 12 visitor hours which may be aggregated continuously, intermittently, or simultaneously by one or more persons.

- S -

Salmonid - The family of fish species including salmon, trout, and char (whitefish).

Salvage cuttings - Intermediate cuttings made to remove trees that are dead or in imminent danger of being killed by injurious agents. (10)

Scarified - Land in which the topsoil has been broken up or loosened in preparation for regenerating by direct seeding or natural seedfall. Also refers to ripping or loosening road surfaces to a specified depth for obliteration or "putting a road to bed." (3)

Second growth - Forest growth that has become established following some interference, such as cutting, serious fire, or insect attack, with the previous Forest crop. (6)

Sediment - Earth material (rocks, gravels, sands, silts, clays) transported, suspended, or deposited by water. (6)

Seed tree cutting - Removal in one cut of the mature timber from an area, except for a small number of seed bearers left singly or in small groups. (3)

Selection cutting - The annual or periodic removal of trees (particularly mature trees), individually or in small groups, from an uneven-aged forest, to realize the yield and establish a new crop of irregular constitution. (3)

Sensitive species - Plant or animal species which are susceptible or vulnerable to activity impacts or habitat alterations. Those species that have appeared in the Federal Register as proposed for classification or are under consideration for official listing as endangered or threatened species, that are on an official State list, or that are recognized by the Regional Forester as needing special management to prevent placement on Federal or State lists. (2)

Sensitivity analysis - A determination of the effects of varying the level of one or more factors, while holding the other factors constant. (6) (10)

Seral - A stage in plant community development.

SHAB, Special Habitats - Areas set aside by the Willamette National Forest Plan to protect unique plant and animal habitats.

Shelterwood - The cutting method that describes the silvicultural system in which, in order to provide a source of seed and/or protection for regeneration, the old crop (the shelterwood) is removed in two or more successive shelterwood cuttings. The first cutting is ordinarily the seed cutting, though it may be preceded by a preparatory cutting, and the last is the final cutting. Any intervening cutting is termed removal cutting. An even-aged stand results. (3)

Silviculture - The art and science of controlling the establishment, composition, and growth of forests. (2)

SALT CREEK WATERSHED ANALYSIS

Site preparation - 1) An activity (such as prescribed burning, disking, and tilling) performed on a reforestation area, before introduction of reforestation, to ensure adequate survival and growth of the future crop; or 2) manipulation of the vegetation or soil of an area prior to planting or seeding. The manipulation follows harvest, wildfire, or construction in order to encourage the growth of favored species. Site preparation may include the application of herbicides; burning, or cutting of living vegetation that competes with the favored species; tilling the soil; or burning of organic debris (usually logging slash) that makes planting or seeding difficult.

Skidding - A general term for hauling loads by sliding, not on wheels, as developed originally from stump to roadside, deck, skidway, or other landing. (3)

Skyline Logging - A system of cable logging in which all or part of the weight of the logs is supported during yarding by a suspended cable.

Slash - The residue left on the ground after tree felling and tending, and/or accumulating there as a result of storm, fire, girdling or poisoning. It includes unutilized logs, uprooted stumps, broken or uprooted stems, the heavier branchwood, etc. (3)

Snag - A standing dead tree.

Soil productivity - The capacity of a soil to produce a specific crop such as fiber or forage under defined levels of management. Productivity is generally dependent on available soil moisture and nutrients, and length of growing season.

Soil resource inventory - See Landtype.

Special Interest Areas - Areas managed to make recreation opportunities available for the understanding of the earth and its geological, historical, archeological, botanical, and memorial features. (6)

Special Forest Products (SFPs) - Forest resources that are not associated with timber sale contracts. May be for commercial or personal use. Some common SFPs include greenery, mushrooms, live plants, cones, berries, etc.

Special Wildlife Habitat - A habitat which is unique and has a special function not provided by plant communities or Successional stages; includes riparian zones, wetlands, cliffs, caves, talus, and meadows.

Stand (tree stand, timber stand) - An aggregation of trees or other vegetation occupying a specific area and sufficiently uniform in species composition, age arrangement, and condition as to be distinguishable from the forest or other vegetation or land cover on adjoining areas. (2)

Stand diversity - Any attribute that makes one timber stand biologically or physically different from other stands. This difference can be measured by, but not limited to: different age classes; species; densities; or non-tree floristic composition.

Stand replacement fire - Fire that kills most or all of a stand of trees, creating space for a new stand to begin.

Standards and Guidelines - Principles specifying conditions or levels of environmental quality to be achieved.

Stream Buffer - Vegetation left along a stream channel to protect the channel or water from the effects of logging, road building, or other management activity.

Stream Class - Classification of streams based on the present and foreseeable uses made of the water, and the potential effects of on-site changes on downstream uses. Four classes are defined:

Class I - Perennial or intermittent streams that: provide a source of water for domestic use; are used by large numbers of anadromous fish or significant sports fish for spawning, rearing or migration; and/or are major tributaries to other Class I streams.

Class II - Perennial or intermittent streams that: are used by fish for

SALT CREEK WATERSHED ANALYSIS

spawning, rearing or migration; and/or may be tributaries to Class I streams or other Class II streams.

Class III - All other perennial streams not meeting higher class criteria.

Class V - All other intermittent streams not meeting higher class criteria. (10)

Stream Structure - The arrangement of logs, boulders, and meanders which modify the flow of water, thereby causing the formation of pools and gravel bars in streams. Generally, there is a direct relationship between complexity of structure and fish habitat. Complex structure is also an indication of watershed stability.

Subdrainage - Areas used for planning and analysis. It is based on tributary drainage boundaries and averaging 2000 to 4000 acres.

Subwatershed - A subdivision of a watershed equivalent to the 6th field subwatersheds as presented in the PACFISH report. These are larger than subdrainages.

Suitability - The appropriateness of applying certain resource management practices to a particular area of land, as determined by an analysis of the economic and environmental consequences and the alternative uses foregone. A unit of land may be suitable for a variety of individual or combined management practices. (1) (2) (FSM 1905)

Succession - A series of changes by which one group of organisms succeeds another through stages leading to a potentially stable climax community.

Suppression - The process of extinguishing or confining fire. (2)

System Road - A road meant to be used in the future with an established maintenance schedule.

- T -

Territory - The area which an animal defends, usually during breeding season, against intruders of its own species.

T. E. and S. species - Threatened, endangered and sensitive species, both plant and animal.

Thermal cover - Cover used by animals to ameliorate effects of weather.

Thinning - A felling made in an immature stand primarily to maintain or accelerate diameter increment and also to improve the average form of the remaining trees without permanently breaking the canopy. An intermediate cutting. (3)

Threatened and Endangered (T&E) species - See Threatened species; see Endangered species.

Threatened species - Those plant or animal species likely to become endangered species throughout all or a significant portion of their range within the foreseeable future. See Endangered species. (2)

Till - An unsorted mixture of clays, silts, sands, gravels and rocks deposited by glaciers.

Tractor logging - Any logging method which uses a tractor as the motive power for transporting logs from the stumps to a collecting point--whether by dragging or carrying the logs. (3)

Transient snow zone - That area where snowfall tends to melt soon after it falls, such that accumulation waxes and wanes through the winter .

Travel Corridor - A route followed by animals along a belt or band of suitable cover or habitat.

Tuff, Tuffaceous - Material made up of volcanic ash deposits.

SALT CREEK WATERSHED ANALYSIS

Turbidity - The degree of opaqueness, or cloudiness, produced in water by suspended particulate matter, either organic or inorganic. Measured by light filtration or transmission and expressed in Jackson Turbidity Units (JTUs).

- U -

Ultra oligotrophic - Very, very clean, clear water.

Underburn - Fire, natural or prescribed, which burns only on the forest floor with an intensity such that dominant trees are typically not killed.

Understory - The trees and other woody species growing under a more-or-less continuous cover of branches and foliage formed collectively by the upper portion of adjacent trees and other woody growth. (6)

- V -

Viewshed - Portion of the Forest that is seen from a major travel route, or high use location.

Visual quality objective (VQO) - Categories of acceptable landscape alteration measured in degrees of deviation from the natural-appearing landscape.

Preservation (P) - Ecological changes only.

Retention (R) - Management activities should not be evident to the casual Forest visitor.

Partial Retention (PR) - Management activities remain visually subordinate to the characteristic landscape.

Modification (M) - Management activities may dominate the characteristic landscape but must, at the same time, follow naturally established form, line, color, and texture. It should appear as a natural occurrence when viewed in foreground or middleground.

Maximum Modification (MM) - Human activity may dominate the characteristic landscape, but should appear as a natural occurrence when viewed as background.

Enhancement - A short-term management alternative which is done with the express purpose of increasing positive visual variety where little variety now exists. (2)

Visual resource - The composite of basic terrain, geologic features, water features, vegetative patterns, and land use effects that typify a land unit and influence the visual appeal the unit may have for visitors. (2)

- W -

Watershed - The entire land area that contributes water to a major drainage system or stream as designated by the FEMAT Report.

Wetlands - Areas that are inundated by surface or ground water often enough to support, and usually do support, primarily plants and animals that require saturated or seasonally saturated soil conditions for growth and reproduction. (E.O. 11990)

Wild and Scenic river - Those rivers or sections of rivers designated as such by Congressional action under the 1968 Wild and Scenic Rivers Act, as supplemented and amended, or those sections of rivers designated as wild, scenic, or recreational by an act of the legislature of the state or states through which they flow. Wild and scenic rivers may be classified and administered under one or more of the following categories:

Wild River Areas - Those rivers or sections of rivers that are free of impoundment's and generally inaccessible except by trail, with watersheds or shorelines essentially primitive and waters unpolluted.

Scenic River Areas - Those rivers or sections of rivers that are free of impoundment's, with watersheds still largely primitive and shorelines largely undeveloped, but accessible in places by roads.

SALT CREEK WATERSHED ANALYSIS

Recreational River Areas - Those rivers or sections of rivers that are readily accessible by road or railroad, that may have some development along their shorelines, and that may have undergone some impoundment or diversion in the past. (2) (6)

WIN, Watershed Improvement needs - A systematic survey of watershed conditions.

Winter Range - An area used by deer and elk during the winter months; usually at lower elevation and/or on south and west exposures.

Woody Material - Organic materials necessary for stream channel stability and maintenance of watershed condition. It includes large logs and root wads.

- X,Y,Z -

Xeric - Dry, referring to soil or site.

Yarding - Hauling timber from the stump to a collection point. (2)

Young Stand Density Management (Pre-commercial Thinning) - Thinning of small trees when no income is derived from the trees and cut trees are generally not removed from the site.

APPENDIX M: Salt Creek Watershed Analysis Team Members

Team Members

David Murdough, Soil Scientist
Barbara Ledbetter, Hydrologic Technician

Team Function

Team Leader
Editor

Terrestrial Group

Dick Davis, Wildlife Biologist
Dede Steele, Wildlife Biologist
Tim Bailey, Silviculturist
Dennis Sullivan, Supervisory Forestry Technician
Mike Williams, Civil Engineering Technician
Kim McMahan, Botanist

Terrestrial Group Leader and Wildlife
Terrestrial Group Leader and Wildlife
Vegetation
Fire
Transportation
Botany

Aquatic Group

Al Johnson, Hydrologist
Nikki Swanson, Fisheries Biologist
Mark Leverton, Geologist

Aquatic Group Leader and Hydrology
Fisheries
Geology

Social Group

Carol Winkler, Archeologist

Cultural and Historical Resources

GIS (Geographic Information Systems) Support

Jim Minogue, GIS Specialist
Karen Meza, GIS Specialist

GIS Coordinator
GIS Analyst

Information Management Support

Debby Murdough, Computer Specialist

Information Coordinator and Computer Support

With Contributions From

Resources Northwest, Inc.

Reference and Current Conditions, Chapter III*
Appendices*

*edits by US Forest Service personnel

APPENDIX N: Public and Agency Contacts

The following individuals, groups, businesses, and/or government agencies provided information during this Watershed Analysis.

State and Federal Agencies:

Oregon Department of Fish and Wildlife; Jeff Ziller, & Bill Castillo

U.S. Forest Service, Willamette Forest

U.S. Geological Survey Water Resources Division; Jo Miller

Oregon Department of Transportation (ODOT)

Oregon Department of Environmental Quality; Roberta Lindberg