

LANDFIRE Biophysical Setting Model

Biophysical Setting 3010800

**Inter-Mountain Basins Big Sagebrush
Shrubland**

☒ This BPS is lumped with: 1125

☐ This BPS is split into multiple models: 1125 describes MZ29 better. 1080 has ARCA13, which doesn't apply in these mapzones. Production is somewhat different, but not enough to split out (Benkobi).

General Information

Contributors (also see the Comments field)

Date 10/3/2006

Modeler 1 Steve Cooper scooper@mt.gov

Reviewer Lakhdar Benkobi lbenkobi@fs.fed.us

Modeler 2

Reviewer Jeff DiBenedetto jdibenedetto@fs.fed.us

Modeler 3

Reviewer George Soehn george_soehn@blm.gov

Vegetation Type

Upland
Savannah/Shrub
Steppe

Dominant Species

ARTRW8
PASM

Map Zone

30

Model Zone

☐ Alaska
☐ California
☒ Northern Plains
☐ N-Cent. Rockies

General Model Sources

☒ Literature
☒ Local Data
☒ Expert Estimate

BOGR2
CHRY9
PSSP6
HECO26
NAVI4
CAFI

☐
☐ Great Basin
☐ Great Lakes
☐ Hawaii
☐ Northeast
☐
☐ Pacific Northwest
☐ South Central
☐ Southeast
☐ S. Appalachians
☐ Southwest

Geographic Range

This system encompasses eastern and central MT, as opposed to throughout the Rocky Mountains, etc as BpS 1125 usually refers to. (This system is lumped with BpS 1125.) 1125 is common throughout MZs 20 and 29 currently (not necessarily historically), except in western part of section 331Da. In MZ29, common historically.

For MZ29, it would occur in northeast WY section 331G, Thunder basin grasslands, northeast of 331Gg.

For MZ29, basin big sagebrush is very uncommon. Have *Artemisia tridentata* ssp. *vaseyana* (BpS 1126) at higher elevations associated with Bighorn, Pryor Mtns and Laramie ranges. Have *Artemisia tridentata* ssp. *wyomingensis* elsewhere where *A. t. ssp. vaseyana* doesn't occur. Mountain big sagebrush occurs in sections M331 associated with Bighorn and Laramie Ranges. *A. t. ssp. wyomingensis* occurs everywhere else.

In MZ29, in southeast MT, but this could be due to a soil anomaly. It probably occurred historically all through the subsections of southeast MT. Also through MZ30 in 331Mi in western Dakotas, 331Md in lower portion. As move north in 331Md, there is less of it. Probably does not occur in 331Mc. Canopy cover of sagebrush is probably <10%.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Biophysical Site Description

This system is Great Plains Sagebrush Steppe for MZ20. For MZ29, we are describing sagebrush wheatgrass steppe, where western wheatgrass is dominant. MZs 20 and 29 are very similar for this type.

Soils are primarily dry from sedimentary processes in this system; soils are less fertile in this system, sometimes more calcareous. The Great Plains expression is found exclusively on "heavy" textured soils derived from shale and mudstones and can be strongly correlated with particular geologic formation or members thereof.

April, May and June have by far the most precipitation and this peaks in late May, early June. This pattern carries throughout the MT portion of the Great Plains though a gradient of more summer precipitation as you progress eastward but still the "spring" peak. It's not until you encounter tallgrass prairie does summer precipitation become predominant.

Wyoming big sagebrush occupies plains, foothills, terraces, slopes, plateaus, basin edges and even lower mountain slopes due to the fact that *Artemisia tridentata* ssp. *vaseyana* is not part of the mix in MZ20 nor in MZ29. Soils are shallow to moderately deep, moderate to well drained and almost exclusively fine textured soils. Wyoming big sagebrush generally occurs in the 5-15in precipitation zones. Soil depth and accumulation of snow enhances these communities in lower precipitation zones (Knight 1994).

In MZ29, *A. t.* ssp. *wyomingensis* can occur from 2200-8000ft.

Bluebunch/ARTR-wyomingensis type is probably an inclusion in this BpS occurring on steep, south aspect slopes, typically badlands slopes/topography.

Vegetation Description

Wyoming big sagebrush is the dominant mid-to late seral species within this plant assemblage.

PASM and ELLA3 are by far the dominant grasses in MZ20 expression of this BpS. In MZ29, PASM, HECO26 and BOGR2 are by far the dominant grasses. Cool season grasses such as Indian ricegrass, bluebunch wheatgrass (Indian ricegrass and bluebunch wheatgrass occur only where coarser textured soils prevail), needle-and-thread (needle and thread has a broad environmental amplitude but is more typically abundant on coarse soils; however, under heavy grazing, it does quite well on fine-textured soils.), blue grama, Sandberg bluegrass, squirreltail, threadleaf sedge and infrequently Thurber's needlegrass. Rhizomatous wheatgrasses, such as western wheatgrass and thickspike wheatgrass, and plains reedgrass, are common species within these MZs 20 and 29. Junegrass also occurs.

Common forbs are species of *Astragalus*, *Crepis*, *Delphinium*, *Phlox* and *Castilleja*, while associated shrubs and shrub-like species can be small green rabbitbrush, MFRInge sagewort, winterfat and broom snakeweed. Other dominant species of forbs include RACO3 and SPCO. Also, LIPU and PHHO occur.

Forbs most important for MZ20 include SPHCOC, DALPUR, PHLHOO, RATCOL and OPUPOL. Other forbs in MZs 10 and 19 include hawksbeard (*Crepis acuminata*), bird's beak (*Cordylanthus* spp.), blue bell (*Mertensia* spp), Rocky Mountain aster (*Aster scopulorum*), *Phlox* species, lupine (*Lupinus* spp) and buckwheat (*Eriogonum* spp). In MZ29, all of the above are probably found except for lupine, which would occur in higher precipitation areas and be associated with mountain big sage.

Herbaceous species usually dominate the site prior to re-establishment. Site re-establishment is by seed

****Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.**

bank, seed production from remnant plants and seeds from adjacent (untreated) plants.

Wyoming big sagebrush in upland sites have fewer understory species relative to the mountain big sagebrush subspecies, though at higher elevations or moister areas of this vegetation community there is a higher potential for herbaceous species, relative to ARTTST (ssp tridentata) sites; no definitive statement on undergrowth herbaceous diversity can be made for ARTTSW (ssp wyomingensis) sites. Herbaceous cover increases transitioning into the mixedgrass prairie, and in open patches.

In MZ29, A.t. ssp. tridentata is not found. A.t. ssp. wyomingensis is found where A.t. ssp. vaseyana is not present. It can occur with greasewood and silver sagebrush, as well as rabbitbrush and saltbush.

Disturbance Description

Many researchers believe fire was the primary disturbance factor within this plant assemblage. Other disturbance factors may include insects, rodents and lagomorphs, drought, wet cycles, gradual changes in climate and native grazing (Wyoming Interagency Vegetation Community 2002). Drought may have been more significant disturbance than native grazing or insects, so was included. Native grazing by large ungulates (eg, bison), and insects were included as occurring every 10yrs but causing no transitions to another class. Heavy-impact grazing in the late closed stage occurs less frequently and causes a transition to an open state.

Following fire or other significant disturbance, herbaceous species will dominate the ecological site post-burning and recovery to prefire canopy cover is quite variable and may generally take 50-120yrs, but occasionally occurs within a decade (Baker, in press). Site re-establishment is by seed production from remnant plants, and seeds from adjacent (untreated) plants. Discontinuity of fuel in Wyoming big sagebrush communities can result in mosaic burn patterns, leaving remnant plants for seed, but can be large expanses of complete mortality (Bushey 1987, Baker, in press). Fire does not stimulate germination of soil-stored Wyoming big sagebrush, but neither does it inhibit its germination (Chaplin and Winward 1982). Regeneration may occur in pulses linked to high precipitation events (Maier et al. 2001).

Overall fire return intervals in Wyoming big sagebrush appear to have ranged from 100-240yrs or more (Baker, in press) for MZ22. In MZ20, some believe that intervals are shorter, with replacement fire occurring approximately every 30yrs in some of the classes (based on BLM Fire Management plans and local expert estimate, Downey). However, there was disagreement with that short interval. It is also said that we are fairly certain of the recovery time required (50-150yrs, mostly around 100yrs). With this slow recovery, if fires returned to the site in 30yrs, eventually the whole landscape would be only class A and maybe B (open) (Cooper, personal correspondence). Therefore, for MZ20, MFRI was modeled at an overall 90yr interval, similar to other adjacent mapzones and similar to BpS 1080 MFRI of 80yrs, which this BpS is thought to be very similar to.

There was some disagreement among MZ20 modelers as to the MFRI of 90yrs for this 1125 system. Up north, where there is a heavy grass component and much less cover of sagebrush than what is down south, and relatively connected topography and a lot of wind, it would burn more frequently (Downey, pers comm). Perhaps that would be considered BpS 1085 instead of BpS 1125. And even though BpS 1085, which is also comprised mainly of Wyoming big sagebrush has an MFRI of 30yrs, these two systems are different as it relates in large part to setting and precipitation patterns, and continuity of fuel. Eastern MT has few breaks, versus mountainous systems that would be much less likely to have the huge sweeping fires. Although the species are the same *Artemisia tridentata* ssp. *wyomingensis* - the systems aren't (Martin, pers comm). The longer MFRI for 1125 was therefore retained.

****Fire Regime Groups are:** I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Benkobi (pers comm) states that in MZ29, fire frequency could range from 36-40yrs (<http://gisdata.usgs.net>). However, MZ29 reviewers did not want to change the model. However, because it was also stated that recovery occurred after at least 60yrs in MZ29, and due to the discrepancy from previous mapzones, the MFRI from MZ20 was retained.

Discontinuity of fuel in Wyoming big sagebrush communities often result in mosaic burn patterns, but large expanses can burn with complete mortality under extreme conditions (Bushey 1987, Baker, in press). Mixed severity fire was originally modeled in this BpS but due to a new understanding of definitions of severity types, it was thought that mixed severity fire does not occur in this system and rather patchy fires do occur, with replacement severity.

MZs 20 and 29, where prescribed burn: after 29yrs, there was still zero recovery of Wyoming big sagebrush (Cooper pers comm). It is thought that the Wyoming big sagebrush communities take longer than 100yrs to recover. In Bighorn battlefield, historically there was much sagebrush. It burned in the mid-80s and there is still no evidence of sagebrush re-establishment 10yrs later.

Antelope, mule deer and pygmy rabbits are native herbivores that browse sagebrush. These were also not included in the model. In MZ29, probably not pygmy rabbits. Sage grouse might also have an impact? It is questionable as to the impact/frequency of antelope and mule deer in MZ29.

Adjacency or Identification Concerns

This type is difficult to distinguish from mixed-grass prairie with a high shrub component. It is possible that with severe disturbance, a state change might occur to mixed-grass prairie - which in turn changes the potential for the site to return to sagebrush. Extensive severe burns for want of an adjacent seedbank would take extensive periods before ARTTSW was again a significant component. The reference condition might have been sagebrush, but now the abiotic factors and biophysical gradients indicate a mixed-grass prairie.

Secondary shrub and herbaceous components may vary considerably across the range of its extent. Wyoming big sagebrush sites may be a mosaic with or abut juniper, ponderosa pine, salt desert shrub and grassland vegetation types across its range. However, the most common accompanying vegetation is Northern Great Plains midgrass prairie.

Broom snakeweed and halogeton may dominate sites disturbed by overgrazing, oil and gas development, or other disturbances. Club moss in this system increases with the intensity and duration of grazing. BROJAP can be an increaser with burning/grazing. There is also BROTEC invasion but that doesn't occur in the Northern Great Plains, except in MZ29.

Juniper increase might be occurring due to lack of fire today, but it is not developing into a true juniper woodland, especially in MZ29.

Shrub cover increases in MZ20 and 29 with overgrazing, and herbaceous layer decreases dramatically.

Might be difficult to distinguish from BpS 1080 and BpS 1085.

Much of 1080 has been lost due to land clearing for agriculture or converted to a cheatgrass or greasewood type. For basin big sagebrush in MZ29, this is the case. For Wyoming big sagebrush in MZ29, much has

****Fire Regime Groups are:** I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

been lost due to burning for modern grazing. The understory is currently more annual bromes due to increased grazing.

Overgrazing has also been an issue in 1080. HESCOM and KOEMAC increase (MZ20) where grazing is intense and protracted. It is questionable as to whether HESCOM increases with grazing (some areas of MZ29), and might rather decrease with overgrazing. With overgrazing in some areas of MZ29, more bluegrasses.

Plant associations are similar btwn 1125 and 1080. Shrubland is perhaps further south. Herbaceous cover is the only distinguishing factor. 1125 is definitely the more prominent historically. 1080 more prevalent in central WY. These (like mixegrass prairie) are distinguished by geography. Therefore, they're being combined for MZ29.

In Bighorns battlefield (around Hardin, MT), historic photos showed dense (up to 20%-30% cover, that is) shrub covered system, but currently, mostly grass - due to fires that burned there (Clark et al 1995 DRAFT).

If adjacent to pine systems, might be seeing more trees currently. (Also if in grass systems). This was seen in historic photographs throughout northern part of MZ29 and through western SD (Clark et al 1995 DRAFT).

Native Uncharacteristic Conditions

Over 45% shrub cover would be uncharacteristic for MZ20 and MZ29. In fact, Wyoming big sagerush in MZ29 would not exceed 40% cover. The only reason it would be this high is in cases of extreme overgrazing or in the absence of fire or changes in fire regime - frequency.

Scale Description

Occurrences may cover between hundreds and thousands of hectares.

Disturbance patch sizes range from 10s-1000s of hectares. The patch and disturbance size gets larger as this shrub BpS intergrades with the grassland BpS, and also gets larger from MZs 19 and 20 into MZ29.

Issues/Problems

Difficult to identify where hybrids occur with other big sagebrush taxa.

Comments

This model for MZ29 was adapted from the same BpS from MZ20 created by Steve Cooper and Shannon Downey and reviewed by Steve Barrett. For MZs 29 and 30, descriptive additions and changes were made. Other reviewers for MZ29 were Bobby Baker and Jim Von Loh.

Model for MZ20 was adapted from the draft model for MZ22 for 1125b Inter-Mountain Basins Big Sagebrush Steppe-Wyoming Big Sagebrush, created by Mark Williams, Vicki Herren and an anonymous contributor and reviewed by Tim Kramer, Eve Warren and Destin Harrell. Changes were made to the description and model.

The model for MZ22 was adapted from Rapid Assessment (RA) model R0SBWYwy created by Tim Kramer (tim_kramer@blm.gov) and reviewed by Bill Baker, Don Bedunah and Dennis Knight.

****Fire Regime Groups are:** I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

For the Rapid Assessment, the workshop code was WYSB. This model was combined with another Rapid Assessment model, ROSBWA (workshop code was WSAG1), modeled by George Soehn (george_soehn@blm.gov) and reviewed by Sarah Heide (sarah_heide@blm.gov) and Krista Gollinick-Waid (krista_waid@blm.gov). The two were combined based on peer-review and the similarity of disturbance regimes and species composition.

The RA Model is based on the original FRCC PNVG (WYSB1) with modifications from Wyoming Interagency Vegetation Committee (2002) and expert estimates. Peer review for the RA model was incorporated 4/30/2005. Additional reviewers were Karen Clause (karen.clause@wy.usda.gov), Ken Stinson (ken_stinson@blm.gov) and Eve Warren (eve_warren@blm.gov).

Vegetation Classes

Class A 35 %		<u>Indicator Species and Canopy Position</u>	<u>Structure Data (for upper layer lifeform)</u>									
Early Development 1 All Structure		NAVI4	<table><tr><td></td><td>Min</td><td>Max</td></tr><tr><td>Cover</td><td>0 %</td><td>80 %</td></tr><tr><td>Height</td><td>Herb 0m</td><td>Herb 0.5m</td></tr></table>		Min	Max	Cover	0 %	80 %	Height	Herb 0m	Herb 0.5m
	Min	Max										
Cover	0 %	80 %										
Height	Herb 0m	Herb 0.5m										
<u>Upper Layer Lifeform</u>		Upper	<input checked="" type="checkbox"/> Upper layer lifeform differs from dominant lifeform. Herbs dominate this class, but shrubs are growing up and do not yet dominate the class. Shrub cover less than five percent belongs in this class.									
<input checked="" type="checkbox"/> Herbaceous		PASM										
<input type="checkbox"/> Shrub		Upper										
<input type="checkbox"/> Tree		BOGR2										
<u>Fuel Model</u>		Lower										
2		CAFI										
<u>Description</u>		Lower										

Herbaceous dominated. In the pre-settlement condition, NAVI4 (in MZ20) and HECO26 in MZ29 would have been a major upper position component. Primarily grasses with forbs. Exact species will vary depending on location. Western wheatgrass, Sandberg bluegrass, plains reedgrass, needle-and-thread, bluebunch wheatgrass, threadleaf sedge, plains junegrass and blue grama would be dominant grasses. Forbs may include Astragalus, Crepis, Castelleja, Delphinium, Agoseris, Phlox and others. There may also be significant component of small green rabbitbrush.

Succession to class B, a mid-development open stage, occurs after 40yrs. This succession was originally modeled at 20yrs; however, it was later decided that that was a minimum age for succession, and it would take more like 40yrs to achieve 5-15% canopy cover of ARTTSW. There is one paper that shows no ARTTSW 15yrs post-fire and another paper for MZ19 that indicates no recovery after as much as 18yrs (Cooper, personal correspondence). In MZ29, recovery occurred after 60yrs.

Insect/disease (0.001 probability or 0.1% of the landscape each year), native grazing (0.1 probability or 10% of the landscape each year) and wind/weather stress (every 100yrs, 0.01 probability or 1% of the landscape each year) occur, but do not cause a transition.

Replacement fire was originally modeled at every 30yrs, based on expert estimate and local observations. - in BLM Fire Management Plans (Downey, personal correspondence). However, this was later changed to 90yrs based on recovery times of this type. This, and the other changes in age range, changed the class percentage from 20% to 35%.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Class B 40 %

Mid Development 1 Open

Upper Layer Lifeform☐ Herbaceous☒ Shrub☐ Tree**Fuel Model**

2

Indicator Species and Canopy Position

ARTRW8

Upper

PASM

Mid-Upper

NAVI4

Mid-Upper

HECO26

Middle

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	20 %
Height	Shrub 0m	Shrub 0.5m
Tree Size Class		

☐ Upper layer lifeform differs from dominant lifeform.**Description**

Sagebrush canopy is greater than five but less than 15%. Understory is well represented by herbaceous species as described for class A. (Montana Academy of Sciences publication - re: in breaks, After 15yrs after fire, no sagebrush yet.)

ARFR4 also present in lower canopy.

Succession to class C, late development closed stage, occurs after 50yrs. (60yrs for MZ29)

Insect/disease (0.001 probability of 0.1% of the landscape each year), native grazing (0.1 probability or 10% of the landscape each year) and wind/weather stress (every 100yrs, 0.01 probability or 1% of the landscape each year) occur, but do not cause a transition to another stage.

Fire was modeled more frequently than in MZ22 based on expert estimate and data from BLM Fire Mangement Plans. Originally, mixed fire was modeled at occurring every 40yrs, maintaining the class in this stage (Downey, personal correspondence). However, this was later removed due to a new understanding of definitions of mixed versus replacement fire. This, and the other changes in age range, changed the class percentage from 55% to 35%. Replacement fire occurs every 90yrs.

Class C 25 %

Late Development 1 Open

Upper Layer Lifeform☐ Herbaceous☒ Shrub☐ Tree**Fuel Model**

2

Indicator Species and Canopy Position

ARTRW8

Upper

PASM

Mid-Upper

NAVI4

Mid-Upper

HECO26

Middle

Structure Data (for upper layer lifeform)

	Min	Max
Cover	21 %	40 %
Height	Shrub 0m	Shrub 0.5m
Tree Size Class		

☐ Upper layer lifeform differs from dominant lifeform.**Description**

Sagebrush canopy is >15 percent. Understory is well represented by herbaceous species as described for class A. This class is more common on drier sites.

Shrub cover max was 30% in MZ20. In MZ29, it was increased to 65% cover by other reviewers. However, it was decided that here it could not be this amount of cover. Modal cover is 15%. The most measured was 32% cover. Some could have been higher cover but not much. Common in literature that grazing/over-grazing increases cover, not the opposite.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

It is probably more common in 20% range. 40% is high, but could be a max (Cooper, diBenedetto, personal comm). Regional lead changed to 40% per comments.

ARFR4 is also present in lower canopy.

Insect/disease (0.001 probability of 0.1% of the landscape each year), native grazing (0.002 probability or 0.2% of the landscape each year) cause a transition to the mid-open stage.

Native grazing (0.1 probability or 10% of the landscape each year) occurs, but does not cause a transition to another stage.

Drought was modeled at an overall interval of 100yrs split between maintaining this stage or taking it to the mid-development stage.

Originally, mixed fire was modeled at occurring every 40yrs, maintaining the class in this stage (Downey, personal correspondence). However, this was later removed due to a new understanding of definitions of mixed versus replacement fire. Replacement fire occurs every 100yrs. This only changed the class percentage from 25% to 30%.

Class D 0 % [Not Used] [Not Used] Upper Layer Lifeform <input type="checkbox"/> Herbaceous <input type="checkbox"/> Shrub <input type="checkbox"/> Tree	<u>Indicator Species and Canopy Position</u> <u>Fuel Model</u>	Structure Data (for upper layer lifeform) <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th style="text-align: center;"><i>Min</i></th> <th style="text-align: center;"><i>Max</i></th> </tr> </thead> <tbody> <tr> <td><i>Cover</i></td> <td style="text-align: center;">%</td> <td style="text-align: center;">%</td> </tr> <tr> <td><i>Height</i></td> <td></td> <td></td> </tr> <tr> <td><i>Tree Size Class</i></td> <td colspan="2"></td> </tr> </tbody> </table> <input type="checkbox"/> Upper layer lifeform differs from dominant lifeform.		<i>Min</i>	<i>Max</i>	<i>Cover</i>	%	%	<i>Height</i>			<i>Tree Size Class</i>		
	<i>Min</i>	<i>Max</i>												
<i>Cover</i>	%	%												
<i>Height</i>														
<i>Tree Size Class</i>														

Description

Class E 0 % [Not Used] [Not Used] Upper Layer Lifeform <input type="checkbox"/> Herbaceous <input type="checkbox"/> Shrub <input type="checkbox"/> Tree	<u>Indicator Species and Canopy Position</u> <u>Fuel Model</u>	Structure Data (for upper layer lifeform) <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th style="text-align: center;"><i>Min</i></th> <th style="text-align: center;"><i>Max</i></th> </tr> </thead> <tbody> <tr> <td><i>Cover</i></td> <td style="text-align: center;">%</td> <td style="text-align: center;">%</td> </tr> <tr> <td><i>Height</i></td> <td></td> <td></td> </tr> <tr> <td><i>Tree Size Class</i></td> <td colspan="2"></td> </tr> </tbody> </table> <input type="checkbox"/> Upper layer lifeform differs from dominant lifeform.		<i>Min</i>	<i>Max</i>	<i>Cover</i>	%	%	<i>Height</i>			<i>Tree Size Class</i>		
	<i>Min</i>	<i>Max</i>												
<i>Cover</i>	%	%												
<i>Height</i>														
<i>Tree Size Class</i>														

Description

Disturbances

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Fire Regime Group:** IV

Historical Fire Size (acres)

Avg

Min

Max

Sources of Fire Regime Data

- ☒ Literature
- ☒ Local Data
- ☒ Expert Estimate

Additional Disturbances Modeled

- ☒ Insects/Disease
- ☒ Native Grazing
- ☐ Other (optional 1)
- ☒ Wind/Weather/Stress
- ☐ Competition
- ☐ Other (optional 2)

Fire Intervals

	Avg FI	Min FI	Max FI	Probability	Percent of All Fires
Replacement	90			0.01111	100
Mixed					
Surface					
All Fires	90			0.01113	

Fire Intervals (FI):

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class.

References

Anderson, J.E. and R.S. Inouye. 2001. Landscape-scale changes in plant species abundance and biodiversity of a sagebrush steppe over 45 years. *Ecological Monographs*. 71(4): 531-556.

Baker, W.L. In press. Fire and restoration of sagebrush ecosystems. *Wildlife Society Bulletin*, in press.

Benkobi, L. and D.W. Uresk. 1996. Seral stage classification and monitoring model for big sagebrush/western wheatgrass/blue grama habitat. In: J.R. Barrow, E.D. McArthur, R.E. Sosebee and R.J. Tausch, compilers. *Proceedings: shrubland ecosystem dynamics in a changing environment; 1996 May 23-25; Las Cruces, NM*. Gen. Tech. Rep. INT-GTR-338. Ogden, UT: USDA Forest Service, Intermountain Research Station.

Bunting, S.C., B.M Kilgore and C.L. Bushey. 1987. Guidelines for prescribed burning sagebrush-grass rangelands in the northern Great Basin. Gen. Tech. Rep. INT-231. Ogden, UT: USDA Forest Service. 33 pp.

Clark, R., . DiBenedetto, J. Losensky. 1995 DRAFT. A description of historic vegetation patterns and trends on the Norther Plains using repeat photography. USDA Forest Service.

Fire Regime Condition Class (FRCC) Interagency Handbook Reference Conditions, Modeler: Doug Havlina, Date: 8/15/03, PNVG Code: WSAG1. 2.

Knight, D.H. 1994. *Mountains and Plains, The Ecology of Wyoming Landscapes*. Yale University Press, New Haven, CT.

Lesica, P., S.V. Cooper and G. Kudray. 2005. Big sagebrush shrub-steppe postfire succession in southwest Montana. Unpublished report to Bureau of Land Management, Dillon Field Office, Montana Natural Heritage Program, Helena, MT. 29 pp. plus appendices.

NatureServe. 2007. *International Ecological Classification Standard: Terrestrial Ecological Classifications*. NatureServe Central Databases. Arlington, VA. Data current as of 10 February 2007.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

- Perryman, L., A.M. Maier, A.L., Hild and R.A. Olson. 2001. Demographic characteristics of three *Artemisia tridentata* Nutt. subspecies. *Journal of Range Management*. 54(2): 166-170.
- Sturges, D.L. 1994. High-elevation watershed response to sagebrush control in southcentral Wyoming. Res. Pap. RM-318. Fort Collins, CO: USDA Forest Service, Rocky Mountain Forest and Range Experiment Station. 19 pp.
- Vale, T.R. 1975. Presettlement vegetation in the sagebrush-grass area of the Intermountain West. *Journal of Range Management*. 28(1): 32-36.
- Welch, B.L and C. Criddle. 2003. Countering Misinformation Concerning Big Sagebrush. Research Paper RMRS-RP-40. Ogden, UT: USDA Forest Service, Rocky Mountain. Research Station. 28 pp.
- Winward, A.H. 1991. A renewed commitment to management of sagebrush grasslands. In: Research in rangeland management. Ag. Exper. Stn. Special Rep. 880. Corvallis, OR: Oregon State University. 7 pp.
- Wyoming Interagency Vegetation Committee. 2002. Wyoming Guidelines for Managing Sagebrush Communities with Emphasis on Fire Management. Wyoming Game and Fish Department and Wyoming BLM. Cheyenne, WY. 53 pp.
- Young, J.A. and R.A. Evans. 1978. Population dynamics after wildfires in sagebrush grasslands. *Journal of Range Management*. 31: 283-289.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.