

Rapid Assessment Reference Condition Model

The Rapid Assessment is a component of the LANDFIRE project. Reference condition models for the Rapid Assessment were created through a series of expert workshops and a peer-review process in 2004 and 2005. For more information, please visit www.landfire.gov. Please direct questions to helpdesk@landfire.gov.

Potential Natural Vegetation Group (PNVG)

R5BSSA Bluestem - Saccahuista

General Information

Contributors (additional contributors may be listed under "Model Evolution and Comments")

Modelers

Ron Masters rmasters@ttrs.org
Susanne Hickey shickey@tnc.org
Judy Teague Judy_Teague@natureserve.org

Reviewers

Doug Zollner dzollner@tnc.org

Vegetation Type

Grassland

General Model Sources

- Literature
 Local Data
 Expert Estimate

Rapid Assessment Model Zones

- | | |
|--|---|
| <input type="checkbox"/> California | <input type="checkbox"/> Pacific Northwest |
| <input type="checkbox"/> Great Basin | <input checked="" type="checkbox"/> South Central |
| <input type="checkbox"/> Great Lakes | <input type="checkbox"/> Southeast |
| <input type="checkbox"/> Northeast | <input type="checkbox"/> S. Appalachians |
| <input type="checkbox"/> Northern Plains | <input type="checkbox"/> Southwest |
| <input type="checkbox"/> N-Cent. Rockies | |

Dominant Species*

SCHIZ QUVI
SCSCL ANTE
PANIC ANVI2
SPSP PAPL3

LANDFIRE Mapping Zones

36
37

Geographic Range

This PNVG is located along the Gulf Coast and inland varying distances from 50 to 150 miles (80-240 km) from south Texas to Louisiana and the mouth of the Mississippi River. To the north this type is bordered by Oak-Hickory forest (Kuchler type 100) in much of Texas and in east Texas Oak-Hickory-Pine (Kuchler type 111). In Louisiana, it is bordered to the north and east by Southern Floodplain Forest (Kuchler type 112) (Kuchler 1964). To the south and west it also joins with the desert grasslands.

Biophysical Site Description

The bluestem-sacahuista is relatively flat, but is characterized by ridge-swale or mound-intermound microtopography. This type is dissected by numerous rivers and streams which result in highly variable species composition (Johnston 1963, Diamond and Smeins 1985, Drawe 1994). A topographic and moisture gradient exists as one progresses inland and out of floodplains. The diversity of embedded edaphic conditions and wetlands within the general type is important and interacted with fire to determine wildlife species distributions. Extended inundation in areas referred to as lagunas adds a disturbance element within 25 km of the coast. These areas are subject to a different successional pattern than that following other types of disturbance (Scifres and Mutz 1975).

Vegetation Description

This type has many of the same vegetation elements of tallgrass prairie but also has a number of additional species, including some tropical grasses. Nearly 1,000 plant species have been identified in this type. The forb community tends to be richer in the coastal prairie than in true tallgrass prairie. This type is considered a shrub-grassland complex rather than a prairie (Johnston 1963, Scifres and Mutz 1975, Drawe 1994). This type is highly variable in species composition because of the dissected nature of the terrain and topography caused by numerous rivers and creeks (Johnston 1963, Diamond and Smeins 1985, Drawe 1994). The species composition is dominated by little bluestem (*Schizachyrium scoparium*), sea coast bluestem (*S. c. var. littoralis*) several Panicums and sacahuista, also known as Gulf cordgrass (*Spartina spartinae*). Sacahuista primarily dominates along floodplains of the numerous rivers and near the coast.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Other important species include bushy bluestem (*Andropogon glomeratus*), other bluestems such as split-beard (*A. ternarius*), broomsedge bluestem (*A. virginicus*), silver bluestem (*Bothriochloa saccharoides*), various *Sporobolus*, and several tropical grasses of the genera *Heteropogon*, *Paspalum*, *Trachypogon* and the previously mentioned *Panicum*. Secondary species vary in importance regionally depending on topography and soil moisture relations and include sideoats grama (*Bouteloua curtipendula*), buffalo grass (*Buchloe dactyloides*) and threeawns (*Aristida* spp.). Several grass-like species that are important include *Carex* spp., *Eleocharis* spp., *Scirpus* spp. Conspicuous forbs include the genera *Ratibida*, *Rudbeckia*, *Liatris*, and *Sagittaria*. Shrubs that are important include honey mesquite (*Prosopis glandulosa*) and various acacias most notably huicache (*Acacia farnesiana*) in Texas, also *Rosa bracteata*, and various oaks (*Quercus* spp). Eastern baccharis (*Baccharis halimifolia*) and wax myrtle (*Myrica cerifera*) are more important to the east. All of these woody plants and others increase in the absence of fire.

Disturbance Description

This type is fire regime group II, with frequent replacement fires, both lightning and anthropogenic in origin (Stewart 1951, Lehmann 1965, Drawe 1980, Stewart 2002; Journey et al 2004). Likely, this type has one of the most frequent fire regimes in North America. Annual burning was described in references to historic accounts (Stewart 1951, Chamrad and Dodd 1973, Stewart 2002:141-144) and in one instance reference was made to burning twice (summer and winter) in the same year (Lehmann 1965:133). These references do not indicate every acre was burned every year but likely some considerable area was burned every year with most of the type being burned at least biannually and some areas burned twice in a given year. Lehmann (1965) also notes accounts about the patches of unburned vegetation and relative green-up compared to burned areas. Fire was likely possible during most seasons and dependant on the availability of dry fine fuels sufficient to carry a fire. Historic accounts from the 1800's depict large burns, but the terrain is dissected by numerous rivers and creeks bordered by trees (Lehmann 1965, Drawe 1994). Therefore this landscape matrix strongly influenced the probable size of burn. A problem with much of the literature on fire in prairies, and therefore a caution, is that it does not include interaction with herbivory (Engle and Bidwell 2001). Bison (*Bison bison*) were historically an important source of disturbance that increased heterogeneity of patches on the landscape. Wild horses were established early on and large herds were noted by early explorers in the southern part of this type (Stewart 2002). Pronghorn antelope historically occurred in the southwestern most part of this type (Nelson 1925) where rainfall amounts dropped considerably. Although historical accounts of large groups (1,000's) of bison do occur, bison herds were of smaller size and more dispersed in this system than herds of the central Great Plains. Bison grazing affects fire patterns and thus the landscape patterns in tallgrass prairie (Risser 1990) and assuredly this system as well. Bison and other grazing/browsing wildlife species preferentially seek out the new growth of recently burned areas affecting patch composition (e.g., Coppedge and Shaw 1998, Jackson 1965, Risser 1990, Steuter 1986, Fuhlendorf and Engle 2004). Burn accounts are in agreement with the patch burn model where small burns are preferentially grazed by bison. Using the fire/bison interaction model first proposed by Steuter (1986) recent modifications propose that anywhere from 1/6 to 1/3 of a 20,000 acre (8,094 hectares) tallgrass landscape likely burned (Fuhlendorf and Engle 2004). Likely this figure is less for coastal prairie because of the dissected terrain. Burning causes earlier green-up and increased nutrient content of native grasses and is preferentially selected by grazing animals (Lehmann 1965, Oefinger and Scifres 1977). Typically following green-up, fire is followed by intensive bison grazing pressure to the point that structural classes shifted over the landscape in response to an interaction between bison grazing pressure and fire (Steuter 1986; Fuhlendorf and Engle 2001, 2004). Heavily grazed and trampled areas would not burn in the next year to three years creating a one-way closed path. Following this type disturbance the patches are dominated with forbs and will not burn in the succeeding dormant and growing season because of lack of fuel. Whereas previous years unburned post-grazing re-growth would be the next patch to burn. Bison grazing influenced fire return intervals. Fire occurrence in turn influenced bison grazing distribution. This model depicts a landscape composed of a continuously shifting mosaic of patches with a short time period of duration. The small patch burn and very frequent fire scenario is essential to perpetuate suitable lek sites and brood rearing habitat for Attwater's prairie chicken (*Tympanicus cupido attwateri*) in this system with long growing seasons, fertile soil and quick recovery time and with habitat requirements (Kessler 1978) similar to other

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

prairie chicken species noted by Sparks and Masters (1996). This species historically occurred as somewhat discrete populations in parts of the blackland prairie and coastal prairie (Lehmann 1965, Chamrad and Dodd 1973, Silvy and Hagen 2004, Silvy et al 2004). Frequent fire is essential to control woody dynamics in this dissected landscape mosaic of rivers and creeks with stringers of bottomland and some upland forests (Denevan 1992; Lehmann 1965, Stewart 1951, 2002) and varying edaphic and moisture conditions (Scifres and Mutz 1975).

Adjacency or Identification Concerns

Scale Description

Sources of Scale Data Literature Local Data Expert Estimate

We (Lee, Judy & Susanne) reviewed maps showing the large rivers to help derive average fire size which would be limited by breaks in fuel (i.e. rivers).

Issues/Problems

Model Evolution and Comments

Lee Elliott (TNC) also assisted with the model development. We combined the live oak savanna, saline prairie communities with this PNVG. In the model we used alternative succession to account for the small percentage of the landscape that would contain live oak savanna. For class C, grazing in combination with drought (wind/weather/stress) moves a small percentage of Class C to class E, through reduced competition between oak and grass. Drought in class E can reduce a small percentage of the oaks thus pushing those areas to class B.

Succession Classes**
Succession classes are the equivalent of "Vegetation Fuel Classes" as defined in the Interagency FRCC Guidebook (www.frcc.gov).

Class A 30 %

Early1 All Struct

Description

Post fire community that is short duration (often weeks-depending on time of burning) before transitioning into one of the other community stages. Succession post inundation with water proceeds in a different manner through a sedge then bunchgrass stage.

Dominant Species* and Canopy Position

SCHIZ4 Upper
 SCSCCL Upper
 PANIC Upper
 SPSP Upper

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 1

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	55 %
Height	no data	Herb Medium 0.5-0.9m
Tree Size Class	no data	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class B 45 %

Mid2 Closed

Description

Mixed forb and grass community either somewhat recovered from bison grazing, or inundation with water or continuing post burn development. Can be somewhat forb dominated with a woody component in areas.

Dominant Species* and Canopy Position

SCHIZ4 Upper
 SCSCCL Upper
 PANIC Upper
 SPSP Upper

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 3

Structure Data (for upper layer lifeform)

	Min	Max
Cover	55 %	100 %
Height	Herb Medium 0.5-0.9m	Herb Tall > 1m
Tree Size Class	no data	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Class C 20%

Mid1 Open
Description

Forb dominated site with sparse bunchgrass clumps, derived from heavy bison grazing and trampling pressure, wallowing and horning, or inundation with water.

Dominant Species* and Canopy Position

RATIB Upper
RUDBE Upper
SCHIZ4 Upper
SCSCL Upper

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 1

Structure Data (for upper layer lifeform)

	Min	Max
Cover	30 %	55 %
Height	Herb Short <0.5m	Herb Medium 0.5-0.9m
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class D 2%

Late1 Closed
Description

Tallgrass dominated but with a persistent woody component, tillering and overall plant vigor reduced by mulching effect from accumulation of ungrazed, unburned plant litter. Over short periods of fire exclusion woody encroachment will rapidly occur. The woody element will also increase following drought and over-utilization of herbaceous plants. Can go from tree seedling/sapling to large trees.

Dominant Species* and Canopy Position

SCHIZ4 Middle
SCSCL Middle
PRGL2 Upper
ACFA Upper

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 3

Structure Data (for upper layer lifeform)

	Min	Max
Cover	55 %	100 %
Height	Herb Medium 0.5-0.9m	Tree Short 5-9m
Tree Size Class	Medium 9-21"DBH	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

This class is composed of tall grass species with thatch buildup and reduced tillering and plant vigor, but the class also has a persistent woody component which initially in this class would be shrub size but quickly growing to small tree sized.

Class E 3%

Late1 Open
Description

Oak mottes/live oak savanna found in areas of sand. Species of live oak changes depending on location in PNVG, Quercus virginiana, north of Brazos River and Q. fusiformis, south of Brazos River. Vegetation can range from tree seedling/sapling to large trees.

Dominant Species* and Canopy Position

QUVI Upper
SCSCL Lower
QUFU Upper
ILVO Middle

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 9

Structure Data (for upper layer lifeform)

	Min	Max
Cover	10 %	40 %
Height	Tree Regen <5m	Tree Medium 10-24m
Tree Size Class	Large 21-33"DBH	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Disturbances

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Disturbances Modeled

- Fire
- Insects/Disease
- Wind/Weather/Stress
- Native Grazing
- Competition
- Other:
- Other

Historical Fire Size (acres)

Avg: 100000
 Min: 100
 Max: 300000

Sources of Fire Regime Data

- Literature
- Local Data
- Expert Estimate

Fire Regime Group: 2

- I: 0-35 year frequency, low and mixed severity
- II: 0-35 year frequency, replacement severity
- III: 35-200 year frequency, low and mixed severity
- IV: 35-200 year frequency, replacement severity
- V: 200+ year frequency, replacement severity

Fire Intervals (FI)

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class. All values are estimates and not precise.

	<i>Avg FI</i>	<i>Min FI</i>	<i>Max FI</i>	<i>Probability</i>	<i>Percent of All Fires</i>
<i>Replacement</i>	3.6	1	2	0.27778	68
<i>Mixed</i>	7.7	2	5	0.12987	32
<i>Surface</i>					
<i>All Fires</i>	2			0.40766	

References

Anderson, R. C. and L. E. Brown. 1986. Stability and instability in plant communities following fire. American Journal of Botany 73:364-368.

Axelrod, D. I. 1985. Rise of the grassland biome, central North America. Botanical Review 51:163-201.

Bailey, D.W., J.E. Gross, A.L. Emilio, L.R. Rittenhouse, M.B. Coughenour, D.M. Swift, and P.L. Sims. 1996. Mechanisms that result in large herbivore grazing distribution patterns. Journal of Range Management 49(5):386-400.

Biondini, M.E., A.A. Steuter, and R.G. Hamilton. 1999. Bison use of fire-managed remnant prairies. Journal of Range Management 52(5):454-461.

Box, T. W., and R. S. White. 1969. Fall and winter burning of south Texas brush ranges. Journal of Range Management 22:373-376.

Brown, James K.; Smith, Jane Kapler, eds. 2000. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 257 p.

Chamrad, A. D. and J. D. Dodd. 1973. Fire in the range of Attwater’s prairie chicken. Proceedings Tall Timbers Fire Ecology Conference 12:257-276.

Coppedge, B.R., and J.H. Shaw. 1997. Effects of horning and rubbing behavior by bison (Bison bison) on woody vegetation in a tallgrass prairie landscape. American Midland Naturalist 138(1):189-196.

Coppedge, B.R., and J.H. Shaw. 1998. Bison grazing patterns on seasonally burned tallgrass prairie. Journal of Range Management 51(3):258-264.

Coppedge, B.R., and J.H. Shaw. 1998. Diets of bison social groups on Tallgrass Prairie in Oklahoma. The Prairie Naturalist 30(1):29-36.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

- Coppedge, B.R., and J.H. Shaw. 2000. American bison Bison bison wallowing behavior and wallow formation on tallgrass prairie. *Acta Theriologica* 45(1):103-110.
- Daubenmire, R. 1968. Ecology of fire in grasslands. *Adv. Ecol. & Research* 5:209-266.
- Denevan, W. M. 1992. The pristine myth: the landscape of the Americas in 1492. *Annals of the Association of American Geographers* 82:369-385.
- Diamond, D. D., and F. E. Smeins. 1985. Composition, classification and species response patterns of remnant tallgrass prairies in Texas. *The American Midland Naturalist* 113(2):294-308.
- Diamond, D. D., D. H. Risking and S. L. Orzell. 1987. A framework for plant community classification and conservation in Texas. *The Texas Journal of Science* 39:203-221.
- Drawe, D. L. 1980. The role of fire in coastal prairie. Pages 101-113 in *Proceedings of a Symposium: Prescribed Range Burning in the coastal prairie and eastern Rio Grande Plains of Texas*. Texas Agricultural Extension Service, The Texas A & M University System, College Station, TX.
- Drawe, D. L. 1994. Bluestem-sacahuista Prairie SRM 711. Pages 91-92 in T. N. Shiflet, ed. *Rangeland cover types of the United States*. Society for Range Management, Denver, CO. 152p.
- Durham, A. J., and M. M. Kothmann. 1977. Forage availability and cattle diets on the Texas coastal prairie. *Journal of Range Management* 30:103-106.
- Engle, D.M., and T.G. Bidwell. 2001. Viewpoint: The response of Central North American prairies to seasonal fire. *Journal of Range Management* 54:2-10.
- Ford, M. F., and J. B. Grace. 1998. The interactive effects of vertebrate herbivory and fire on a coastal marsh in Louisiana, the Pearl River. *Wetlands* 18:1-8.
- Ford, M. F., and J. B. Grace. 1998. Effects of herbivores on vertical soil accretion, shallow subsidence and soil elevation changes in coastal Louisiana. *Journal of Ecology* 86:974-982.
- Fuhlendorf, S. D. and D. M. Engle. 2001. Restoring heterogeneity on rangelands: ecosystem management based on evolutionary grazing patterns. *BioScience* 51:625-632.
- Fuhlendorf, S. D. and D. M. Engle. 2004. Application of the fire-grazing interaction to restore a shifting mosaic on tallgrass prairie. *Journal of Applied Ecology* 41:604-614.
- Gough, L. and J. B. Grace. 1998. Effects of flooding, salinity, and herbivory on coastal plant communities, Louisiana, United States. *Oecologia* 117:527-535.
- Gough, L. and J. B. Grace. 1999. Predicting effects of environmental change on plant species density: experimental evaluations in a coastal wetland. *Ecology* 80:882-890.
- Grace, J. B. and B. H. Pugsek. 1997. A structural equation model of plant species richness and its application to a coastal wetland. *American Nat.* 149:436-460.
- Grace, J. B. and G. R. Guntenspergen. 1999. The effects of landscape position on plant species density: evidence of past environmental effects in a coastal wetland. *Ecoscience* 6:381-391.

Grace, J. B. and H. Jutila. 1999. The relationship between species density and community biomass in grazed and ungrazed coastal meadows. *Oikos* 85:398-408.

Grace, J. B., L. Allain, and C. Allen. 2000. Factors associated with plant species richness in a coastal tallgrass prairie. *Journal of Vegetation Science* 11:443-452.

Grace, J. B., L. Allain, and C. Allen. 2000. Vegetation associations in a rare community type -coastal tallgrass prairie. *Plant Ecology* 147:105-115.

Hansmire, J. A., D. L. Drawe, D. B. Wester and C. M. Britton. 1988. Effect of winter burns on forbs and grasses of the Texas coastal prairie. *The Southwest Naturalist* 33:333-338.

Jackson, A. S. 1965. Wildfires in the Great Plains grasslands. *Proc. Tall Timbers Fire Ecology Conference* 4:241-259.

Johnston, M. C. 1963. Past and present grasslands of southern Texas and northeastern Mexico. *Ecology* 44:456-466.

Jutila, H. M., and J. B. Grace. 2002. Effects of disturbance on germination and seedling establishment in a coastal prairie grassland: a test of the competitive release hypothesis. *Journal of Ecology* 90:291-302.

Jurney, D., R. Evans, J. Ippolito, and V. Bergstrom. 2004. The role of wildland fire in portions of southeastern North America. Pages 95-116 in R. T. Engstrom and W. J. de Groot (eds). 22nd Tall Timbers Fire Ecology Conf. Proceedings. Kanaskas, Alberta.

Kessler, W. B. 1978. Attwater prairie chicken ecology in relation to agricultural practices. Ph.D. Thesis, Texas A&M University, College Station. 158p.

Komarek, E. V. 1965. Fire ecology, grasslands and man

Kucera, C. L. 1978. Grasslands and fire. Pages 90-111 in *Proc. Conference Fire regimes and ecosystem properties*. Gen. Tech. Rep. WO-26. USDA For. Ser., Washington, DC.

Kuchler, A. W. 1964. Potential natural vegetation of the conterminous United States. Special Publication 36. American Geographical Society, New York, New York.

Lehmann, V. W. 1965. Fire in the range of Attwater's prairie chicken. *Proceedings Tall Timbers Fire Ecology Conference* 4:127-143.

Lonard, R. I., F. W. Judd, and E. H. Smith. 2003. Recovery of vegetation following a wild fire on the margins of tidal flats, Padre Island National Seashore, Texas. *Texas Journal of Science* 55:347-364.

Lynch, B. D. 1962. Study of a grassland mosaic at Austin, Texas. *Ecology* 43:679-686.

McAtee, J. W., C. J. Scifres, and D. L. Drawe. 1979. Improvement of Gulf cordgrass range with burning or shredding. *Journal of Range Management* 32:372-375.

Moore, C. T. 1972. Man and fire in the central North American grassland 1535-1890: a documentary historical geography. Ph.D. Dissertation, University of California, Los Angeles. 155p.

Nelson, E. W. 1925. Status of the pronghorn antelope, 1922-1924. U.S. Department of Agriculture Bulletin

No. 1346. Washington, D.C. 64p.

Oefinger, R. D. and C. J. Scifres. 1977. Gulf cordgrass production, utilization and nutritional value following burning. Texas A&M University, Texas Agricultural Experiment Station Bulletin. College Station, TX. 19p.

Plumb, G.E., and J.L. Dodd. 1993. Foraging ecology of bison and cattle on a mixed prairie: implications for natural area management. *Ecological Applications* 3(4):631-643.

Risser, P. G. 1990. Landscape processes and the vegetation of the North American grassland. in S. L. Collins and L. L. Wallace, eds. *Fire in North American tallgrass prairies*. University of Oklahoma Press, Norman. 175p.

Sampson, F. B., F. L. Knopf and W. R. Ostlie. 2004. Great Plains ecosystems; past, present and future. *Wildlife Society Bulletin* 32:6-15.

Sauer, C. O. 1950. Grassland climax, fire and man. *Journal of Range Management* 3:16-21.

Schmidt, Kirsten M, Menakis, James P., Hardy, Colin C., Hann, Wendel J., Bunnell, David L. 2002. Development of coarse-scale spatial data for wildland fire and fuel management. Gen. Tech. Rep. RMRS-GTR-87. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 41 p. + CD.

Scifres, C. J., and K. W. Duncan. 1982. Brownseed paspalum response to season of burning. *Journal of Range Management* 35:251-253.

Scifres, C. J., and J. L. Mutz. 1975. Secondary succession following extended inundation of Texas Coastal Rangeland. *Journal of Range Management* 28:279-282.

Scifres, C. J., J. W. McAtee, and D. L. Drawe. 1980. Botanical, edaphic and water relationships of gulf cordgrass (*Spartina spartinae*) and associated communities. *The Southwestern Naturalist* 25:397-410.

Scifres, C. J., J. L. Munz, R. E. Whitson, and D. L. Drawe. 1982. Interrelationships of huisache canopy cover with range forage on the coastal prairie. *Journal of Range Management* 35:558-562.

Shaw, J.H., and M. Lee. 1997. Relative abundance of bison, elk, and pronghorn on the southern plains, 1806-1857. *Plains Anthropologist* 42(159, Memoir 29):163-172.

Silvy, N. J. and C. A. Hagen. 2004. Introduction: management of imperiled prairie grouse species and their habitat. *Wildlife Society Bulletin* 32:2-5.

Silvy, N. J., M. J. Peterson, and R. R. Lopez. 2004. The cause of the decline of pinnated grouse: the Texas example. *Wildlife Society Bulletin* 32:16-21.

Smeins, F. E., and D. D. Diamond. 1983. Remnant grasslands of the Fayette Prairie, Texas. *The American Midland Naturalist* 110:1-13.

Sparks, J. C. and R. E. Masters. 1996. Fire seasonality effects on vegetation in mixed-, tall- and southeastern pine-grassland communities: a review. *Transactions of the 61st North American Wildlife and Natural Resources Conference* 61:246-255.

- Steuter, A. A. 1986. Fire behavior and standing crop characteristics on repeated seasonal burns-northern mixed prairie. Pages 54-59 in A. L. Koonce, ed., Prescribed burning in the Midwest: State-of-the-art, Proceedings of a symposium. University of Wisconsin, Stevens Point. 162 p.
- Stewart, O. C. 1951. Burning and natural vegetation in the United States. *Geographical Review* 41:317-320.
- Stewart, O. C. 1963. Barriers to understanding the influence of use of fire by aborigines. *Proc. Tall Timbers Fire Ecology Conference* 2:117-126.
- Stewart, O. C. 2002. Forgotten fires, Native Americans and the transient wilderness. Edited by H. T. Lewis and M. K. Anderson. University of Oklahoma Press, Norman. 364p.
- Tharpe, B. C. 1925. Structure of Texas vegetation east of the 98th meridian. *University of Texas Bulletin* 2606, University of Texas, Austin.
- Vogl, R. J. 1974. Effects of fire on grasslands. Pages 139-194 in T. T. Kozlowski, and C. E. Ahlgren (eds.). *Fire and ecosystems*. Academic Press, New York.
- U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (2002, December). Fire Effects Information System, [Online]. Available: <http://www.fs.fed.us/database/feis/>.
- Vinton, M.A., and D.C. Hartnett. 1992. Effects of bison grazing on *Andropogon gerardii* and *Panicum virgatum* in burned and unburned tallgrass prairie. *Oecologia* 90:374-382.
- Wright, H. A., and A. W. Bailey. 1982. *Fire ecology: United States and southern Canada*. John Wiley and Sons, New York. 501p.

PERSONAL COMMUNICATION (if applicable):

Sam Fuhlendorf, Assistant Professor, Oklahoma State University