

THE ROLE OF BIOCONTROL OF EMERALD ASH BORER

IN PROTECTING ASH REGENERATION AFTER INVASION



Jian J. Duan¹
Roy G. Van Driesche²
Leah S. Bauer³
Richard Reardon⁴
Juli Gould⁵
Joseph S. Elkinton²

¹USDA, Agricultural Research Service, Newark, DE

²Dept. Environmental Conservation, Univ. Massachusetts, Amherst, MA

³USDA Forest Service, Lansing, MI

⁴USDA Forest Service, Morgantown, WV

⁵USDA, Animal and Plant Health Inspection Service, Buzzards Bay, MA



Forest Health Assessment and
Applied Sciences Team



The Forest Health Technology Enterprise Team (FHTET) was created in 1995 by the Deputy Chief for State and Private Forestry, USDA Forest Service, to develop and deliver technologies to protect and improve the health of American forests.

FHTET became Forest Health Assessment and Applied Sciences Team (FHAASST) in 2016. This booklet was published by FHAASST as part of the technology transfer series.

<http://www.fs.fed.us/foresthealth/technology/>

This publication is available online at:

https://www.fs.fed.us/foresthealth/technology/pdfs/FHAASST-2017-02_Biocontrol_role_EAB_regeneration.pdf

For additional copies of this publication, contact:

Chuck Barger University of Georgia, Center for Invasive Species and Ecosystem Health, (mailing) 2360 Rainwater Road, (physical) 4601 Research Way Room 116, Tifton, GA 31793; 229-386-3298; cbarger@uga.edu

Jian Duan USDA ARS, 501 South Chapel St., Newark, DE 19713; 302-737-6780; Jian.duan@ars.usda.gov

Booklet front cover images: Ash seedlings grow quickly after EAB kills the overstory ash trees, thereby favoring the regeneration and recruitment of native hardwoods over invasive species (Photo: USDA FS). *Spathius galinae*, the most recently released EAB parasitoid, externally parasitizes EAB larvae, protecting large ash trees from EAB damage (Photo: USDA ARS)

KEY FINDINGS

THE GOAL OF THE EAB BIOCONTROL PROGRAM IS TO PROTECT ASH REGENERATION AND RESTORE CANOPY COVER OVER TIME

- Long-term monitoring in Michigan and several northeastern states has documented increasing parasitism and reduced EAB attack rates.
- Ash regeneration is currently benefiting from releases of introduced parasitoids, which now cause 20–80% parasitism of EAB larvae in ash saplings (1–2 inch dia.) and young trees (5–8 inch dia.).
- *Spathius galinae* is well established at release sites in Michigan, New York, and Connecticut, where it attacks 13–49% of EAB larvae. It spread at least 1 mile from points of release, where surveys have been done, within 2 years of release. This parasitoid will begin protecting most large ash trees over the next five years and will provide continuity of protection for young ash trees as they mature, allowing ash regeneration in EAB-damaged forests.



Emerald ash borer (EAB), *Agrilus planipennis* (Photo: Pennsylvania Department of Conservation and Natural Resources – Forestry Archive)

BACKGROUND INFORMATION

Emerald ash borer (EAB) is an invasive Asian beetle that is destroying ash in forests over much of eastern North America because of the high susceptibility of our native ash and a lack of effective natural enemies (**Fig. 1**). To increase mortality of EAB larvae and eggs, the USDA (FS, ARS, APHIS) is carrying out a biological control program based on importation of parasitoids from the beetle's native range in northeastern China and the Russian Far East. Four such parasitoids have been approved for release since 2007 (**Figs. 2–5**), and these species are being reared at the USDA APHIS EAB Biocontrol Facility in Brighton, Michigan, for release in the United States. Three of these natural enemies are now established at various locations and are helping lower EAB densities.

Detailed information on EAB, biocontrol, release site selection, and how to request parasitoids is in the “EAB Biocontrol Release and Recovery Guidelines,” available at www.MapBiocontrol.org.



Fig. 1. Typical damage to native ash trees caused by an infestation of the invasive Asian beetle, Emerald Ash Borer (Photo: USDA FS)





Fig. 2. Egg parasitoid *Oobius agrili* (Photo: USDA ARS)



Fig. 3. Larval parasitoid *Tetrastichus planipennis* (Photo: USDA ARS)




Fig. 4. Larval parasitoid *Spathius agrili* (Photo: USDA ARS)



Fig. 5. Larval parasitoid *Spathius galinae* (Photo: USDA ARS)

How Will Asian Parasitoids Help Protect Ash?



We know that in Asia, parasitoids cause high mortality of EAB, while in North America our native parasitoids only sometimes do so. While North American ash are less resistant than Asian ash, American ash can tolerate some level of EAB attack.

Conversely, woodpeckers, which are scarce in northeastern

Asia, are common in North America and kill about 30–50% of surviving large EAB larvae. The goal of the EAB biocontrol program is to lower survival of EAB eggs and larvae by establishing more effective Asian parasitoids. The combination of these introduced parasitoids, together with moderate levels of tree resistance, high woodpecker predation, and some attack by native parasitoids, will reduce EAB densities and lower the number of attacks per tree to levels that ash trees can survive. This lower attack rate is already happening at Michigan study sites and is allowing ash saplings (1–2 inch dia.) and young ash (5–8 inch dia.) to survive and produce seed for ash regeneration. We have made considerable progress over the last twelve years documenting the impacts of these factors, as described below. Moreover, in the aftermath of EAB in southeast Michigan forests where *T. planipennis* is protecting young ash, recruitment and growth of native hardwoods are favored over invasive plant species.



What Gains Has the EAB Biocontrol Program Made?

- **Release of four EAB parasitoids from Asia.**
 - *Oobius agrili*, egg parasitoid from China, released since 2007
 - *Tetrastichus planipennisi*, internal larval parasitoid from China, since 2007
 - *Spathius agrili*, external larval parasitoid from China, since 2007
 - *Spathius galinae*, external larval parasitoid from Russia, since 2015
- **Establishment of three parasitoid species confirmed.** *Oobius agrili* and *T. planipennisi* are widely established in various areas, and releases continue as EAB enters new areas. *Spathius galinae* is established at several sites in Connecticut, Massachusetts, Michigan, and New York after just two years of releases.
- ***Oobius agrili*** causes up to 22% parasitism of EAB eggs at monitored Michigan study sites and is spreading slowly.
- ***Tetrastichus planipennisi*** is spreading rapidly, causing ~20% parasitism in young ash trees (**Fig. 6**) and 50–80% parasitism in ash saplings (**Fig. 7**).
- ***Spathius galinae*** attacks 13–49% of EAB larvae at release sites in Michigan and Connecticut, and has spread up to 1 mile from points of release.
- **90% reduction in EAB attacks on surviving trees.** In Michigan, EAB attack on surviving ash (larvae per m² of phloem area) is now <10% of the peak in 2009 (**Fig. 8**).
- **Avoidance of plant invasions.** In southern Michigan, where forests are recovering from the loss of ash overstory trees, native hardwoods are favored over invasive plants where ash seedlings, saplings, and young trees are protected by *T. planipennisi*.






Fig. 6. (A). EAB larval parasitism by the introduced parasitoid *Tetrastichus planipennisi* in young ash trees (5–8 inch dia.) at release and non-release control plots, after parasitoid releases in monitored Michigan study sites starting in 2007 (arrows show number of adult parasitoids released by year) (Duan et al. 2015. *J. Appl. Ecol.* 52:1246–1254); **(B).** *T. planipennisi* larvae developed inside the parasitized EAB larva (Inset photo: USDA ARS)




Fig. 7. EAB larval parasitism by *Tetrastichus planipennisi* in ash saplings (1–2 inch dia.) from 2013–2015 at the monitored Michigan study sites (Duan et al. 2017. *For. Ecol. Manag.* 394:64–72)




Fig. 8. EAB larval densities declined in the surviving ash trees at Michigan study sites after death of the larger ash trees (2008–2009). The parasitoids, released from 2007–2010, became established in the surviving ash and EAB parasitism increased from 2012–2014 (see Fig. 6) (Duan et al. 2015. *J. Appl. Ecol.* 52:1246–1254)



What is Likely to Happen in the Next Few Years?

Based on (1) our work during the past 12 years in Michigan and 3–6 years in the northeastern United States, (2) the likely imminent spread of EAB to the South and West, and (3) the species of ash in those not-yet-invaded areas, we expect

- **That** small ash (1–8 inch dia.), now and in the near future, will be protected by *O. agrili* and *T. planipennisi*, allowing ash regeneration in the northeastern and central United States.
- **That** within 5–10 years, as current young ash reach mid-size (9–15 inch dia.), *S. galinae* will be abundant and able to protect these trees as they enter the mid-size cohort, allowing ash regeneration (**Fig. 9**) and regrowth to continue.



Fig. 9. Ash seedlings grow quickly after EAB kills the overstory ash trees, thereby favoring the regeneration and recruitment of native hardwoods over invasive species. (Photo: USDA FS, also used on booklet front cover)

What Do We Need for the Not-Yet-Invaded Regions in the South and West?

■ **New regions may need different parasitoids.**

States just recently being invaded in the South (Louisiana and nearby states) or likely to be invaded soon in the West (California through Oregon) and Southwest (Arizona and nearby states) may need DIFFERENT ASIAN PARASITOIDS (due to climatic differences between these states and northeastern China or the Russian Far East, the source of the parasitoids we are now releasing). Efforts to find and collect EAB parasitoids in southern and western China are underway.

■ **South.** In 2017, work began in southern China (at about the latitude of mid-Florida) in a newly located natural ash stand in an area potentially matching Louisiana. To date, however, no EAB populations have been detected in this part of China.

■ **West.** In 2017, a new EAB population was found in western China in an area with a climate similar to that of the Interior Great Basin (Utah and surrounding states). This EAB population needs to be sampled for parasitoids.

■ **Southwest.** Several rare species of ash are native to Arizona and surrounding southwestern states. Conservation and restoration plans should be developed before EAB invades the region as these native ash species have small distributions and are expected to suffer high mortality from EAB.



The use of trade, firm, or corporation names in this publication is for the information and convenience of the reader. Such use does not constitute an official endorsement or approval by the U.S. Department of Agriculture or the Forest Service of any product or service to the exclusion of others that may be suitable.

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotape, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at http://www.ascr.usda.gov/complaint_filing_cust.html and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: program.intake@usda.gov.

USDA is an equal opportunity provider, employer, and lender.

Booklet back cover image: Young ash trees and saplings shown here are protected by introduced EAB parasitoids, especially by the larval parasitoid *Tetrastichus planipennis*, which attack about 20% of large EAB larvae in young trees (5–8 inch dia.) and 50–85% in saplings (1–2 inch dia.) (Photo: USDA FS)



Federal Recycling Program
Printed on Recycled Paper