



United States Department of Agriculture

Draft Federally Recognized Species: Threatened, Endangered, Proposed and Candidate Species Resource Assessment

Tongass National Forest Plan Revision



Forest
Service

Alaska
Region

Tongass
National Forest

December 2024

Cover Photo: Humpback Whale and calf. Hannes Klostermann, HK-Underwater.com.

In accordance with Federal civil rights law and US Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotape, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at www.ascr.usda.gov/complaint_filing_cust.html and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by: (1) mail: US Department of Agriculture Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue SW, Washington, DC 20250-9410; (2) fax: (202) 690-7442; or (3) email: program.intake@usda.gov.

USDA is an equal opportunity provider, employer, and lender.

Federally Recognized Species: Threatened, Endangered, Proposed and Candidate Species Resource Assessment

Tongass National Forest Plan Revision

Forest Service, Alaska Region

Prepared by: Bonnie Bennetsen
ESA Consultation Biologist
Pacific Planning Service Group

This page intentionally left blank.

Table of Contents

Introduction.....	5
Resource Importance	5
Resource History and Current Management Direction.....	9
Resource History	9
Current Management Direction	9
Scope and Scale of Assessment	9
Status and Trends.....	10
Current Status of Habitat Conditions and Trends (for all federally recognized species)	10
Stressors and Drivers	17
Key Takeaways.....	20
References.....	22
Glossary	29

Tables

Table 1. Federally recognized species relevant to the Tongass. ESU = evolutionary significant unit. DPS = distinct population segment.	5
Table 2. Steller Sea Lion designated critical habitat within or near the Tongass. Estimated area of overlap (acres) is approximate and subject to change as shoreline geospatial data improves. LUD = land use designation. NA = not applicable (site does not overlap with the Tongass).....	6

Figures

Figure 1. Steller Sea Lion designated critical habitat, other Steller Sea Lion haulout sites, and Humpback and Sperm Whale Biologically Important Areas within and near the Tongass.	7
Figure 2. Existing marine access facilities across the Tongass, including USFS and non-USFS permitted log transfer facilities, USFS recreation buoys, and other USFS facilities.	8
Figure 3. Humpback whale.	11
Figure 4. Stellar Sea Lion.	14
Figure 5. Sunflower sea star.....	16

This page intentionally left blank.

Introduction

This assessment report presents the federally listed threatened and endangered species, species proposed for Federal listing, and candidate species, together with the ecological conditions needed to support these species and status of ecological conditions in the plan area. The 2012 Planning Rule states that assessments must “identify and evaluate existing information relevant to the plan area” for threatened, endangered, proposed, and candidate species (36 CFR 219.6(b)(5)). Species evaluations include information regarding each species’ occurrence, habitat status, and population status in the plan area and other appropriate scale(s), to the extent that these are known. Each species evaluation also considers drivers, threats, and stressors for that species or its habitat, and where possible, historic condition along with expected trends for that species or habitat. Not all species have the same kind or amount of information available, so not all species evaluations will appear the same in organization and content.

Resource Importance

The Tongass National Forest features a narrow mainland strip of steep, rugged mountains and icefields dominated by largely intact, pristine temperate rainforest, and over 1,000 offshore islands (USDA 2016). Together, the islands and mainland result in approximately 11,000 miles of meandering shoreline, with numerous bays and coves. A system of seaways separate the many islands and provides a protected waterway called the Inside Passage.

Federally recognized, important species on the Tongass National Forest include the threatened Mexico distinct population segment (DPS) of the humpback whale; the endangered fin and sperm whales; the western DPS of the Steller sea lion and its designated critical habitat; the endangered short-tailed albatross; and the proposed sunflower sea star (Table 1).

Table 1. Federally recognized species relevant to the Tongass. ESU = evolutionary significant unit. DPS = distinct population segment.

Class	Species	Scientific Name	ESU/DPS	ESA Status
Marine Mammal	Humpback Whale	<i>Megaptera novaeangliae</i>	Mexico	Threatened
Marine Mammal	Fin Whale	<i>Balaenoptera physalus</i>	NA	Endangered
Marine Mammal	Sperm Whale	<i>Physeter microcephalus</i>	NA	Endangered
Marine Mammal	Steller Sea Lion	<i>Eumetopias jubatus</i>	Western	Endangered + Critical Habitat
Bird	Short-Tailed Albatross	<i>Phoebastria albatrus</i>	NA	Endangered
Invertebrate	Sunflower Sea Star	<i>Pycnopodia helianthoides</i>	NA	Proposed

Marine waters in Southeast Alaska are characterized geologically by a narrow continental shelf and deep channels (Weingartner et al. 2009). Winds, freshwater runoff, tides, and cross-shelf exchange control the regional oceanography (Weingartner et al. 2009). This complex setting leads to heterogeneous physical processes and numerous and diverse marine biological habitats (Weingartner et al. 2009), generally in highly productive and functional conditions. Nearshore marine waters further benefit from the

surrounding relatively pristine and largely undeveloped Tongass temperate rainforest, its freshwater inputs and productive ecosystems.

The Tongass supports important nearshore marine waters utilized by federally recognized species. These waters are important feeding habitat for all relevant federally recognized species, and breeding and resting habitat for the western DPS of the Steller sea lion and the sunflower sea star. There are approximately 1,200 acres of designated critical habitat for the endangered western DPS of the Steller sea lion within Tongass National Forest System lands above mean high water (Table 2, Figure 1), but note that this estimate of overlap is subject to change as our geospatial shoreline mapping improves.

Table 2. Steller Sea Lion designated critical habitat within or near the Tongass. Estimated area of overlap (acres) is approximate and subject to change as shoreline geospatial data improves. LUD = land use designation. NA = not applicable (site does not overlap with the Tongass).

Site	Tongass	LUD Category	Estimated Critical Habitat on the Tongass (acres)
Benjamin Island	No	NA	NA
Biali Rock	Yes	Wilderness	26
Biorka Island	Yes	Natural Setting	2
Cape Addington	Yes	Natural Setting	108
Cape Cross	Yes	Wilderness	303
Cape Fairweather	No	NA	NA
Cape Ommaney	Yes	Natural Setting	116
Coronation Island	Yes	Wilderness	53
Forrester Island	No	NA	NA
Gran Point	Yes	Natural Setting	297
Graves Rock	No	NA	NA
Hazy Islands	No	NA	NA
Lull Point	Yes	Natural Setting	244
Sunset Island	Yes	Natural Setting	44
Timbered Island	Yes	Wilderness	15
White Sisters	Yes	Wilderness	11
TOTAL	11 sites overlapping Tongass	All Tongass sites in Natural Setting or Wilderness	1219

Given the coastal nature of the Tongass, many authorized and permitted activities may indirectly involve marine transport or development and maintenance of shoreline and marine infrastructure and access facilities. There are several existing USFS and non-USFS log transfer facilities, USFS other marine access facilities, and USFS recreation buoys across Tongass shorelines and nearshore waters (Figure 2).




Figure 1. Steller Sea Lion designated critical habitat, other Steller Sea Lion haulout sites, and Humpback and Sperm Whale Biologically Important Areas within and near the Tongass.




Figure 2. Existing marine access facilities across the Tongass, including USFS and non-USFS permitted log transfer facilities, USFS recreation buoys, and other USFS facilities.

Resource History and Current Management Direction

Resource History

Federal lands comprise about 95 percent of Southeast Alaska, with about 80 percent in the Tongass National Forest (and most of the rest in Glacier Bay National Park and Preserve). The remaining land is held in state, Native Corporation, and local community private ownerships.

Most of the area of the Tongass is wild and undeveloped. Approximately 74,000 people live in the towns, communities, and villages of Southeast Alaska, most of which are located on islands or along the mainland coasts (USDA 2016). The communities of Southeast Alaska depend on the Tongass National Forest ecosystem services, including employment in wood products, commercial fishing and fish processing, recreation, tourism, and mining and mineral development. Many residents also depend on subsistence hunting and fishing to meet their basic needs. Most of these uses require marine transport. Commercial marine traffic is substantial in Southeast Alaska as well, and includes cruise ships, cargo ships, intra- and inter-state ferries, commercial and sport fishing vessels, tour boats, and tug boats and barges. In 2018, 644 different commercial vessels > 65 feet in length plus tugs with Automatic Identification System stations were recorded in Southeast Alaska; an additional 473 were excluded that did not enter State waters within 3 miles of shoreline (Miller et al. 2019). Most of these were fishing vessels 65-150 feet in length. Together these 644 vessels travelled nearly 2.3 million miles in Southeast Alaska in 2018, with most miles (29 percent) by cruise ships (Miller et al. 2019). These vessels travelled into nearly every waterway in Southeast Alaska and especially the primary channels. This study did not report the numerous smaller watercraft used for commercial and recreational boating and marine transit needs in Southeast Alaska, especially focused around communities.

Current Management Direction

Legal and Regulatory Compliance

Current Forest Plan direction includes standards and guidelines to help protect and maintain federally recognized marine mammal habitats consistent with the Marine Mammal Protection Act, Endangered Species Act, and National Marine Fisheries Service guidelines for approaching marine mammals. The level of indirect influence from Tongass authorized and approved activities on marine transit has not been quantified, but likely involves very minor contributions compared to other sources of marine traffic such as from public recreation, non-USFS-permitted cruise industry and other tourism, and commerce.

Scope and Scale of Assessment

For this assessment, federally recognized species relevant to the Tongass were evaluated in the context of categories directed and defined by the National Forest System Land Management Planning Final Rule and Record of Decision (hereafter 2012 Planning Rule), detailed in the Code of Federal Regulations (36 CFR 219 2012). The Forest Service Handbook provides specific, detailed direction called directives for implementing planning rules.

Most native species' needs would be covered under broader ecosystem plan components and are not specifically discussed or evaluated in detail. Rather, we focus on the condition, status, and trends for “at-risk species.” Under the 2012 Planning Rule, at-risk species are defined as:

- Federally recognized threatened, endangered, proposed, and candidate species; and

-
- Species of Conservation Concern (SCC).

The 2012 Planning Rule requires the identification of SCC, which are “species, other than federally recognized threatened, endangered, proposed, or candidate species, that are known to occur in the plan area and for which the Regional Forester has determined that the best available scientific information indicates substantial concern about the species’ capability to persist over the long- term in the plan area” (36 CFT 219.9 (c)). This topic is discussed in detail in the Species of Conservation Concern assessment section. The directives also call for identifying information regarding fish, wildlife, and plant species that may be commonly enjoyed and used by the public for hunting, fishing, trapping, gathering, observing, or sustenance, and assessing the conditions and trends of these species and their habitats. These considerations are discussed in the Subsistence and Other Harvest assessment section, focusing on the contributions of those species to social and economic sustainability and subsistence of rural communities in Southeast Alaska.

The 2012 Planning Rule recognizes that it may not be possible to maintain a viable population of some at-risk species within the plan area due to circumstances beyond the authority of the Forest Service or due to limitations in the inherent capability of the land. Examples might be migratory species where viability is primarily affected in other locations, climate-sensitive species affected by warming temperatures, or where the plan area has limited ecological capacity to provide sufficient habitat to sustain the species.

Status and Trends

Current Status of Habitat Conditions and Trends (for all federally recognized species)

The current condition of beach, estuary, and terrestrial marine habitats within and adjacent to the Tongass National Forest has been moderately degraded in localized areas, especially on lands of other ownership in cities and towns. However, Tongass habitats and the vast majority of shoreline and marine habitats within Southeast Alaska remain intact, function properly, and continue to be productive for wildlife and fish. Common vegetation and habitat management activities include timber harvest, young-growth restoration projects, and aquatic organism passage and riparian restoration projects. There are also active, permitted mines in the area. Other actions or authorized activities are related to recreation, public use, maintenance, and monitoring. All of these activities may require marine transport, associated beach landings, and occasionally shoreline or marine infrastructure.

Humpback Whale

The humpback whale shown in Figure 3 was listed as endangered in 1970, first under the Endangered Species Conservation Act and subsequently under the Endangered Species Act (ESA) when it passed in 1973.



Figure 3. Humpback whale.

From the late 1800s through the early 1900s, humpback whales were extensively commercially hunted. Signs of recovery since the end of commercial whaling led NMFS in 2016 to conduct a global status review and publish a rule to change the status of humpback whales under the ESA. Under the final rule, the Mexico DPS (which includes a small portion of humpback whales found in Southeast Alaska) was delisted to threatened, and the Hawaii DPS (which includes most humpback whales found in Southeast Alaska) was removed from listing (81 FR 62260; September 8, 2016). Updated probability of occurrence data indicates individual humpback whales in Southeast Alaska have a 2% probability of being from the threatened Mexico DPS and a 98% probability of being from the recovered Hawaii DPS (Wade 2021, NMFS 2021). In the Gulf of Alaska, individual humpback whales have an 11% probability of being from the threatened Mexico DPS and an 89% probability of being from the recovered Hawaii DPS (Wade 2021, NMFS 2021). The endangered Western North Pacific DPS also has a chance of occurring off the coast of Yakutat (Wade 2021, NMFS 2021). There is evidence of mixing between the Mexico and Hawaii DPSs on their wintering grounds (Darling et al. 2022), and movement and genetics data suggest further population structure within the Mexico DPS (Martien et al. 2021).

Critical habitat was designated for the Mexico DPS in 2021, but none was identified within waters surrounding the Tongass. The humpback whale status review (Bettridge et al. 2015), the final 2016 rule revising the species-wide listing under the ESA (81 FR 62260, September 8, 2016), the recovery outline for the Mexico and other DPSs (NMFS 2022a) along with the original recovery plan (NMFS 1991), and information on the NMFS (2024a): online species account, as well as other best available scientific information provide science on this species and a foundation for this species' assessment.

Distribution within the boundaries of the Tongass

The Mexico DPS spends winters along the Pacific coast of mainland Mexico and in the Revillagigedo Archipelago, transits along the coast of Baja California, and spends summers feeding throughout the North Pacific from California to the Kamchatka Peninsula in Russia (NMFS 2022a: Calambokidis et al. 2008, Titova et al. 2018, Wade 2021). Humpback whales, including predominantly Hawaii DPS but also Mexico DPS individuals, are regularly sighted in the inside marine waters of Southeast Alaska, including shallow coastal areas around the Tongass. This species generally migrates long distances from tropical waters where they calve during the winter to cooler, northern coastal waters where they feed during the

rest of the year. The highly productive fjords of Southeast Alaska are an important feeding area. Nearly all of the Inside Passage marine water channels surrounding the Tongass have been identified by NMFS as a humpback whale Biologically Important Area that supports concentrated feeding activities (Figure 1). Humpback whales are typically observed in nearshore waters of Southeast Alaska from approximately May through December, with peak numbers during late August and September and substantial numbers remaining through early winter, although individuals can be seen every month of the year.

The local distribution of humpback whales in Southeast Alaska appears to be correlated with the density and seasonal availability of prey, particularly herring (*Clupea harengus*) and krill (*euphausiids*). Important local feeding areas include Glacier Bay and adjacent portions of Icy Strait, Stephens Passage/Frederick Sound, Seymour Canal and Sitka Sound. Glacier Bay and Icy Strait appear to be important early in the season, when whales prey heavily on herring and other small, schooling fishes. Frederick Sound is important later in summer, when whales feed on swarming euphausiids. During autumn and early winter, humpbacks move out of the Sound to areas where herring are abundant, particularly Seymour Canal. Late fall and winter habitat in SE Alaska appears to correlate with areas with over-wintering herring, such as lower Lynn Canal, Tenakee Inlet, Whale Bay, Ketchikan, Sitka Sound, and Glacier Bay/Icy Strait. Other areas of Southeast Alaska may also be important for humpbacks, including Cape Fairweather, Sumner Strait, Dixon Entrance, the west coast of Prince of Wales Island, and offshore banks such as the Fairweather Grounds.

Population status and trend

North Pacific humpback whale populations were estimated at approximately 15,000 individuals prior to commercial whaling overexploitation and subsequently declined to an estimated 1,200 by the end of commercial whaling (Teerlink et al. 2015). Populations have bounded back since then, and recent estimates range from 16,293 to 21,808 (CV=0.04; NMFS 2022a: Calambokidis et al. 2008; Barlow et al. 2011; Wade 2021), with annual growth rates estimated at 6.8 percent from 1966 to 2006 (NMFS 2022a: Calambokidis et al. 2008).

There is no current abundance estimate for the Mexico DPS, although an estimated 3,477 (CV=0.101) whales from the Mexico DPS feed off the U.S. West Coast (NMFS 2022a: Calambokidis and Barlow 2020, Curtis et al. 2022). In Southeast Alaska, where Hawaii and Mexico DPSs overlap, past population estimates were 300 in 1979-1983 (Baker et al. 1985), 547 in 1986 (95% CI 504-590; Baker et al. 1992), 404 in 1985-1992 (Straley 1994), and 961 for northern Southeast Alaska in 2000 (95% CI 657-1076; Straley et al 2009). While the current population trend for the Mexico DPS is also unknown, Calambokidis and Barlow (2020) reported an approximate 8.2% annual growth rate from 1989-2018 for humpback whales off California and Oregon waters, where whales from the Mexico and Central America DPSs overlap (NMFS 2022a).

Abundance and calf production rates of humpback whales recently declined in Glacier Bay and Icy Strait in Southeast Alaska, presumably due to disruption of lower trophic level prey after a marine heatwave in the Gulf of Alaska in 2014-2016 (Neilson and Gabriele 2021). The marine heatwave caused dramatic changes in population dynamics and the Glacier Bay National Park and Preserve long-term dataset (1985-2020) indicated sharp declines in humpback whale survival and reproductive success coinciding with the marine heatwave (Neilson and Gabriele 2021, Gabriele et al. 2022). Evidence of a decline in the encounter rate of humpback whales and the number of calves in Prince William Sound (Arimitsu et al. 2021) indicates this decline likely occurred widely throughout the Gulf of Alaska. There is therefore considerable uncertainty about current actual population size and trends and it is unclear if the Mexico DPS is still increasing overall (Young et al. 2023).

Fin Whale

Fin whales, *Balaenoptera physalus*, have been listed as endangered since 1970 under the precursor to the ESA and have retained this status since the ESA was passed in 1973. Though this species was not initially targeted by commercial whalers due to its speed and open ocean habits, as whaling methods advanced, efforts turned to fin whales and decimated their populations. Critical habitat has not been designated for fin whales. The original ESA listing (35 FR 8491; June 2, 1970), the recovery plan (NMFS 2010), the fin whale Northeast Pacific stock assessment (Muto et al. 2021a), information on the NMFS (2023a): online species account, as well as other best available scientific information provide science on this species and a foundation for this species' assessment.

Distribution within the boundaries of the Tongass

Fin whales are typically found in deep, offshore waters. They occur year-round in a wide range of locations, but the density of individuals in any one area changes seasonally. Most migrate from the Arctic (or Antarctic) feeding areas in the summer to tropical breeding and calving areas in the winter, though the location of winter breeding grounds is not known. Fin whales tend to travel in the open seas, away from the coast. In Alaska, fin whales are typically found in offshore marine waters of the Bering Sea, Chukchi Sea, Cook Inlet, Aleutian Islands, Gulf of Alaska and Southeast Alaska in proximity to the open ocean. There have been two sightings in lower Clarence Strait (Dahlheim et al. 2009) and they only rarely occur in the Inside Passage. They have been sighted off the coast of Prince of Wales Island, Coronation Island, Sitka Sound, Yakutat, and Chatham Strait.

Population status and trend

Fin whales are divided into four stocks for management purposes, the Alaska (Northeast Pacific) being one of these. Reliable, recent population estimates and trends are not available for the Northeast Pacific stock. The best available estimate from 2013 indicates a minimum stock population estimate of 2,554 fin whales, but this is viewed as an underestimate (Muto et al. 2021a). Zerbini et al. (2006) estimated annual increase of a portion of the stock at 4.8 percent between 1987 and 2003, though caution is advised in interpreting this result due to uncertainties related to the initial population and population structure of fin whales in the area (Muto et al. 2021a).

Sperm Whale

Sperm whales were listed as endangered under the Endangered Species Conservation Act in 1970 (35 Federal Register 233: 18319) and have been listed under the ESA since it became law in 1973. Sperm whales were a primary target of the commercial whaling industry from 1800 to 1987, which nearly decimated all sperm whale populations. While whaling is no longer a major threat, sperm whale populations are still recovering. The original ESA listing (35 FR 8491; June 2, 1970), the recovery plan (NMFS 2010), the fin whale Northeast Pacific stock assessment (Muto et al. 2021b), and information on the NMFS (2023b): online species account, as well as other best available scientific information provide science on this species and a foundation for this species' assessment.

Distribution within the boundaries of the Tongass

In general, sperm whales are considered deep water species, usually found in waters 600 meters or deeper and are uncommon in waters less than 300 meters. Acoustic surveys, from fixed autonomous hydrophones, detected the presence of sperm whales year-round in the Gulf of Alaska, though they appear to be approximately two times as common in summer than in winter (Mellinger et al. 2004, Muto et al. 2021b). This seasonality of detections is consistent with the hypothesis that sperm whales generally move to higher latitudes in summer and to lower latitudes in winter (Muto et al. 2021b). NMFS identified a Biologically Important Area for sperm whales in the Gulf of Alaska, off the coast of the Tongass (Figure

1). Sperm whales have also been documented using the inside waters of Southeast Alaska. The Southeast Alaska Sperm Whale Avoidance Project (Straley et al. 2015) tracked the movement of several individuals into inside marine waters surrounding the Tongass. The study showed several tagged whales migrating through Chatham Strait as far north as the Lynn Canal with one individual moving through Icy Strait. In 2019 a dead sperm whale washed up on the east side of Lynn Canal, north of Berners Bay near Juneau showing evidence consistent with a ship strike (NMFS 2019).

Population status and trend

Current and historical abundance estimates and associated population trends of sperm whales in the North Pacific are based on limited data and are considered unreliable (Muto et al. 2021b). Sperm whale numbers were heavily depleted by commercial whaling and populations are still recovering and likely increasing. North Pacific stock abundance for sperm whales prior to exploitation is believed to have been around 1,260,000 and reduced to around 930,000 by the late 1970s (no confidence intervals available; Allen and Angliss 2015, Muto et al. 2021b). Estimates for a large area of the eastern temperate North Pacific were produced from line-transect and acoustic survey data by Barlow and Taylor (2005); the acoustic data produced an estimate of 32,100 sperm whales (coefficient of variation (CV) = 0.36; Muto et al. 2021b). From surveys in the Gulf of Alaska in 2009 and 2015, Rone et al. (2017) estimated 129 (CV = 0.44) and 345 sperm whales (CV = 0.43) in each year, respectively (Muto et al. 2021b). Again, caution in interpreting these data is warranted (Muto et al. 2021b).

Steller Sea Lion

Steller sea lions (Figure 4) were listed as threatened under the ESA in 1990 due to unexplained widespread population declines (55 FR 49204). In 1997 NMFS reclassified the population into two distinct population segments; the threatened eastern DPS which inhabits waters east of 144 °W longitude (a line near Cape Suckling, AK) through Southeast Alaska, British Columbia, down to California and the endangered western DPS, which includes all Steller sea lions originating from rookeries west of this line into the Aleutians, Pacific Rim, and Japan (62 FR 24345; 62 FR 30772). The eastern DPS has recovered and was delisted in 2013 (78 FR 66139), but the western DPS remains listed as endangered.



Figure 4. Steller Sea Lion.

Distribution within the boundaries of the Tongass

Jemison et al. (2013) found that there is regular movement of western DPS Steller sea lions across the 144° W longitude boundary. Most of the cross-boundary movements are temporary with individuals returning to their natal area for breeding, but some females from the western DPS have likely emigrated permanently and have given birth to pups at White Sisters and Graves Rocks rookeries. Most confirmed sightings of western DPS animals have been in northern areas of Southeast Alaska, especially north of Frederick Sound (Jemison et al. 2013). Several confirmed sightings have been recorded at haulouts in Frederick Sound; and there are a few confirmed observations of western DPS animals south of a line from Hazy Island, through Sumner Strait, to the northern tip of Wrangell Island (NMFS 2013).

Sea lions obtain 100 percent of their diet from the marine environment but depend on terrestrial environments for rookeries (birthing areas) and haulouts (non-birthing resting and loafing areas). Haulouts and rookeries tend to be preferentially located on exposed rocky shoreline and wave-cut platforms (Ban 2005). Some rookeries and haulouts are also located on gravel beaches. Terrestrial sites tend to be associated with marine waters that are relatively shallow and well-mixed, with average tidal speeds and gradual bottom slopes (Ban 2005). Rookeries are nearly exclusively located on offshore islands and reefs. Sites used as rookeries in the breeding season may also be used as haulouts during other times of year. Some haulouts are used year-round while others only on a seasonal basis (NMFS 2008). When not on land, Steller sea lions are seen near shore and out to the edge of the continental shelf and beyond.

Population status and trend

The western DPS declined by 75 percent between 1976 and 1990 and another 40 percent between 1991 and 2000. Then, between 2000 and 2004 there was a 12% increase in numbers of non-pups counted in the Alaskan range of the western DPS, the first increase observed during more than two decades of systematic surveys. The observed increase, however, was not evenly spread among all regions of Alaska. Increases were noted in the eastern and western Gulf of Alaska and in the eastern and central Aleutian Islands, while the decline persisted through 2004 in the central Gulf of Alaska and the western Aleutian Islands. Western DPS Steller sea lions increased in abundance between 2012 and 2015 in Alaska, continuing the trend that began in the early 2000s (Fritz et al. 2016).

Between 2015 and 2017, pup counts declined in the eastern (-33%) and central (-18%) Gulf of Alaska (Sweeney et al. 2017), likely due to changes in availability of prey associated with the marine heat wave from 2014 to 2016 (Bond et al. 2015, Peterson et al. 2016, von Biela et al. 2019, Yang et al. 2019). In 2019, pup counts rebounded to 2015 levels, but non-pup counts in the eastern, central, and western Gulf of Alaska regions declined (Sweeney et al. 2019). The population currently appears to be relatively stable in the Gulf of Alaska (Young et al. 2023), though signs of recovery have slowed or ceased altogether in the western DPS, and in Southeast Alaska (mostly non-listed eastern DPS, but some western DPS in northern portions) appear to be declining in recent years (Sweeney et al. 2023).

Steller sea lion critical habitat

Critical habitat was designated in 1993, based on the location of terrestrial rookery and major haulout sites, spatial extent of foraging trips, and prey availability (58 CFR § 226.202 at 58 FR 45278, 1993; redesignated and amended in 1999 at 64 FR 14067). Steller sea lion critical habitat includes all designated major haulouts and rookeries, as well as terrestrial, aquatic, and air zones that extend 3,000 feet landward, seaward, and skyward, respectively, from each. NMFS has designated three rookeries and 11 major haulouts along Tongass shorelines (Table 2, Figure 1). In total approximately 1,200 acres of Steller sea lion critical habitat occur within the 3,000 feet terrestrial zone on Tongass lands above mean high water. Note that this estimate is approximate and subject to change as our shoreline geospatial mapping

improves. In addition, there are two other major haulouts designated in southeast Alaska within the boundaries of Glacier Bay National Park at Graves Rock and Cape Fairweather.

The physical and biological habitat features of Steller sea lion critical habitat support reproduction, foraging, rest, and refuge that are essential to the conservation of the Steller sea lion. Factors that influence the suitability of terrestrial sites include substrate, exposure to wind and waves, the extent and type of human activities and disturbance in the region, and proximity to prey resources (Mate 1973). The use of traditional rookery sites and the link of territorial males, postpartum females, and pups here, make them particularly vulnerable to harassment. In some cases, haulout sites can be completely abandoned after repeated disturbances.

The remote locations of most rookeries and haulouts help to reduce the frequency of harassment, but disturbance by air and watercraft continues to occur. Steller sea lions are vulnerable to harassment and disruption of essential life functions (e.g., breeding, pup care, and rest) at rookeries and haulouts throughout their range. In addition to boat and airplane traffic, other Tongass activities such as timber harvest, mineral extraction, and subsistence harvest in the vicinity of critical habitat may disrupt the essential life functions that occur there.

Sunflower sea star

On August 18, 2021, the Center for Biological Diversity petitioned NMFS to list the sunflower sea star (*Pycnopodia helianthoides*) under the ESA, shown in Figure 5. NMFS determined that the proposed action may be warranted (86 FR 73230, December 27, 2021) and began a full status review to evaluate overall extinction risk for the species. NMFS determined that the sunflower sea star is likely to become an endangered species within the foreseeable future throughout its range and on March 16, 2023, published a proposed rule to list the sunflower sea star as a threatened species (88 FR 16212). NMFS did not propose at that time to designate critical habitat.



Figure 5. Sunflower sea star.

Distribution within the boundaries of the Tongass

The sunflower sea star is a large (up to 1 m in diameter), fast-moving (up to 160 cm/minute), many-armed (up to 24) echinoderm native to the west coast of North America. It occupies waters from the intertidal to at least 435 m deep but is most common at depths less than 25 m and rare in waters deeper than 120 m. Sunflower sea stars occur over a broad array of soft-, mixed-, and hard-bottom habitats from the Aleutian Islands, AK, to Baja California, Mexico, but are most abundant in waters off eastern Alaska and British Columbia. The sunflower sea star hunts a range of bivalves, gastropods, crustaceans, and other

invertebrates using chemosensory stimuli and will dig for preferred prey in soft sediment. While generally solitary, they are also known to seasonally aggregate, perhaps for spawning purposes.

Population status and trend

Prior to 2013, the global abundance of sunflower sea star was estimated at several billion animals, but from 2013–2017 sea star wasting syndrome (SSWS) reached pandemic levels, killing an estimated 90 percent or more of the population. Declines in the northern portion of its range were less pronounced than in the southern portion, but estimated mortality still exceeded 60 percent of the population.

Stressors and Drivers

The primary marine stressors in the region include changes in climate trends, coastal development, floating marine debris, marine-based recreation, and recreational and commercial fishing. Marine climate changes include increasing ocean temperatures, acidification, algal blooms and associated toxins, and noise as well as changes in ocean currents, hydrology, tidewater glaciers and sea ice, marine species distributions and food webs, and sea levels countered by isostatic rebound from glacial recession. These changes will be compounded for some species, especially the whales and sea lion, by increasing marine recreation and marine debris in the region.

Sources of habitat degradation occur primarily from marine-based recreation, tourism, and commerce, commercial and recreational fisheries, shoreline and marine infrastructure development, and global marine debris. Vessel traffic and associated underwater noise, disturbance at resting sites, potential for ship strikes of marine mammals, and other forms of habitat degradation is thought to have increased 10-fold in Southeast Alaska since the 1980s (Jansen et al. 2015).

Marine debris has also been increasing globally and regionally (National Marine Fisheries Service (NMFS) 2016), along with associated entanglement risks (NMFS 2014). There is likely to be additional habitat degradation associated with climate change as detailed in the species' Population-level drivers and stressors sections below.

Humpback Whale

Population-level drivers and stressors

Humpback whales in Southeast Alaska today face threats from entanglement in fishing gear, vessel-based harassment, vessel strikes, underwater noise, competition with commercial fisheries, and impacts from changing climate. Because the humpback whale inhabits shallow coastal areas, it is increasingly exposed to human activity, making it more susceptible to most of these stressors. The ability of humpback whales to recover from threats is limited by a slow growth rate, long gestation periods, high maternal investment in offspring, and low fecundity (NMFS 2022a). Further, despite being a highly mobile species, humpback whales may be constrained by the high level of site fidelity they exhibit to both wintering and feeding areas (NMFS 2022a).

Humpback whales can become entangled by many different gear types, including moorings, traps, pots, or gillnets. Once entangled, if they can move the gear, the whale may drag and swim with attached gear for long distances, ultimately resulting in fatigue, compromised feeding ability, or severe injury, which may lead to reduced reproductive success or even death. There is evidence to suggest that most humpback whales experience entanglement over the course of their lives but are often able to shed the gear on their own. Studies of characteristic wounds and scarring indicate that humpback whales in the region experience a high rate of interaction with fishing gear (20-71%; Neilson et al. 2009). However, the portion of whales that become entangled and do not survive is unknown.

Whale watching vessels, recreational boats, and other vessels may cause stress and behavioral changes in humpback whales, leading to vessel-based harassment, though regulations aimed at preventing and minimizing such impacts exist (50 CFR 224.103). Because humpback whales are often found close to shore and generally surface in an active state, they tend to be popular whale watching attractions. Humpback whales are the central attraction for the whale watching industry in Southeast Alaska.

Inadvertent vessel strikes can injure or kill humpback whales. Humpback whales are vulnerable to vessel strikes throughout their range, but the risk is much higher in coastal areas with heavier ship traffic. Vessel size, speeds, and nighttime travel further influence lethality (Neilson et al. 2012, Webb and Gende 2015). Mean annual mortality due to vessel strikes between 2016 and 2020 was estimated at 1.75 whales in Southeast Alaska and 0.14 in the Gulf of Alaska (Young et al. 2023). Much of the vessel traffic is seasonal during summer months, though some occurs year-round (Neilson et al. 2012). There is also generally more vessel traffic in near shore areas putting near shore species like humpback whales at higher risk for collisions (Neilson et al. 2012). This and other vessel stressors are likely increasing along with increases in vessel traffic and a growing whale watching industry (NUKA 2012, Neilson et al. 2012, Jansen et al. 2015).

This population is likely exposed to moderate levels of underwater noise resulting from human activities, such as commercial and recreational vessel traffic and near-shore blasting or pile driving for shoreline infrastructure. Overall population-level effects of exposure to underwater noise are not well established, but exposure is likely chronic in more heavily trafficked waters of Southeast Alaska. As vessel traffic and other activities are expected to increase, the importance of this stressor is expected to increase. Increasing ocean acidification with climate change is likely to further exacerbate underwater noise, as more acidic marine waters decrease sound absorption for frequencies less than or equal to 10 kHz (Hester et al. 2008). Projected acidity of mid-century marine waters are expected to be 40 percent less effective in absorbing sounds in these frequencies (Hester et al. 2008).

Recent work suggests that commercial fisheries of forage fish on the U.S. West Coast may negatively impact baleen whales by affecting a primary prey, though there is high uncertainty and many caveats (NMFS 2022a: Kaplan et al. 2013; Koehn et al. 2017). There is suspected interaction with the herring fishery in Southeast Alaska, but impacts to humpback whales are considered modest and impacts stable because the herring fishery is regulated.

The impacts of climate change on whales are largely unknown, but it is considered one of the largest threats facing high latitude regions where many humpback whales forage. Changes are occurring from increasing ocean temperatures, acidification, algal blooms and associated toxins, and acidification effects on underwater noise as well as changes in ocean currents, hydrology, tidewater glaciers and sea ice, marine species distributions and food webs, and sea levels countered by isostatic rebound from glacial recession. Notably, a marine heatwave from 2014 to 2016 raised concerns about the indirect impacts of climate change and future interactions between multiple threats (NMFS 2022a). The heatwave pushed some prey species, like anchovies, closer to shore (Santora et al. 2020). This subsequently shifted whale distribution and increased whale foraging in nearshore waters where there is more overlap with commercial and recreational fixed-gear fisheries, including that for Dungeness crab. At the same time, there was a change in fishing effort in central California due to a harmful algal bloom at the start of the Dungeness crab fishery season (Santora et al. 2020; Saez et al. 2021). The delay at the start of the Dungeness fishing season in the fall increased fishing effort in the spring, which coincided with the arrival of humpback whales (Saez et al. 2021). These factors resulted in increased exposure of humpback whales to fishing gear and caused significantly more entanglements in the region.

Further, in Southeast Alaska, the marine heatwave caused dramatic changes in population dynamics (Gabriele et al. 2022), including sharp declines in humpback whale survival and reproductive success. The main driver of these declines was thought to be changes to prey availability and quality, and subsequent food limitation. Any resulting changes in prey distribution could lead to changes in foraging behavior, nutritional stress, and diminished reproduction for humpback whales. Additionally, changing water temperature and currents could impact the timing of environmental cues important for navigation and migration.

Fin Whale

Population-level drivers and stressors

Although the main direct threat to fin whales was addressed by the International Whaling Commission moratorium on commercial whaling, several threats remain today. These include collisions with vessels, entanglement in fishing gear, effects of increasing anthropogenic noise in the ocean environment, and reduced prey abundance due to overfishing and/or climate change (Allen and Angliss 2015; Muto et al. 2021a). Collisions with vessels is a major threat; fin whales are considered the most vulnerable species to ship strikes after North Atlantic Right Whales. The projected increase in ship traffic in offshore and arctic waters arising from the opening of trans-polar shipping routes (as arctic sea ice continues to decline) will increase the risk of vessel strike and increase ambient noise and pollution. Fin whales can also become entangled in fishing gear, either swimming off with the gear attached or becoming anchored. Once entangled, whales may drag and swim with attached gear for long distances, ultimately resulting in fatigue, compromised feeding ability, or severe injury, which may lead to reduced reproductive success or death. Underwater noise negatively affects whale populations, interrupting their normal behavior and driving them away from areas important to their survival. Increasing evidence suggests that exposure to intense underwater sound in some settings may cause some whales to strand and ultimately die.

The impacts of change in climate on baleen whales may result from altered oceanographic conditions, as well as the timing and distribution of sea ice coverage. Changes in prey distribution could lead to changes in foraging behavior, nutritional stress, and diminished reproduction for fin whales. Fin whales experienced deleterious effects of a marine heat wave event along with other large whale species in 2015 and 2016 (Young et al. 2023). Additionally, changing water temperature and currents could impact the timing of environmental cues important for navigation and foraging.

Sperm Whale

Population-level drivers and stressors

The sperm whale recovery plan (NMFS 2010) cites a number of potential threats to Pacific stocks but all of these threats are considered low or expected to have little effect on their recovery. Four of the most common threats cited for Alaska sperm whales are interactions with commercial fishing, whale watching, acoustic disturbance and ship strikes. Very little information is available for understanding the frequency of ship strikes but general consensus is that the threat is low (NMFS 2010). Neilson et al. (2012) found that out of the 89 defined whale strikes documented from 1978-2011 only one of those was a sperm whale and the fate of that whale is unknown. A minimum estimate of the mean annual mortality caused by ship strikes is 0.2 (Young et al. 2023). The level of effects on sperm whales from ship noise is not fully understood either but effects are expected to be like those described for humpback whales (NMFS 2010).

Steller Sea Lion

Population-level drivers and stressors

Threats to Steller sea lions include effects of fisheries on prey; climate change influences on sea level rise, temperature changes, ocean acidification, and harmful algal blooms; disease and parasites, potentially exacerbated by climate change; toxic substances such as mercury, organochlorines, and others that bioaccumulate in top predators; human-caused injuries and mortality from active fishing gear, disturbance at terrestrial sites, entanglement in marine debris or fishing gear, illegal feeding, illegal shooting, and vessel strikes; and predation from killer whales, sharks and humans. These threats are further detailed in the NMFS (2023c): [online species account](#) and the Steller sea lion recovery plan (NMFS 2008). The recovery plan (NMFS 2008) ranked environmental variability in prey, competition with fisheries, and predation by killer whales as potentially high threats, toxic substances as medium threat, and incidental take due to interactions with active fishing gear, Alaska native subsistence harvest, illegal shooting, entanglement in marine debris, disease and parasites, disturbance from vessel traffic and tourism, and disturbance due to research activities as low threats. Our understanding of the impacts of these threats on Steller sea lion survival and reproduction and which threats are the largest impediments to Steller sea lion recovery is currently limited (NMFS 2008).

Sunflower sea star

Population-level drivers and stressors

Species-level impacts from sea star wasting syndrome (SSWS), both during the pandemic and on an ongoing basis, have been identified as the major threat affecting the long-term persistence of the sunflower sea star. Although disease, specifically SSWS, is the primary known threat to this species, activities that affect near-shore or deeper benthic communities could also impact this species. Other threats include habitat degradation and destruction, especially in nearshore, urbanized areas, unregulated harvest of the species in some jurisdictions, and both direct and indirect consequences of climate change (NMFS 2022b). Elevated ocean temperatures likely contributed to the precipitous declines of this species as warmer temperatures are associated with increased transmission and mortality rates of SSWS (NMFS 2022b). Further, ocean acidification can increase physiological stress and decrease survival in a broad array of marine organisms, especially in shell-forming prey of this species (NMFS 2022b). Tongass permitted or authorized activities most likely to influence this federally recognized species include shoreline infrastructure development and bark accumulation at log transfer facilities, though the mobility of this species and localized nature of these activities decreases risk.

Key Takeaways

There are no federally recognized plants or fish relevant to the Tongass. Threatened, endangered, candidate and proposed wildlife and invertebrate species relevant to the Tongass are all marine species, including three whales, one sea lion, one seabird, and one sea star (Table 1).

The Tongass:

- Supports but generally does not encompass nearshore marine waters used as foraging habitat by all federally recognized species relevant to the Tongass, and as breeding and resting habitat by the endangered western distinct population segment (DPS) of the Steller sea lion and proposed sunflower sea star.
- Provides approximately 1,200 acres of designated critical habitat for the endangered western DPS of the Steller sea lion within Tongass National Forest Service lands above mean high water.

Critical habitat for this species consists of 3,000 feet landward, seaward, and skyward from three major rookeries (breeding sites) and 11 major haul-outs (resting sites) within and near the Tongass (Table 2, Figure 1).

- Indirectly influences marine systems via Forest Service authorized and permitted activities that may involve boat or float plane transit or involve nearshore, shoreline, or marine infrastructure.
- Is subject to stressors caused by changing climate including increasing ocean temperatures, acidification, algal blooms and associated toxins, and noise as well as changes in ocean currents, hydrology, tidewater glaciers and sea ice, marine species distributions and food webs, and sea levels countered by isostatic rebound from glacial recession. These stressors are being and will continue to be compounded for some species, especially the whales and sea lion, by recent increases in marine recreation in the region.
- Current Forest Plan direction includes measures to protect federally recognized marine mammal habitats consistent with the Endangered Species Act and Marine Mammal Protection Act.

This page intentionally left blank.

References

- Allen, B.M. and R.P. Angliss. 2014. Alaska marine mammal stock assessments, 2014. NOAA Technical Memorandum NMFS-AFSC-301. 304 pp.
- Arimutsu, M.L., J.F. Piatt, S. hatch, R.M. Suryan, S. Batten, M.A. Bishop, R.W. Campbell, H. Coletti, D. Cushing, K. Gorman, R.R. Hopcroft, K.J. Kuletz, C. Marsteller, C. Mckinstry, D. McGowan, J. Moran, S. Pegau, A. Schaefer, S. Schoen, J. Straley, V. R. von Beila. 2021. Heatwave-induced synchrony within forage fish portfolio disrupts energy flow to top pelagic predators. *Global Change Biology* 27: 1859-1878.
- Baker, C.S. 1985. Population characteristics and migration of summer and late season humpback whales (*Megaptera novaeangliae*) in Southeastern Alaska. *Marine Mammal Science* 1(4); 304-323
- Baker, C.S, J.M Straley and A. Perry. 1992. Population characteristics of individually identified humpback whales in southeastern Alaska: Summer and fall 1986. *Fishery Bulletin* 90(3), 430-437
- Ban, S.S. 2005. Modelling and characterization of Steller sea lion haulouts and rookeries using oceanographic and shoreline type data. Thesis, Master of Science, University of British Columbia.
- Barlow, J., J. Calambokidis, E.A. Falcone, C.S. Baker, A.M. Burdin, P.J. Clapham, J.K.B. Ford, C.M. Gabriele, R. LeDuc, D.K. Mattila, T.J. Quinn II, L. Rojas-Bracho, J.M. Straley, B.L. Taylor, J. Urban R., P. Wade, D. Weller. 2011. Humpback whale abundance in the North Pacific estimated by photographic capture-recapture with bias correction from simulation studies. *Marine Mammal Science*, 27(4): 793-818.
- Barlow, J. and B.L. Taylor. 2005. Estimates of sperm whale abundance in the northeastern temperate Pacific from a combined acoustic and visual survey. *Marine Mammal Science*, 21(3):429-445.
- Bettridge, S., C.S. Baker, J. Barlow, P.J. Clapham, M. Ford, D. Gouveia, D.K. Mattila, R.M. Pace III, P.E. Rosel, G.K. Silber, P.R. Wade. 2015. Status review of the humpback whale (*Megaptera novaeangliae*) under the Endangered Species Act. NOAA Technical Memorandum NOAA-TM-NMFS-SWFSC-540.
- Bond, N.A., M.F. Cronin, H. Freeland and N. Mantua. 2014. Causes and impacts of the 2014 warm anomaly in the NE Pacific. *Geophysical Research letters*, Vol. 42, 3414-3420.
- Calambokidis, J. and J. Barlow. 2020. Updated abundance estimates for blue and humpback whales along the U.S. West Coast using data through 2018. NOAA Technical Memorandum NOAA-TM-NMFS-SWFSC-634. September 2020.
- Calambokidis, J., E.A. Falcone, T.J. Quinn, A.M. Burdin, P.J. Clapham, J.K.B. Ford, C.M. Gabriele, R. LeDu, D. Mattila, L. Rojas-Bracho, J.M. Straley, B.L. Taylor, J. Urban R., D. Weller, B.H. Witteveen, M. Yamaguchi, A. Bendlin, D. Camacho, K. Flynn, A. Havron, J. Huggins and N. Maloney. 2008. SPLASH: Structure of populations, levels of abundance and status of humpback whales in the North Pacific. Final Report for Contract AB133F-03-RP-00078, Cascadia Research, for U.S. Department of Commerce. May 2008.
- Cram, J., N. Kendall, A. Marshall, T. Buchrens, T. Seamons, B. Leland, K. Ryding and E. Neatherlin. 2018. Steelhead at risk report: Assessment of Washington's steelhead populations. 181 pp.
- Curtis, K.A., J. Calambokidis, K. Audley, M.G. Castaneda, J. De Weerd, A. J. Garcia Chavez, F. Garita, P. Martinez-Loustalot, J.D. palacios-Alfaro, B. Perez, E. Quintana-Rizzo, P.R. Barragan, N.

- Ransome, K. Rasmussen, J. Urban R., F.V. Zurita, K. Flynn, T. Cheeseman, J. Barlow, D. Steel and J. Moore. 2022. Abundance of humpback whales (*Megaptera novaeangliae*) wintering in Central America and southern Mexico from a one-dimensional spatial capture-recapture model. NOAA Technical Memorandum NOAA-TM-NMFS-SWFSC-661.
- Dahlheim, M.E., J.M. Waite, and P.A. White. 2009. Cetaceans of Southeast Alaska: Distribution and seasonal occurrence. Publications, agencies and staff of the U.S. Department of Commerce. Paper 157.
- Darling, J.D., K. Audley, T. Cheeseman, B. Goodwin, E.G. Lyman and R.J. Urban. 2022. Humpback whales (*Megaptera novaeangliae*) attend both Mexico and Hawaii breeding grounds in the same winter: mixing in the northeast Pacific. *Biology Letters*, 18.
- Fritz, L., K. Sweeney, R. Towell and T. Gelatt. 2016. Aerial and ship-based surveys of Stellar sea lions (*Eumetopias jubatus*) conducted in Alaska in June-July 2013 through 2015, and an update on the status and trend of the Western District population Segment in Alaska. NOAA Technical Memorandum NMFS-AFSC-321.
- Gabriele, C.M., C.L. Amundson, J.L. Neilson, J.M. Straley, C.S. Baker, S.L. Danielson. 2022. Sharp decline in humpback whale (*Megaptera novaeangliae*) survival and reproductive success in southeastern Alaska during and after the 2014-2016 Northeast Pacific marine heatwave. *Mammalian Biology* 102:1113-1131.
- Hester, K.C., E.T. Peltzer, W.J. Kirkwood and P.G. Brewer. 2008. Unanticipated consequences of ocean acidification: A noisier ocean at lower pH. *Geophysical Research Letters*, Vol. 35.
- Huff, D.D, S.T. Lindley, B.K. Wells and F. Chai. 2012. Green sturgeon distribution in the Pacific Ocean estimated from modeled oceanographic features and migration behavior. *PLOS ONE*, Vol. 7, Issue 9, September 2012.
- Jansen, J.K., P.L Boveng, J.M. Ver Hoef, S.P. Dahle, J.L. Bengtson. 2015. Natural and human effects on harbor seal abundance and spatial distribution in an Alaskan glacial fjord. *Marine Mammal Science*, 31(1): 66-89.
- Jemison, L.A., G.W. Pendelton, L.W. Fritz, K.K. Hastings, J.M. Maniscalco, A.W. Trites, T.S. Gelatt. 2013. Inter-population movements of Stellar sea lions in Alaska with implications for population separation. *PLOS ONE*. Vol. 8, Issue 8, August 2013.
- Kaplan, I.C., C.J. Brown, E.A. Fulton, I.A. Gray, J.C. Field and A.D.M. Smith. 2013. Impacts of depleted forage species in the California current. *Environmental Conservation* 40(4): 380-393.
- Koehn, L.E., T.E. Essington, K.N. Marshall, W.J. Sydeman, A.I. Szoboszlai and J.A. Thayer. 2017. Trade-offs between forage fish fisheries and their predators in the California current. *ICES Journal of Marine Science*, 74(9): 2448-2458.
- Lindley, S.T., M.L. Moser, D.L. Erickson, M. Belchick, D.W. Welch, E.L. Rechisky, J.T. Kelly, J. Heublein, and A.P. Klimley. 2008. Marine migration of North American Green Sturgeon. *Transactions of the American Fisheries Society*, 137:182-194.
- Martien, K.K., B.L. Taylor, F.I. Archer, K. Audley, J. Calambokidis, T. Cheeseman, J. De Weerd, A.F. Jordan, P. Martinez-Loustalot, C.D. Ortega-Ortiz, E.M. Patterson, M. Ransome, P. Ruvelas, J.U. Ramirez and F. Villagas-Zurita. 2021. Evaluation of Mexico distinct population segment of

- humpback whales as units under the Marine Mammal Protect Act. NOAA Technical Memorandum NOAA-TM-NMFS-SWFSC-658
- Mate, B.R. 1973. Population kinetics and related ecology of the Northern Sea Lion, *Eumetopias jubatus*, and the California Sea Lion, *Zalophus californianus*, along the Oregon Coast. Dissertation, University of Oregon. March 1973.
- McElhany, P., M.H. Tuckelshaus, M.J. Ford, T.C. Wainwright, and E.P. Bjorkstedt. 2000. Viable salmonid populations and the recovery of evolutionarily significant units. U.S. Department of Commerce, NOAA, Technical Memorandum NMFS-NWFSC042, 156 p.
- Mellinger, D.K., K.M. Stafford, and C.G. Fox. 2004. Seasonal occurrence of sperm whale (*Physeter macrocephalus*) sounds in the Gulf of Alaska, 1999-2001. *Marine Mammal Science*, 20(1): 48-62.
- Miller, S., T. Robertson, B. Higman, A. Chartier and S. Fletcher. 2019. Southeast Alaska vessel traffic risk analysis. Report to Alaska Department of Environmental Conservation, October 2019. Nuka Research and Planning Group, LLC.
- Mora, E.A., R.D. Battleson, S.T. Lindley, M.J. Thomas, R. Bellmer, L.J. Zarri, A.P. Klimley. 2018. Estimating the annual spawning run size and population size of the Southern Distinct Population Segment of Green sturgeon. *Transactions of the American Fisheries Society*, 147: 194-203.
- Muto, M.M. 2021a. Fin whale (*Balaenoptera physalus*): Northeast Pacific Stock. Alaska Marine Stock Assessments 2020. NOAA Technical Memorandum NOAA-TM-AFSC-421.
- Muto, M.M. 2021b. Sperm whale (*Physeter macrocephalus*): Northeast Pacific Stock. Alaska Marine Stock Assessments 2020. NOAA Technical Memorandum NOAA-TM-AFSC-421.
- National Marine Fisheries Service. Undated. North American green sturgeon brochure.
- National Marine Fisheries Service. 2024a. Humpback whale (*Megaptera novaeangliae*) fact sheet.
- National Marine Fisheries Service. 2024b. Green sturgeon (*Acipenser medirostris*) fact sheet.
- National Marine Fisheries Service. 2023a. Fin whale (*Balaenoptera physalus*) fact sheet.
- National Marine Fisheries Service. 2023b. Sperm whale (*Physeter macrocephalus*) fact sheet.
- National Marine Fisheries Service. 2023c. Steller sea lion (*Eumetopias jubatus*) fact sheet.
- National Marine Fisheries Service. 2022a. Recovery outline for the Central America, Mexico, and western North Pacific distinct population segments of humpback whales. ESA Recovery Outline.
- National Marine Fisheries Service. 2022b. Endangered Species Act status review report: sunflower sea start (*Pycnopodia helianthoides*). Final report to the National Marine Fisheries Service, Office of Protected Resources. October 2022.
- National Marine Fisheries Service. 2021. Occurrence of Endangered Species Act (ESA) listed humpback whales off Alaska. Revised August 6, 2021.
- National Marine Fisheries Service. 2013. Status review of the Eastern Distinct Population Segment of Steller sea lion (*Eumetopias jubatus*). Protect Resources Division, Alaska Region, National Marine Fisheries Service, Juneau, Alaska.

- National Marine Fisheries Service. 2010. Final recovery plan for the Fin whale (*Balaenoptera physalus*). National Marine Fisheries Service, Silver Spring, MD. 121 pp.
- National Marine Fisheries Service. 2008. Recovery plan for the Steller Sea Lion, Eastern and Western distinct population segments (*Eumetopias jubatus*). Revision. March 2008.
- National Marine Fisheries Service. 1991. Final recovery plan for the Humpback Whale, *Megaptera novaeangliae*. Prepared by the Humpback Whale recovery team. November 1991.
- National Oceanic and Atmospheric Administration. 2019. Alaska NOAA team examines dead endangered sperm whale. Accessed at: <https://www.fisheries.noaa.gov/feature-story/alaska-noaa-team-examines-dead-endangered-sperm-whale#> on July 8, 2024.
- National Oceanic and Atmospheric Administration Marine Debris Program. 2016. Report on marine debris impacts on coastal and benthic habitats. Silver Spring, MD.
- National Oceanic and Atmospheric Administration Marine Debris Program. 2014. Report on the entanglement of marine species in marine debris with an emphasis on species in the United States. Silver Springs, MD. 28 pp.
- National Park Service. 2021. Glacier Bay and Icy Strait humpback whale population monitoring: 2020 update. Informational brochure. Glacier Bay National Park & Preserve. July 2021.
- Neilson, J.L., C.M. Gabriele, A.S. Jensen, K. Jackson and J.M. Straley. Summary of reported whale-vessel collisions in Alaskan waters. Journal of Marine Biology, Vol. 2012.
- Neilson, J.L., J.M. Straley, C.M. Gabriele and S. Hills. 2009. Non-lethal entanglement of humpback whales (*Megaptera Novaeangliae*) in fishing gear in northern Southeast Alaska. Journal of Biogeography 36, 452-464.
- Northwest Fisheries Science Center. 2014. Status review update for Pacific salmon and steelhead listed under the Endangered Species Act: Pacific Northwest.
- NUKA Research and Planning Group. 2012. Southeast Alaska vessel traffic study. July, 2012, Revision 1.
- Peterson, W., N. Bond and M. Robert. 2016. The blob (part three): Going, going, gone? PICES Press Vol. 24, No. 1. North Pacific Marine Science Organization.
- Rone, B.K., A.N. Zerbini, A.B. Douglas, D.W. Weller and P.J. Clapham. 2016. Abundance and distribution of cetaceans in the Gulf of Alaska. Marine Biology, 164:23.
- Saez, L., D. Lawson and M. DeAngelis. 2021. Large whale entanglements off the U.S. West coast, from 1982-2017. NOAA Technical Memorandum NMFS-OPR-63A. March 2021.
- Santora, J.A., N.J. Mantua, I.D. Schroeder, J.C. Field, E.L. Hazen, S.J. Bograd, W.J. Sydeman, B.K. Wells, J. Calambokidis, L. Saez, D. Lawson and K.A. Forney. 2020. Habitat compression and ecosystem shifts as potential links between marine heatwave and record whale entanglements. Nature Communications, 11: 536.
- Straley, J., V. O'Connel, J. Liddle, A. Thode, L. Wild, L. Behnken, D. Falvey and C. Lunsford. 2015. Southeast Alaska sperm whale avoidance project (SEASWAP): a successful collaboration among scientists and industry to study depredation in Alaskan waters. ICES Journal of Marine Science, 72(5), 1598-1609.

- Straley, J.M., T.J. Quinn II and C.M. Gabriele. 2009. Assessment of mark-recapture models to estimate the abundance of a humpback whale feeding aggregation in Southeast Alaska. *Journal of Biogeography* 36, 427-438.
- Straley, J.M., C.M. Gabriele, S. Baker. 1994. Annual reproduction by individually identified humpback whales (*Megaptera novaeangliae*) in Alaskan Waters. *Marine Mammal Science* 10(1): 87-92.
- Sweeney, K. L., Birkemeier, B., Luxa, K., and Gelatt, T. 2023. Results of the Steller sea lion surveys in Alaska, June-July 2022. AFSC Processed Report 2023-02. Alaska Fisheries Science Center, NOAA, NMFS.
2022. AFSC Processed Rep. 2023-02, 32 p. Alaska Fish. Sci. Cent., NOAA, Natl. Mar. Fish. Serv., 7600 Sand Point Way NE, Seattle WA 98115.
- Sweeney, K., B. Birkemeier, K. Luxa and T. Gelatt. 2019. Results of Steller sea lion surveys in Alaska, June-July 2019. National Marine Fisheries Service, Memorandum to the record.
- Sweeney, K., L. Fritz, R. Towell and T. Gelatt. 2017. Results of Steller sea lion surveys in Alaska, June-July 2017. National Marine Fisheries Service, Memorandum to the record.
- Teerlink, S.F., O. von Ziegesar, J.M. Straley, T.J. Quinn II, C.O. Matkin, E.L. Saulitis. 2014. First time series of estimated humpback whale (*Megaptera novaeangliae*) abundance in Prince William Sound. *Environmental Ecological Statistics*, 22: 345-368.
- Titova, O.V., O.A. Filatova, I.D. Fedutin, E.N. Ovsyanikova, H. Okabe, N. Kobayashi, J.M.V. Acebes, A.M. Burdin and E. Hoyt. 2018. Photo-identification matches of humpback whales (*Megaptera novaeangliae*) from feeding areas in Russian Far East seas and breeding grounds in the North Pacific. *Marine Mammal Science*, 34(1): 100-112.
- Von Biela, V.R., M.L. Arimitsu, J.F. Piatt, B. Heflin, S.K. Schoen, J.L. Trowbridge, C.M. Clawson. 2019. Extreme reduction in nutritional value of a key forage fish during the pacific marine heatwave of 2014-2016. *Marine Ecology Progress Series*, 613: 171-182.
- Wade, P.R. 2021. Estimates of abundance and migratory destination for North Pacific humpback whales in both summer feeding areas and winter mating and calving areas. International Whaling Commission meeting paper.
- Webb, K.R. and S.M. Gende. 2015. Activity patterns and speeds of large cruise ships in Southeast Alaska. *Coastal Management*, 43: 67-83.
- Weingartner, T., L. Eisner, G.L. Eckert and S. Danielson. 2009. Southeast Alaska: oceanographic habitats and linkages. *Journal of Biogeography* 36, 387-400.
- Williams, T.H., B.C. Spence, D.A. Boughton, R.C. Johnson, L.G. Crozier, N.J. Mantua, M.R. O'Farrell, S.T. Lindley. 2016. Viability assessment for Pacific salmon and steelhead listed under the Endangered Species Act: Southwest. NOAA Technical Memorandum NOAA-TM-NMFS-SWFSC-564.
- Young, N.C., A. A. Brower, M. M. Muto, J. C. Freed, R. P. Angliss, N. A. Friday, P. L. Boveng, B. M. Brost, M. F. Cameron, J. L. Crance, S. P. Dahle, B. S. Fadely, M. C. Ferguson, K. T. Goetz, J. M. London, E. M. Oleson, R. R. Ream, E. L. Richmond, K. E. W. Shelden, K. L. Sweeney, R. G. Towell, P. R. Wade, J. M. Waite, and A. N. Zerbini. 2023. Alaska marine mammal stock assessments, 2022. NOAA Technical Memorandum NMFS-AFSC-474. July 2023.

- Zerbinin, A.N., J.M. Waite, J.L. Laake, and P.R. Wade. 2006. Abundance, trends and distribution of baleen whales off Western Alaska and the central Aleutian Islands. *Deep Sea Research* 1(53): 1772-1790

This page intentionally left blank.

Glossary

Critical habitat

Specific areas designated as critical by the Secretary of Interior or Commerce for the survival and recovery of species listed as Threatened or Endangered pursuant to the Endangered Species Act.

Endangered species

Any species of animal or plant that is in danger of extinction throughout all or a significant portion of its range. Plant or animal species identified and defined in accordance with the 1973 Endangered Species Act and published in the Federal Register.

Estuary

An ecological system at the mouth of a stream where fresh water and salt water mix, and where salt marshes and intertidal mudflats are present. The landward extent of an estuary is the limit of salt-intolerant vegetation, and the seaward extent is a stream's delta at mean low water.

Habitat

The sum total of environmental conditions of a specific place occupied by a wildlife or plant species or a population of each species.

Marine waters

Of, or belonging to, or caused by, the sea.

Threatened species

A plant or animal species likely to become an endangered species within the foreseeable future throughout all or a significant portion of its range. Threatened species are identified and defined in accordance with the 1973 Endangered Species Act and published in the Federal Register.