


## 5.1 Managing insect pests with buffers

Buffers can provide habitat for beneficial insects that prey on insect pests of crops. To encourage predation, both the life cycle of the beneficial predator and pest should be understood.

### Key design considerations

- Provide plant diversity and structure in the buffer.
- Protect buffer from disturbances (e.g., pesticides, tillage).
- Predation of insect pests generally increases with the percentage of buffer habitat in the area.
- Locate buffers throughout the fields and landscape to encourage dispersal of beneficial insects.
- Buffers may provide habitat for some pest insects but this can be reduced by selecting appropriate plants (see section 5.2).
- Beetle banks are long, planted berms that provide habitat for beneficial insects (see below).



5.1 Protection and Safety

## 5.1 References

- Altieri, M.A. 1983. Vegetational designs for insect-habitat management. *Environmental Management*. 7: 3-7.
- Altieri, M.A.; Nicholls, C.I. 2004. Effects of agroforestry systems on the ecology and management of insect pest populations. In: Gurr, G.M.; Wratten, S.D.; Altieri, M.A., eds. *Ecological engineering for pest management*. Ithaca, NY: Cornell University Press:143-154.
- Altieri, M.A.; Nicholls, C.I.; Fritz, M.A. 2005. Manage insects on your farm: a guide to ecological strategies. Beltsville, MD: Sustainable Agriculture Network. 130 p. <http://www.sare.org/publications/insect/insect.pdf>. [Date accessed: September 25, 2007].
- Altieri, M.A.; Todd, J.W. 1981. Some influences of vegetational diversity on insect communities of Georgia soybean fields. *Protection Ecology*. 3: 333-338.
- Altieri, M.A.; Whitcomb, W.H. 1979. The potential use of weeds in the manipulation of beneficial insects. *Horticultural Science*. 14: 12-18.
- Asteraki, E.J.; Hart, B.J.; Ings, T.C.; Manley, W.J. 2004. Factors influencing the plant and invertebrate diversity of arable field margins. *Agriculture, Ecosystems and Environment*. 102: 219-231.
- Bhar, R.; Fahrig, L. 1998. Local vs. landscape effects of woody field borders as barriers to crop pest movement. *Conservation Ecology*. 2(2): 3 <http://www.consecol.org/vol2/iss2/art3>. [Date accessed: August 14, 2007].
- Bianchi, F.J.J.A.; Booij, C.J.H.; Tscharntke, T. 2006 Sustainable pest regulation in agricultural landscapes: a review on landscape composition, biodiversity and natural pest control. *Proceedings of the Royal Society B*. 273: 1715-1727.
- Bianchi, F.J.J.A.; van der Werf, W. 2003. The effect of the area and configuration of hibernation sites on the control of aphids by *Coccinella septempunctata* in agricultural landscapes: a simulation study. *Environmental Entomology*. 32: 1290-1304.
- Bianchi, F.J.J.A.; van Wingerden, W.K.R.E.; Griffioen, A.J. [and others]. 2005. Landscape factors affecting the control of *Mamestra brassicae* by natural enemies in brussel sprout. *Agriculture, Ecosystem and Environment*. 107: 145-150.
- Collins, K.L.; Boatman, N.D.; Wilcox, A. [and others]. 2002. Influence of beetle banks on cereal aphid predation in winter wheat. *Agriculture, Ecosystems and Environment*. 93: 337-350.
- Collins, K.L.; Boatman, N.D.; Wilcox, A.; Holland, J.M. 2003. A 5-year comparison of overwintering polyphagous predator densities within a beetle bank and two conventional hedgebanks. *Annals of Applied Biology*. 143: 63-71.

- Corbett, A.; Rosenheim, J.A. 1996. Impact of natural enemy overwintering refuge and its interaction with the surrounding landscape. *Ecological Entomology*. 21: 155-164.
- Den Belder, E.; Elderson, J.; van den Brink, W.J.; Schelling, G. 2002. Effect of woodlots on thrips density in leek fields: a landscape analysis. *Agriculture, Ecosystems and Environment*. 91: 139-145.
- Dennis, P.; Fry, G.L.A. 1993. Field margins: can they enhance natural enemy population densities and general arthropod diversity on farmland. *Agriculture, Ecosystems and Environment*. 40: 95-115.
- Denys, C.; Tscharntke, T. 2002. Plant-insect communities and predator-prey ratios in field margin strips, adjacent crop fields, and fallows. *Oecologia*. 130: 315-324.
- Dix, M.E.; Johnson, R.J.; Harrell, M.O. [and others]. 1995. Influences of trees on abundance of natural enemies of insect pests: a review. *Agroforestry Systems*. 29: 303-311.
- Elliot, N.C.; Kieckhefer, R.W.; Michels, G.W., Jr.; Giles, K.L. 2002. Predator abundance in alfalfa fields in relation to aphids within-field vegetation, and landscape matrix. *Environmental Entomology*. 31: 253-260.
- Epila, J.S.O. 1988. Wind, crop pests and agroforest design. *Agricultural Systems*. 26: 99-110.
- Frank, S.D.; Shrewsbury, P.M. 2004. Effect of conservation strips on the abundance and distribution of natural enemies and predation of *Agrotis ipsilon* on golf course fairways. *Environmental Entomology*. 33: 1662-1672.
- Gurr, G.M.; Wratten, S.D.; Luna, J.M. 2003. Multi-function agricultural biodiversity: pest management and other benefits. *Basic and Applied Ecology*. 4: 107-116.
- Hawkins, B.A.; Cornell, H.V. 1994. Maximum parasitism rates and successful biological control. *Science*. 266:1886.
- Holland, J.; Fahrig, L. 2000. Effect of woody borders on insect density and diversity in crop fields: a landscape-scale analysis. *Agriculture, Ecosystems and Environment*. 78: 115-122.
- Jobin, E.; Choiniere, L.; Belanger, L. 2001. Bird use of three types of field margins in relation to intensive agriculture in Quebec, Canada. *Agriculture, Ecosystems and Environment*. 84: 131-143.
- Kemp, J.C.; Barrett, G.W. 1989. Spatial patterning: impact of uncultivated corridors on arthropod populations within soybean Agroecosystems. *Ecology*. 70: 114-128.

Kinross, C.; Wratten, S.D.; Gurr, G.M. 2004. Pest management and wildlife conservation: compatible goals for ecological engineering? In: Gurr, G.M.; Wratten, S.D.; Altieri, M.A., eds. Ecological engineering for pest management. Ithaca, NY: Cornell University Press: 199-218.

Kirk, D.A.; Evenden, M.D.; Mineau, P. 1996. Past and current attempts to evaluate the role of birds as predators of insect pests in temperate agriculture. In: Nolan, V., Jr.; Ketterson, E.D., eds. Current Ornithology 13. New York: Plenum Press: 175-269.

Kruess, A.; Tscharntke, T. 1994. Habitat fragmentation, species loss, and biological control. *Science*. 264: 1581-1584.

Landis, D.A.; Wratten, S.D.; Gurr, G.M. 2000. Habitat management to conserve natural enemies of arthropod pests in agriculture. *Annual Review of Entomology*. 45: 175-201.

Lee, J.C.; Menalled, F.D.; Landis, D.A. 2001. Refuge habitats modify impact of insecticide disturbance on carabid beetle communities. *Journal of Applied Ecology*. 38: 472-483.

MacLeod, A.; Wratten, S.D.; Sotherton, N.W.; Thomas, M.B. 2004. Beetle banks as refuges for beneficial arthropods in farmland. *Agricultural and Forest Entomology*. 6: 147-158.

Marino, P.; Landis, D.A. 1996. Effect of landscape structure on parasitoid diversity and parasitism in agroecosystems. *Ecological Applications*. 6: 276-284.

Menalled, F.D.; Costamagna, A.C.; Marino, P.C.; Landis, D.A. 2003. Temporal variation in the response of parasitoids to agricultural landscape structure. *Agriculture, Ecosystems and Environment*. 96: 29-35.

Menalled, F.D.; Marino, P.C.; Gage, S.H.; Landis, D.A. 1999. Does agricultural landscape structure affect parasitism and parasitoid diversity? *Ecological Applications*. 9: 634-641.

Nicholls, C. I.; Parrella, M.; Altieri, M. 2001. The effects of a vegetational corridor on the abundance and dispersal of insect biodiversity within a northern California organic vineyard. *Landscape Ecology*. 16: 133-146.

Nicholls, C.I.; Altieri, M.A. 2004. Agroecological bases of ecological engineering for pest management. In: Gurr, G.M.; Wratten, S.D.; Altieri, M.A., eds. Ecological engineering for pest management. Ithaca, NY: Cornell University Press: 33-54.

Norris, R.F.; Kogan, M. 2000. Interactions between weeds, arthropod pests and their natural enemies in managed ecosystems. *Weed Science*. 48: 94-158.

- Panzer, R. 2002. Compatibility of prescribed burning with the conservation of insects in small, isolated prairie reserves. *Conservation Biology*. 16: 1296-1307.
- Pasek, J. 1988. Influence of wind and windbreaks on local dispersal of insects. *Agriculture, Ecosystems and Environment*. 22/23: 539-554.
- Pfiffner, L.; Wyss, E. 2004. Use of sown wildflower strips to enhance natural enemies of agricultural pests. . In: Gurr, G.M.; Wratten, S.D.; Altieri, M.A., eds. *Ecological engineering for pest management*. Ithaca, NY: Cornell University Press: 165-186.
- Pickett, C.H.; Bugg, R.L. 1998. Enhancing biological control: habitat management to promote natural enemies of agricultural pests. Berkeley, CA: University of California Press. 421 p.
- Pywell, R.F.; James, K.L.; Herbert, I. [and others]. 2005. Determinants of overwintering habitat quality for beetles and spiders on arable farmland. *Biological Conservation*. 123: 79-90.
- Pywell, R.F.; Warman, E.A.; Hulmes, L. [and others]. 2006. Effectiveness of new agri-environment schemes in providing foraging resources for bumblebees in intensively farmed landscapes. *Biological Conservation*. 129: 192-206.
- Rodenhouse, N.L.; Barrett, G.W.; Zimmerman, D.W.; Kemp, J.C. 1992. Effects of uncultivated corridors on arthropod abundances and crop yields in soybean agroecosystems. *Agriculture, Ecosystems and Environment*. 38: 179-191.
- Russell, K.N.; Ikerd, H.; Droege, S. 2005. The potential conservation value of unmowed powerline strips for native bees. *Biological Conservation*. 124: 133-148.
- Saska, P.; Vodde, M.; Heijerman, T. [and others]. 2007. The significance of a grassy field boundary for the spatial distribution of carabids within two cereal fields. *Agriculture, Ecosystems and Environment*. 122: 427-434.
- Schimdt, M.H.; Thies, C.; Tscharntke, T. 2004. Landscape context of arthropod biological control In: Gurr, G.M.; Wratten, S.D.; Altieri, M.A., eds. *Ecological engineering for pest management*. Ithaca, NY: Cornell University Press: 55-63.
- Sootherton, N.W. 1995. Beetle banks – helping nature to control pests. *Pesticide Outlook*. 6: 13-17.
- Stamps, W.T.; Dailey, T.W.; Gruenhagen, N.M. 2007. Infestation of European corn borer, *Ostrinia nubilalis* in Midwestern USA fields with herbaceous borders. *Agriculture, Ecosystems and Environment*. 121: 430-434.

- Thies, C.; Steffan-Dewenter, I.; Tscharntke, T. 2003. Effects of landscape context on herbivory and parasitism at different spatial scales. *Oikos*. 101: 18-25.
- Thies, C.; Tscharntke, T. 1999. Landscape structure and biological control in agroecosystems. *Science*. 285: 893-895.
- Thomas, C.F.G.; Marshall, E.J.P. 1999. Arthropod abundance and diversity in differently vegetated margins of arable fields. *Agriculture, Ecosystems and Environment*. 72: 131-144.
- Thomas, M.B.; Wratten, S.D.; Sotherton, N.W. 1991. Creation of 'island' habitats in farmland to manipulate populations of beneficial arthropods: predator densities and emigration. *Journal of Applied Ecology*. 28: 906-917.
- Thomas, M.B.; Wratten, S.D.; Sotherton, N.W. 1992. Creation of 'island' habitats in farmland to manipulate populations of beneficial arthropods: predator densities and species composition. *Journal of Applied Ecology*. 29: 524-531.
- Thomas, S. 2000. Progress on beetle banks in UK arable farming. *Pesticide Outlook*. 11: 51-53.
- Thomas, S.; Noordhuis, R.; Holland, J.; Goulson, D. 2002. Botanical diversity of beetle banks: effects of age and comparison with conventional arable field margins in southern UK. *Agricultural, Ecosystems and Environment*. 93: 403-412.
- Tremblay, A.; Mineau, P.; Stewart, R.K. 2001. Effects of bird predation on some insect populations in corn. *Agriculture, Ecosystems and Environment*. 83: 143-152.
- Tscharntke, T.; Steffan-Dewenter, I.; Kruess, A.; Thies, C. 2002. Contribution of small habitat fragments to conservation of insect communities of grassland-cropland landscapes. *Ecological Applications*. 12: 354-363.
- Van Emden, H.F. 1965. The role of uncultivated land in the biology of crop pests and beneficial insects. *Scientific Horticulture*. 17: 121-136.
- Van Emden, H.F.; Williams, G.F. 1974. Insect stability and diversity in agro-ecosystems. *Annual Review of Entomology*. 19: 455-475.
- Varchola, J.M.; Dunn, J.P. 2001. Influence of hedgerow and grassy field borders on ground beetle activity in fields of corn. *Agriculture, Ecosystems and Environment*. 83: 153-163.
- Wilkinson, T.K.; Landis, D.A. 2005. Habitat diversification in biological control: the role of plant resources. In: Wackers, F.L.; van Rijn, P.C.J.; Bruin, J., eds. *Plant provided food for carnivorous insects*. Cambridge, MA: Cambridge University Press: 305-325.

With, K.A., Pavuk, D.M.; Worchuck, J.L. [and others]. 2002. Threshold effects of landscape structure on biological control in agroecosystems. *Ecological Applications*. 12: 52-65.

Woodcock, B.A.; Potts, S.G.; Pilgrim, E. [and others]. 2007. The potential of grass field margin management for enhancing beetle diversity in intensive livestock farms. *Journal of Applied Ecology*. 44: 60-69.

Wratten, S.D.; van Emden, H.F. 1995. Habitat management for enhanced activity of natural enemies of insect pests. In: Glen, D.M.; Greaves, M.P.; Anderson, H.M., eds. *Ecology and integrated farming systems*. Bristol, UK: John Wiley and Sons: 117-145.