### SPRUCE BUDWORM (CHORISTONEURA FUMIFERANA)

PERFORMANCE IN RELATION TO FOLIAR CHEMISTRY

OF ITS HOST PLANTS

William John Mattson

Principal Insect Ecologist U.S. Department of Agriculture, Forest Service North Central Forest Experiment Station 202 Pesticide Research Center Michigan State University East Lansing, MI 48824

Scott S. Slocum and C. Noah Koller Research Assistants University of Minnesota Department of Entomology St. Paul, MN 55108

Spruce budworm growth was best on balsam fir, poorest on lowland black spruce, and intermediate on upland white and black spruce. Growth was consistently, positively linked to foliar N and negatively linked to Fe, K, and select terpenes. Survival rates were not strongly, nor consistently linked to any of the measured foliar chemical traits.

The population dynamics of the spruce budworm, Choristoneura fumiferana, is clearly an ecosystem/ biome level process. The process has many parts, all of which interact to some degree, and none of which is really well understood individually, not to mention how they operate together. The process starts, as we see it, by a budworm larva taking a bite out of a plant. The plant has a complex array of nutritive and nonnutritive chemicals that together affect budworm behavior, metabolism and ultimately growth, reproduction and/or death. Our primary objective has been to examine variations in budworm performance among different host tree species in relation to the foliar chemistry of these trees. The ultimate goal is to identify the major host factors regulating budworm performance.

This paper reports on our preliminary analyses of the growth and survival of budworms in relation to total nitrogen, mineral elements, mono-terpenes, and phenolics in its diet on three tree species: balsam fir, Abies balsamea; white spruce, Picea glauca; and black spruce, Picea mariana.

This study was conducted in two locations, the main plot being about 20 miles south of International Falls, Minnesota, in Koochiching County, and the secondary plot about 15 miles west of Cloquet, Minnesota, in St. Louis County.

# Methods

In 1981, we selected 122 trees for the purpose of monitoring the performance of larval budworms in relation to foliar chemistry of these trees. Among this set were 20 white spruce, 25 black spruce, and 58 balsam fir which were divided into four classes: 24 small (1-5 m), 16 medium (5-10 m), 8 large (10-15 m), and 10 stress. Nineteen of the medium/ large category were selected at random for the purpose of studying both early and late season (i.e., late summer) budworm performance.

On each tree we selected five branches at midcrown level, approximately 70° apart so as to encircle the tree. Each branch was then enclosed with a 36" long fine-mesh cloth sleeve cage which served as an enclosure for 15 second-stage larvae placed therein. Larvae were obtained from the laboratory colony of the Insect Rearing Service of the Forest Pest Management Institute, Environment Canada, Canadian Forestry Service in Sault Ste. Marie, Ontario. Each branch contained at least 30 new shoots. Budworms were removed at the pupal stage and subsequent moths were collected in the laboratory and frozen within 24 hours after emergence in preparation for freeze-drying to constant weight.

We started the experiment in all cases after shoot elongation had begun (approximately 300 degree days, using 2.8°C as the threshold and March 1 as the beginning date). Budworms normally emerge from hibernacula at about 200 degree days in northern Minnesota (Bean and Wilson 1964) whereas in New Brunswick it may be closer to 100 (Cameron et al., 1968). Our records indicate that balsam "bud break" occurred at about 238 DD in our main plot. In 1981, we placed larvae on black spruce about 1 month later (6/22/81) than on balsam and white spruce so as to avoid any possible adverse effects of the late phenology of black spruce. In 1982 all tree species received larvae at 300 DD. Furthermore, in 1981, we also placed second stage larvae on 19 medium/large balsam on July 24. This is about the time that budworm first stage larvae are normally preparing for overwintering.

Foliage chemistry was measured only once during the larval period, at approximately the commencement of the fifth larval stage. This was done because the fifth and sixth stages consume 90-95 percent of the total diet (Miller 1977, Retnakaran 1983). Foliage was gathered from all sides of the tree at midcrown, immediately stored in coolers on dry ice and then frozen at the laboratory until used for analysis. Except for terpene analyses, the foliage was separated from stems, then lyophilized and ground to pass through a 40-mesh screen on a Wiley mill. It was then stored dry in a glass container in the dark until analysis. Details of the various analyses can be obtained from the authors on request.

Because the field studies did not permit control of temperatures during larval development, temperature effects may be confounded with tree effects when comparing budworm performance on different plots and at different times. In all cases we know the daily maximum, minima, and mean temperatures for our experiments as well as the precipitation values. Once we understand the budworm's temperature-growth responses we will be able to remove possible temperature effects from host effects.

Table 1.--Mean adult female and male weights (mg dwt) reared on different hosts in 1981 and 1982

Host	plant	19 Mean	81 SD	198 Mean	82 SD	198 Mean	31 SD	198 Mean	32 SD	"a/
		Fem	ales	Fema	ales	Mal	les	Ma	Les	
BFir	stress	21.44	5.64	22.13	5.11	10.24	1.92	12.91	2.95	10
BFir	lge	19.28	4.91	22.42	4.32	9.66	2.02	12.40	2.71	8
BFir	med	20.42	3.90	20.65	5.74	9.76	1.76	11.99	3.44	16
BFir	sm1	17.83	3.99	18.78	4.43	9.33	1.69	10.87	2.38	24
WSpr	1ge	16.81	5.50	18.49	6.33	7.31	1.93	12.43	3.02	10
WSpr	sml	15.00	6.05	16.17	6.21	8.20	2.81	9.26	2.93	10 . /
BSpr	upld	15.06	10.37	19.04	6.01	8.12	1.89	10.67	2.78	$10/20^{-c}$
BSpr	lowld	8.32	3.38	11.10	3.56	4.92	1.42	6.73	1.79	15
BFir	late	13.56	3.09	na <sup>b/</sup>	na	7.41	1.78	na	na	18

 $\frac{a}{b}$ Number of trees in each species class.  $\frac{b}{c}$ No data available. C Ten trees in 1981, twenty in 1982.

# Results

#### Differences Among Species

Budworms clearly attained the highest adult weights on balsam fir (18-22 mg dwt), followed by white spruce (15-18.5 mg) or upland black spruce (15-19 mg) and then lowland black spruce (8-11 mg) (Table 1). The pattern was consistent between years for both sexes. The fact that white spruce-reared budworms were smaller than those from fir is contrary to results from eastern North America where such spruce-reared budworms are usually about equal to (Blais 1957, Greenbank 1963) or significantly larger (Koller and Leonard 1982, and see T. Thomas in this proceedings) than those from balsam fir. The upland black spruce data agree with previous research which suggests that it produces slightly smaller insects than does balsam fir but not necessarily less than does white spruce (Blais 1957, Greenbank 1963).

Variability in female weights was about 50 percent greater on the spruces than on balsam fir as estimated by the coefficients of variation  $(SD/\tilde{x})$  (Table 1). This was the result of the fact that some spruce trees consistently produced very small insects ( $\leq 11$  mg) while others consistently produced large insects ( $\geq 20$  mg) similar to those on large balsam fir.

Generation survival rates (from second instars to the adult stage) in 1981 revealed that balsam fir was a superior host followed by white spruce and then black spruce (Table 2). The pattern changed in 1982, however, showing that white spruce and lowland black spruce were not different from balsam. Upland black spruce, however, again showed the lowest survival rates. We cannot explain differences in the survival patterns between years except to hypothesize that weather conditions may have been responsible. Mean daily temperatures were about  $2.4^{\circ}$ C higher during the larval period in 1981 than in 1982 (16.7° vs. 14.3°). Mean daily maxima averaged nearly the same (23.8° vs. 22.9°) in both years but daily minima were about 4°C higher in 1981 (9.5° vs. 5.7°). Table 2.--Survival rates and proportion of females from different hosts in 1981 and 1982

	Surviva	al rate	Propo fema	rtion les
Host plant	1981	1982	1981	1982
BFir stress	.290	•282	•48	.51
BFir md/lge	.305	.276	.55	.48
BFir sml	.330	.327	.54	.51
WSpr all	.232	.351	.53 ./	.51
BSpr upld	.130	.179	.71* <sup>a</sup> /	.45
BSpr lowld	.120	.27.5	•60*	.51
BFir late	.157	na	.53	na

 $\frac{a}{significantly}$  (p  $\leq$  .05) different from .50 ratio.

# Differences Among Age Classes

We studied budworm performance on three age/size classes of trees (3-5 m, 5-10 m, 10-15 m) known as small(s), medium(m), and large(1), respectively. All trees were nearly open grown so most had full tree crowns. Large trees were bearing male flowers in the upper half crown level, whereas only few of the medium and none of the small trees were flowering.

In both balsam and white spruce, there was a consistent trend for small trees to produce the smallest insects. This pattern held between years and for both sexes. For example, the female mean dry weights (mg) for each size class shown in the following tabulation are significantly different ( $p \leq .05$ ) from one another:

1981BFir: 20.35(m) 19.26(1) 17.83(s)1982BFir: 22.42(1) 20.65(m) 18.78(s)1981WSpruce: 16.81(1) 15.09(s)1982WSpruce: 18.49(1) 16.17(s)

The smallest difference between mean female weights on small and the large-medium classes ranged from 1.4 to 1.9 mg on balsam and 1.7 to 2.3 mg on white spruce. The differences imply that insects growing on large-medium trees averaged at least 10 percent larger than those from small trees and in some cases as much as 20 percent. Survival rates, however, did not vary significantly among age classes (Table 2). Therefore, although small balsam and white spruce produced smaller insects, they did not produce lower survival rates than larger trees. In general, for both balsam and white spruce, mean insect weight gains and survival rates per tree had no correlation with one another.

# Differences Owing to Phenology

On July 24, 1981, we "planted" second stage budworm larvae on 19 medium-large balsams. This date is nearly 2.5 months later than budworm second stage larvae normally emerge for feeding in northern Minnesota. The larvae on these trees attained only about two-thirds the size (13.56 mg) of normal early-season larvae (Table 1). Furthermore, survival was only about half (16%) of that experienced by the similar early-season cohort (Table 2). This was not unexpected because late summer foliage conditions are drastically different from those of early season. The fact that budworms performed as well as they did, however, was surprising. Mean daily temperatures during this late season experiment were about 2°C higher (18.9° vs. 16.7°) than during the early season experiments.

### Differences Owing to Stress Treatments

Ten trees (7 medium and 3 large) were trenched at a radius of 10 feet from the trunk in the spring of 1980 down to depth of about 30 inches. This depth reached well into the mineral soil layer or hit bedrock since the study plot was on top of a rock outcrop covered with shallow soils. The ground area under each tree canopy was then covered with black polyethylene plastic. Trees were monitored for moisture stress by pressure bombing twigs that were collected near the bottom of the crown. Pressure bomb readings were taken once per week and two times per day (before 8:00 a.m. and after 2:00 p.m.), during the 6-week larval growth period. The following tabulation shows the mean seasonal pressure bomb differences (stress-control) at both a.m. and p.m. samples:

	a.m. difference	p.m. difference
1981	0.50 bars	0.89
1982	0.45 bars	0.78

Mean differences in water potential between stress and control trees were usually less than one bar--suggesting that the stress treatment was relatively weak. The differences were nevertheless statistically significant on half of the sample periods each summer.

Stress trees produced significantly larger female and male insects than the control trees (all medium and large firs combined) (Table 1). However, in 1982 only males were significantly larger than those from the control trees. The mean weight differences (stress-control) for each sex are shown in the following tabulation:

		Females	Males	
1981	(Stress-Control):	1.31 mg	0.53	
1982	(Stress-Control):	0.79 mg	0.80	

Thus, so far, the induced stress treatment has had only a minor enhancing effect on budworm growth. However, it has not had any effect on survival rates (Table 2).

#### Budworm Performance in Relation to Foliar Nutrients

Theoretically an insect's performance should vary in relation to its dietary needs and a diet's deviations from these needs. In the case of the spruce budworm, no one really yet knows what the particular details of its needs are. Mattson and Koller (1981) proposed that the minimally optimal foliar N value for female performance is about 2.1 percent dwt. Harvey (1974) proposed that the minimally optimal levels of soluble sugars in the female's diet is about 4 percent fwt or 20 percent dwt. Albert et al, (1982) found that peak behavioral (i.e., feeding) response to soluble sugars (sucrose) in the diet occurs at about .03M or less than 1 percent fwt. Thus diets deviating negatively from these values probably will produce smaller insects and/or longer growth and feeding periods. Requirements for other important dietary components are really unknown.

To obtain a perspective on the budworm's needs for the mineral elements, we analyzed the nutrient levels in the bodies of adult males and females collected from the trees in our study and then compared their body levels to their food levels (Table 3). The data reveal that the elemental concentrations showing highest deviations from foliar levels are as follows for males and females:

# Females: Na>N>P>Cu>Zn>Fe>Mg>K>Ca>Mn Males: Na>N>Cu>P>Zn>Fe>Mg>K>Ca>Mn

Sodium clearly had the highest magnification factor (MF) (60.9 and 13.4), but owing to possible errors in its measurement on our emission spectrometer we consider the results very tentative. Nitrogen, P, and Cu were clearly the next highest, with all elements down to Fe showing MF's greater than two. All the rest were smaller than one. MF is concentration in insect body divided by concentration in foliage. Studying the budworm's utilization efficiencies of mineral elements in a low-salt McMorran (1965) meridic diet gave nearly the same results as the above magnification factor array:

# Cu>N>Zn>P>K>Mg>Fe>Ca>Mn

The utilization efficiencies array is somewhat different from MF array because the artificial diet does not exactly match the levels of elements contained in the foliage. The point is, though, that any variations in budworm performance on different hosts will most likely be due to variations in the limiting nutrients--those having highest magnification factors or utilization ratios.

Table 3.--Concentrations (ppm dwt) of mineral elements in the bodies of male and female spruce budworm adults, and balsam fir foliage (6/22/82), and the ratios of insect/foliage elemental concentrations

Item	%N	Р	K	Mg	Ca	Zn	Fe	Na	Cu	Mn
SBW Female Mean SD	7.97 .44	7,539 308	9,932 800	861 101	290 97	113 22	73 43	56 21	12 4	4 3
SBW <u>Male</u> Mean SD	9.51 .40	9,569 642	10,855 892	987 83	340 73	117 23	87 51	128 38	19 8	6 4
Fir foliage Mean SD	1.26	1,872 108	12,738 1,704	927 109	3,671 646	34 4	31 4	9 2	4 1	303 103
♀SBW/foliage	6.3	4.0	.78	.92	.08	3.3	2.4	6-1	3.3	.01
SBW/foliage	7.5	5.1	.85	1.06	•09	3.4	2.8	13.9	5.4	•02

Growth in Relation to N and Mineral Elements

Balsam fir. Regressing male and female dry weights against the foliar elemental concentrations they experienced as fifth and sixth stage larvae revealed that nitrogen was the only variable consistently and positively related to growth (Table 4). Calcium had a significant positive effect on female weights in 1981 but this is probably spurious owing to the fact that the larvae need only about 300 ppm calcium or less in their diets and foliage has about 15-fold this level. Calcium, of course, could be related to some other important foliar variable which in turn affects larval growth. For example, it was negatively correlated (p < .05) with both tannins and phenolics. Calcium may form chelates with many kinds of phenolics, perhaps rendering them less deleterious to a leaf consumer. The effects of the other elements (K, Fe, Cu) were, to our surprise, all negative. For example, based on the body/foliage magnification factors, we expected that Fe and Cu were in relative short supply, but their negative correlation with weight gain implies otherwise. However, as in the case of calcium, their correlation need not imply direct cause and effect but some indirect effect. K, for example, was significantly positively correlated with 10 mono- and sesqui-terpene species in balsam fir and with the terpene grand sums. Fe, on the other hand, was negatively correlated with four terpenes and Cu positively with three. In general, all foliar elements tended to show a negative correlation with total phenols and condensed tannins.

Late season results were unlike the early season results in that K and Cu were now positively correlated with weight gain (Table 4). For both elements, however, late season levels were less on the average than they were in early season (e.g., K: 8000 vs. 10,482). Not only were they less, they were much more variable (e.g., CV-K = .22 vs. CV-K = .08). Late season Cu levels were probably below optimal, for more than half of the late season trees had less than 0.5 ppm in their foliage. Early season trees, on the other hand, averaged about 4 ppm--none going below 2.6 ppm. When Cu occurred at similar levels (4 ppm) in the meridic diet, budworms more completely (ca. 60-75 percent) extracted it than other elements that occurred at levels comparable to early season foliage. N was not a significant late season nutrient variable for both sexes but only for males, probably owing to its relatively uniform concentration among late season trees (CV-N = .08 vs. CV-N = .14).

Table 4.--Significant variables in the regression of mean male and female adult dry weights per tree on foliar mineral element concentrations in different host trees and years

Host species	s Signi	fican	t Vari	ables	R <sup>2</sup>	n <sup>a/</sup>
Females						
BFir-79	+N	-Fe			.69	12
BFir-81	+N	-Fe	-к	+Ca	.44	50
BFir-late	+K				.17	18
Males						
BFir-79	+N	-Cu			.47	12
BFir-81	+N	-Fe	-к		.18	50
BFir-late	+N	-Fe	+Cu		.33	18
Females						
WSpr-81	+N	-к	-Zn		.61	18
BSpr-81	+N	-К			.66	17
Males						
WSpr-81					00	18
BSpr-81	+N	-Mn	-Cu		.76	17

 $\frac{a}{N}$  Number of trees in the regression.

Spruces. As with the firs, N was the only element showing a consistent, positive relation to weight gain for both sexes (Table 4). All other elements were negative; thereby corroborating the pattern seen for early season balsam fir. The consistently negative contribution of K stands out because K levels in black spruce were even lower in most cases than they were for late season balsam (6000 ppm vs. 8000 ppm). Thus one is obliged to conclude that K itself is not directly affecting weight gains but indirectly through its effect on some other plant traits. For example, plant K was positively associated with every terpene species but one in both black and white spruce. We hasten to add that none of the associations was statistically significant though. This is probably at least partially due to a small data set which will be enlarged in the next few months. As before, the negative contribution of Cu is difficult to explain because copper was close to its minimally optimal level in the foliage diet. On the other hand, manganese may be approaching deleterious levels in the foliage because it's about one-hundred fold more abundant there than in the insect's body. Moreover, excessively high levels of one mineral element can interfere with the absorption and utilization of other elements and nutrients (Maynard et al. 1979). No explanation is readily available for zinc's negative contribution because its level in the foliage (46 ppm) is hardly excessive. In fact, at this level in the artificial diet it is highly utilized and supports good growth.

Survival in Relation to N and Mineral Elements

Regressing generation survival (the arcsin transformation of the survival rate) against foliar elements revealed that none accounted for more than 50 percent of the observed variation and in most cases, they accounted for only about one-third of the variation (Table 5). Moreover, there was no consistency between years or between species. Therefore, we feel that the observed results may be entirely an artifact.

We were also surprised to learn that insect weight gains (FWT, MWT) and survival rates per tree were not significantly correlated with one another except on black spruce (\*) where the relationship was negative (p  $\leq$  .05) contrary to expectation:

	BFir	WSpr	BSpr	Late BFir
FWT	.11	.13	51*	•23
MWT	03	28	56*	.16

The latter result may suggest that foliage was in short supply and hence higher survival meant less food per insect and thus lower growth. On the other hand, it also could suggest that the tree traits governing survival and growth of budworm are linked in opposing directions, or perhaps not at all in the case of fir and white spruce. Table 5.--Significant variables in the regression of generation survival rates per tree (2d→adult) on foliar mineral elements and phenolics (PH) in different host trees and years

Host species	Sign	ifica	nt va	riables	R <sup>2</sup>	a/ n
BFir-79 BFir-81 BFir-late	+Fe +Mn +N	-PH +Cu			.32 .31 .18	39 50 18
WSpr-81 BSpr-81	+Cu +Mg	-к	-Fe	+РН	•52 •33	18 18

 $\frac{a}{N}$  Number of trees in regression

### Budworm Performance in Relation to Allelochemics

### Terpenes

Preliminary analyses reveal that several terpenes are significantly negatively correlated with weight gain for both sexes in both balsam fir and white spruce. In the case of balsam fir, the following six species of monoterpenes were negatively correlated ( $p \leq .05$ ) with growth except where noted (ns):

	alpha-	beta-		beta-
	pinene	pinene	camphene	phellandrene
MWT	36	37	43	38
FWT	32	32	40	25
	bornyl acetate	terpinol	terpe ene grd s	ene Sum
MWT	24	30	62	
FWT	16 ns	24	45	

None of the terpenes in balsam fir, in fact, were significantly positively correlated with growth.

In the case of the white spruce, five different compounds were significantly negatively correlated ( $p \leq .05$ ) with growth except where noted (ns) as shown in the following tabulation:

		Sesquite	erpenes &	monoterpene	alcohols	3
	camphor	#1	#2	#14	#35	
MWT FWT	33 ns 71	69 64	19 ns 66	32 ns 68	57 ns 65	

Only one terpene showed a positive relation to growth. The fact that white spruce and balsam may have different terpenes regulating budworm performance is not extraordinary because the two trees have different kinds as well as amounts of the individual terpenes. For example, balsam at mid-June had a terpene grand sum of about 6,700 ppm fwt vs. 836 ppm for white spruce. In other words, balsam has roughly 8-fold more terpenes. Black spruce was similarly terpene-rich having about 6,200 ppm fwt in mid-June. This suggests that if it's the total amount of terpenes that are

deterent, then budworm performance should be better on white spruce than on either balsam fir or black spruce, all other things being equal. The data, however, show the contrary. What is the explanation? Particular species-unique terpene compounds may be especially deterent at low levels in white spruce (e.g., camphor). On the other hand, all other things are not equal. For example, lower plant nitrogen levels may make white spruce significantly less suitable than fir in spite of its lower terpene levels. White spruce, for example, had 20 percent less N in mid-June than did balsam fir (e.g., 1981: 1.24 percent vs. 1.52 percent). We know that this is due to the fact that white spruce grows faster than balsam fir and thereby dilutes its foliar nutrient levels similarly faster (Fig. 1).

Going back to the hypothesis of the effects of terpene grand sums on budworm growth leads us to differences between small and medium/large balsams. As we said earlier, small trees produced 10 to 20 percent smaller insects than did medium/large trees. This may have been due to the fact that small trees had 42 percent more terpenes than did medium/large trees at mid-June when larvae were in the 5th/6th instars (9,547 ppm fwt vs. 6,700 ppm). The N levels in these two age classes were nearly identical (e.g., 1981: 1.50 vs. 1.55), so this is not a large potential source of variation. We also know that mineral element differences are not a likely explanation of the difference because they appeared to be available in sufficient amounts in both age classes, barring any negative interactions with tannins and phenolics. Similarly, there were



Figure 1.--Seasonal change in foliar nitrogen content (percent dwt) of three balsam and three white spruce with respect to degree day accumulations (2.8°C as base), and the phenology of the spruce budworm

no significant differences between the two groups for total phenols (2.64 percent vs. 2.59) or tannins (3.8 vs. 4.5 percent). Therefore, until the final analyses are complete, we suggest that the budworm growth difference between balsam age classes is due in large part to their terpene differences.

Small white spruce also had smaller insects and more total terpenes than did large white spruce (994 ppm fwt vs. 667 ppm). But, in this case, male and female weights were not correlated with terpene grand sums. The most likely explanation for differences in insect weight gain between white spruce age classes is the difference in levels of the one or more of individual terpene compounds listed above. There were no significant differences in tannins, total phenolics, and N levels between the two age classes just as was the case for balsam.

#### Phenolics-tannins

Incorporating phenolic and tannin estimates along with the mineral element data in the multiple regressions did not change any earlier conclusions for balsam fir but did in the case of male weight gain on white spruce. In this case, tannin was the only variable significantly correlated with male weight (r = -.46). This reflects the fact that no other variables were previously related to male weight gains.

The simple correlation coefficients between phenolics, tannins and weight gains on fir and spruce were not significant in all but two cases (\*) as shown below:

	Bal	sam	White Spruce		
	FWT	MWT	FWT	MWT	
Phenolics	29*	.00	.19	.02	
Tannins	06	.13	20	46*	

In general, levels of tannins and phenolics were negatively correlated with nearly all mineral elements in fir and white spruce. In the case of spruce, however, only one of the correlations was significant. This is in striking contrast to balsam where only five out of eighteen negative correlations were not significant in both 1979 and 1981, 2 years for which we have substantial data sets. The following tabulation shows the significant ( $p \leq .05$ ) tannin and phenolic correlations with N in balsam fir foliage:

	N-79	N-81
Phenolics	62	45
Tannins	na	42

Palmer (1982) found the same negative relationship between phenolics and N in <u>Populus tremuloides</u> foliage.

Previous studies on the effects of phenolics and tannins on the growth performance (via lowered protein utilization) of folivorous insects have almost unanimously come to the same general conclusion: there seems to be little or no measurable deleterious effect, except perhaps at extraordinarily high tannin concentrations (Lawson et al. 1982, Bernays 1981, Fox and McCauley 1977).

While our analyses are too preliminary to come to any firm conclusions, we believe that there may also be long-term, subtle effects that have been overlooked in the typical growth bioassays which usually measure insect performance (nutritionally and behaviorally) for only one or two instars, or at most for the whole feeding period of one generation. Tannins and/or phenols could have some chronic, subtle effects, such as through the chelation of micro-nutrients like Cu, Zn, and Fe--which then renders them less available to the insects. Such effects would have to be monitored over two or more successive generations to see the impacts of chronic micronutrient deficiencies. The chelation of such micronutrients may also be an important plant defense against microorganisms and could also thereby affect essential gut microbes or pathogens in those consumers having such microsymbionts (Swinburne 1981, Radhakrishnan and Sivaprasod 1980, Roy and Mukherjee 1979, Shieh <u>et</u> al. 1968, Emery 1982).

# Budworm Growth Response to N Levels

The analyses have indicated that budworm weight gains are consistently but not exclusively linked to variations in plant foliar N levels. In those cases where N was not implicated, there was usually only very little variation in N levels among plants. Therefore, to add strength to the N hypothesis, we planned two experiments to raise the level of nitrogen offered to larvae, thereby hoping to elicit increased growth. The first experiment was designed to raise dietary N for just the first three instars (2, 3, and 4), whereas the second was designed to raise N for the whole larval life span, but particularly for late stage (5, 6) larvae.

# Diet/Foliage Transfer Experiment

In the early spring before budbreak occurs, second stage budworm larvae mine 1-year-old or sometimes 2-year-old needles for 1 to 2 weeks (Blais 1979, McGugan 1954). After molting to the third instar, larvae then move to the newly opening buds or male flowers for feeding. During the needle mining period budworm subsist on very little N, such needles, usually having between 0.95 and 1.1 percent N. The newly flushed foliage, on the other hand, is very N rich (as high as 8 percent) but very emphemeral because shoots and needles are rapidly expanding thereby causing a precipitous decline in most foliar elements (Fig. 1), probably owing to a dilution effect. The decline takes the form of a negative exponential (%N = aX)) as has also been reported by Shaw and Little (1977). By the fifth larval stage N levels are < 1.5 percent in balsam, and somewhat less in white spruce owing to its faster development and nutrient dilution (Fig. 1). Since the fifth and sixth larval stages eat about 95 percent of the whole food budget for the larval period, (Retnakaran 1983, Miller 1977) we asked whether a consistently high N diet for the first three feeding stages (2, 3, 4) would have any enhancing effect on ultimate weight gain if these larvae were then transferred at the 5th instar to a low N foliage diet. To test this hypothesis we simultaneously grew insects on meridic diet (McMorran 1965) having about 4.4 percent N and also on the natural hosts, balsam and white spruce. At the fifth larval stage, ten females were

transferred from the diet to each of 48 balsam fir and 20 white spruce trees for completion of feeding. At pupation all insects were removed and later weighed for adult body size. The results on balsam showed that diet/foliage reared larvae were significantly larger (9 percent) than foliage/foliage reared insects (20.51 vs. 18.87 mg). Furthermore, plotting mean female weights of each group from a tree against the foliar N values they experienced as fifth and sixth stage larvae gave nearly identical regressions except for the intercepts which reflect their mean differences in body size:

diet/foliage	FWT =	12.06 +	5.42%N	$r_{2}^{2} =$	.19
foliage/foliage	FWT =	10.37 +	5.46%N	$r^2 =$	.22

Thus the early diet ration seems to have "bumped up" final adult weights.

On the other hand, the same experiment on white spruce gave opposite results; diet/foliage insects were about 9 percent smaller (14.18 vs. 15.50 mg where .1 ). There was no significant regression of either group on foliar N values but the two groups nevertheless performed similarly on the set of 20 spruce as evidenced by the significant regression between their respective mean adult weights:

$$FWT (d/f) = 6.72 + .481 FWT (f/f) r^2 = .30$$

These results are in direct contrast with those reported by Thomas (this proceedings) who found that diet to current white spruce foliage transfers (at the 6th larval stage) produced larger insects (ca. 21 mg) than similar transfers to balsam fir (ca. 15 mg). The explanation for this divergence from our results could be due to several factors: different populations of insects, different physical environments, and different host materials. The latter one entails at least two variables. Thomas used excised branches of both fir and spruce whereas we used intact branches. Perhaps even more important is the fact that the eastern populations of white spruce and fir are known to differ (phytochemically) from the more western populations such as the one we studied in northern Minnesota (Wilkinson et al. 1971, von Rudloff 1975, Zavarin and Snajbeck 1972, and Lester 1974). For example, von Rudloff (1975) reported that camphor levels are nearly twice as high in some western populations of white spruce than in eastern ones. However, until more is also known about the nutrient levels of the eastern host materials, it's impossible to explain the causes of the apparent differences in budworm performance.

To recapitulate, the balsam fir transfer experiment suggested that giving young larvae a high protein diet enhanced their performance by an amount that was constant, when the N effect on late larvae was accounted for. On the other hand, the spruce transfer experiment suggested that the early high protein diet was not enhancing, for these larvae attained smaller size than those having spent their early instars on spruce. Nevertheless both diet and tree insect groups performed similarly with respect to the trees they were on. The following tabulation summarizes the transfer effects on adult female dry weights relative to the meridic diet control:

Maan famala	diet/wsp	wsp/wsp	bfir/bfir	
weight (mg)	14.18	15.50	18.87	
Maan formala	diet/b	fir diet.	/diet_	
weight (mg)	20.5	1 26-3	31	

Obviously, feeding for the fifth and sixth larval stage on a tree was not as good as feeding on the diet. White spruce insects achieved about half, and balsam insects about two-thirds of their potential size. Why the meridic diet experience enhanced growth on fir but disenhanced it on spruce is an enigma. Perhaps early season spruce foliage is superior to the artificial diet. On the other hand, later in the season, it is plainly inferior. One can speculate that there must have been significant transfer shock going from diet to trees, especially in the case of white spruce. Perhaps the shock was due to higher metabolic costs to operate the budworm's mixed-function oxidase system on spruce than on fir (Brattsden 1983). This assumes, of course, that there were allelochemics that elicited higher MFO activity on spruce than on fir or that there were present significant MFO inhibitors in the spruce foliage which may have rendered spruce allelochemics more deleterious. On the other hand, the effect may have occurred at the behavioral level, for spruce foliage seems to harden off faster than balsam, and it might also possess important feeding deterrents (e.g., pungenin) (Heron 1965).

# Fertilization Study

To elucidate the budworm growth response ( $\Delta$ WT) to changes in foliar N ( $\Delta$ N), we first regressed budworm weight gains in 1981 on foliar N levels. Secondly, we fertilized our most N-impoverished trees, the lowland black spruce (3-5 m tall and 30-45 years old) and half of the small balsam which are comparable to trees used in two earlier fertilization studies (Shaw et <u>al</u>. 1978, Shaw and Little 1972). In each case we had 15 treated and 15 control trees. We applied 600 lbs. N/acre (urea) around the root zone of each tree during the first week of May 1982.

Male and female weight gains (MWT, FWT) in 1981 were clearly linear functions of foliar N (using all tree species and age classes):

FWT	in	3.45	+	9.844	(%N)	$r_2^2$	=	•54	n	=	114
MWT	=	3.13	+	3.959	(%N)	$r^2$	=	•44	n	20	113.

Scatter plots of the data revealed that female weight gains showed a clear positive trend over the full range of N values (0.47-2.05%). Furthermore, female weights increased with even higher N levels (2-4%) administered in the form of casein and wheatgerm in artificial diets. Male weights, on the other hand, showed little tendency to increase with foliar N levels above 1.5\%. Moreover, administering even higher levels of N (2-4%) in the artificial diet brought about a weak response, 1-2 mg, suggesting that above 1.5\% N males have a shallow response potential if any. The linear regressions imply that the w ~: gain response of budworms to an increment of N is constant, e.g.,  $\Delta FWT/\Delta N = 9.84$ , the slope of the regression line. In other words, for each unit increase in foliar N there is a concomitant 9.8 mg increase in female weight. In the case of the fertilization study, our treatment of 600 lbs. N/acre elevated foliar N levels by 0.40 and 0.45% for the black spruce and balsam fir trees, respectively. These N changes both elicited 1.38 mg changes in female mean weights as shown in the in the following tabulation:

	FWT AFWT	<u>MWT</u> <u>A MWT</u>	<u>% N</u> <u>∆ %N</u>
BfirCk	18.78	10.87	1.31
BfirF	20.16	11.24	1.76
BSprCk	11.10	6.73	0.85
BSprF	12.48	7.69	1.25

The urea treatment caused significant (p  $\leq$  .05) increases in female weights on both tree species but significant increases in male weights only on black spruce. The fact that the males did not respond to urea fertilization on balsam lends support to our suspicion that males have a low optimal dietary N requirement (perhaps about 1.5%) and that near this level their response is nearly flat.

Using the fertilization data for females only, we calculate that  $\Delta$  FWT/ $\Delta$ N for balsam and black spruce are 3.07 and 3.45, respectively. These values are not much different from those derived from the fertilization experiments of Shaw et al. (1978). They reported that high and low urea treatments raised foliar N values (on June 23) by 0.8 and 0.4%, respectively. These increments in turn elicited female adult dry weight gains (using formula of Mattson et al. 1982) of 3.33 and 1.13 mg. Thus  $\Delta FWT/\Delta N$  was 4.16 and 2.83 for the high and low urea treatments, respectively. Shaw et al.'s high calcium nitrate treatment gave a △FWT/△N value of 2.86. Therefore, pooling Shaw et al.'s and our values suggests that  $\Delta$  FWT/ $\Delta$ N averages about 3.27 and ranges from 2.83-4.16. In other words, FWT = a + 3.27 (%N). This implies that for every 1% increment in foliar N, there will be a corresponding 3.27 mg dwt increment in adult female weights.

On the other hand, our earlier regression analysis suggested that AFWT/AN should be about 9 instead of 3. What's the explanation for this discrepancy? The explanation might lie in the fact that changing foliar N through fertilization results in many other changes in foliar chemistry that are not all enhancing. For example, fertilization is also known to raise the levels of mono- and sesqui-terpenes. Moreover, the fertilization studies reported herein (Shaw et al. and ours) were done on young balsam fir which we have already shown to have significantly higher terpene levels than older trees. Thus the budworm responses ( $\Delta$ FWT) on small trees might be significantly less than on older trees which have lower levels of terpenes. Similarly, the lowland black spruce which we fertilized probably has

levels of terpenes at least as high and total phenolics levels that are higher than the small fir. Thus the overall 1981 regression may have a higher slope or predicted weight increment per unit of N because the pooled data consists of such species as white spruce and medium/large balsam which may have higher levels of N relative to terpenes, thereby giving higher weight gains per unit N increment than would small balsams and lowland black spruce.

### Changes in Fecundity

In order to obtain some idea of the potential impact of different dietary regimes on budworm population dynamics, we used the fecundity/pupal size equation of Miller (1963) and the adult dry weight/pupal size equation of Mattson <u>et al</u>. (1982) to project changes in female dry weight into changes in egg output. The resulting formula for fecundity in relation to body size is as follows:

$$F = -442.1 + 216.7 (FWT)^{.37}$$
  
 $\Delta F = 81.04 (FWT^{.63}) \Delta FWT$ 

The second equation says that changes in egg output  $(\Delta F)$  increase directly with changes in body size  $(\Delta FWT)$ . In other words, a 2 mg change in body size elicits exactly twice the output of a 1 mg change and so on, holding initial body size (FWT) constant.

The question yet to be answered is how large are the differences in fecundity between insects having different sizes. For example, female budworms from small balsam averaged 32 less eggs than similar females on medium balsam (187 vs. 219) (Table 6). Similarly insects from small white spruce averaged 26 less eggs than females from large white spruce (148 vs. 174). In the case of black spruce, females from the lowland trees averaged 117 fewer eggs than those from upland trees (32 vs. 149).

In the case of the fertilization experiments, we estimated that increasing foliar nitrogen levels by 1.0% would result in roughly a 3 mg increment in female weight. This translates into 41 more eggs for females that weighed 17 mg before fertilization. If foliar N increased only 0.5%, the result would be about half as many eggs, i.e., 20 more per female.

The significance such differences in egg output have on the insect's population dynamics cannot, of course, be answered. These are questions that must be addressed through an ecosystem level model which incorporates all of the major factors regulating budworm natality and mortality.

#### CONCLUSIONS

Although the study is not yet completed, there are some consistencies that seem substantial enough to warrant recapitulation.

There are clear differences in budworm growth between small and large tree classes, larger trees producing larger and hence more fecund insects. Furthermore, there are differences between species--balsam giving rise to larger insects than white spruce. This pattern seems not to hold in eastern North America where the reverse is true. The explanation for this inconsistency may reside in the fact that the phytochemistry of the eastern and western tree populations are different owing to limited gene exchange, different geological histories, and different environments. There is also, of course, the possibility that the insect populations are substantially different as well.

Budworm survival rates did not vary among tree size/age classes and appeared to be highest on balsam and lowest on black spruce. Survival rates and budworm weight gain per tree were not correlated except in the case of black spruce where the association was negative. This implies that the plant traits which affect weight gain are independent of those that affect survival except perhaps in black spruce where they could be negatively linked. Budworm survival rates were not consistently linked to any variables that were measured.

Insect weight gains per tree were consistently. positively linked to foliar N and negatively to Fe and/or K. The negative associations with Fe and K are surprising because neither element occurs at levels high enough to be minimally optimal much less toxic or noxious. Nevertheless, iron levels in the insect body show a tendency to decrease with increasing levels in the diet. Iron, in fact, is the only element showing this inverse behavior. Moreover, iron concentrations in the insect are also negatively correlated with insect size. What this implies is not clear. It may mean that insects sequester less Fe per unit body weight when diets are better for growth. Fe concentration is also positively correlated with (1/N) of the diet. Since total consumption is usually positively linked to (1/N), Fe uptake by the insect may be related to the total amount of Fe passing through the digestive system.

Potassium levels by themselves may not be inhibitory to budworm growth but K is linked positively to foliar terpene levels which apparently are inhibitory.

Terpene levels were negatively linked to budworm weight gains both on balsam and on white spruce. In the case of balsam, terpene grand sums (all molecular species pooled) showed the strongest correlations whereas in white spruce individual compounds not the grand sums had the highest linkage to weight gain. Difference in budworm performance between tree age classes may be largely due to higher levels of terpenes in the younger trees. Because plant terpene profiles are known to exhibit significant east-west variation across North America, it is likely that budworm performance on these trees will similarly vary.

The significance of phenolics and tannins in the performance of spruce budworm is still uncertain. The early data suggest little effect on either survival or weight gain. However, the spectrum of phenolic compounds in balsam and the two spruces has not yet been examined so it is entirely possible that one or more of them could have significant behavioral or physiological effects. Pungenin, for example, in white spruce could be an important feeding deterent.

Finally, it is clear that changing insect N intake through artificial diet to foliage transfers and fertilization results in enhanced insect growth, at least on balsam fir. The data suggest that growth increment per unit nitrogen increment ( $\Delta$ FWT/ $\Delta$ N) is about 3, as long as the minimally optimal level of N in the diet has not yet been reached. We suspect that the growth increment per unit nitrogen increment ( $\Delta$ FWT/ $\Delta$ N) will vary with different host species and age classes owing to different background levels of nutrients and allelochemics.

### Literature Citations

- Albert, P. J.; Cearley, C.; Hanson, F.; Parisella, S. Feeding responses of eastern spruce budworm larvae to sucrose and other carbohydrates. J. Chem. Ecol. 8:: 233-239; 1982.
- Bean, J. L.; Wilson, L. F. Comparing various methods of predicting development of the spruce budworm, <u>Choristoneura fumiferana</u>, in northern Minnesota. J. Econ. Ent. 57: 925-928; 1964.
- Bernays, E. Plant tannins and insect herbivores: an appraisal. Ecol. Entomol. 6: 353-360; 1981.
- Blais, J. R. Some relationships of the spruce budworm, <u>Choristoneura fumiferana</u> (Clemens) to black spruce, <u>Picea mariana</u> (Moench) Voss. For. Chron. 13: 364-372; 1957.
- Blais, J. R. Rate of defoliation in relation to spruce budworm attack and timing of spray application. Can. J. For. Res. 9: 354-361; 1979.
- Brattsden, L. B. Cytochrome P-450 involvement in the interaction between plant terpenes and insect herbivores. In: ACS Symposium Series 208, Plant Resistance to Insects, P. A. Hedin, ed., Washington, D.C. 1983: 173-195.
- Cameron, D. G.; McDougall, G. A.; Bennett, C. W. Relation of spruce budworm development and balsam fir shoot growth to heat units. J. Econ. Ent. 61: 857-858; 1968.
- Emery, T. Iron metabolism in humans and plants. Amer. Scientist 70: 626-632; 1982.
- Fox, L. R.; Macauley, B. J. Insect grazing on Eucalyptus in response to variations in leaf tannins and nitrogen. Oecologia 29: 145-62; 1977.
- Harvey, G. T. Nutritional studies of eastern spruce budworm (Lepidoptera: Tortricidae) I. soluble sugars. Can. Ent. 106: 353-365; 1974.

- Heron, R. J. The role of chemotactic stimuli in the feeding behavior of spruce budworm larvae on white spruce. Can. J. Zool. 43: 247-269; 1965.
- Koller, C. N.; Leonard, D. E. Comparison of energy budgets for spruce budworm, <u>Choristoneura</u> <u>fumiferana</u> (Clemens) on balsam fir and white spruce. Oecologia 49: 14-20; 1981.
- Lawson, D. L.; Merritt, R. W.; Klug, M. J.; Martin, J. S. The utilization of late season foliage by the orange striped oakworm, <u>Anisota senatoria</u>. Ent. exp.& appl. 32: 242-248; 1982.
- Lester, D. T. Geographic variation in leaf and twig monoterpenes of balsam fir. Can. J. For. Res. 4: 55-60; 1974.
- Maynard, L. A.; Loosli, J. K.; Hintz, H. F.; Warner, R. G. Animal nutrition. New York: McGraw-Hill; 1979; 602 p.
- McMorran, A. A synthetic diet for the spruce budworm, <u>Choristoneura fumiferana</u> (Clemens) (Lepidoptera: Tortricidae). Can. Ent. 97: 58-62; 1965.
- Mattson, W. J.; Koller, C. N. Spruce budworm performance in relation to matching selected chemical traits of its hosts. In: Proc. IUFRO Symposium, The Role of the Insect/Plant Relationship in the Population Dynamics of Forest Pests, Irkutsk, U.S.S.R. Aug. 24-28, 1981. In press.
- Mattson, W. J.; Koller, C. N.; Slocum, S. S. Adult female spruce budworm, <u>Choristoneura</u> <u>fumiferana</u>, (Lepidoptera: Tortricidae), dry weight in relationship to pupal fresh weight and case diameter. Great Lakes Entomol. 15: 185-187; 1982.
- Mattson, W. J.; Lorimer, N.; Leary, R. A. Role of plant variability (trait vector dynamics and diversity) in plant/herbivore interactions. In: Resistance to diseases and pests in forest trees. H. M. Heybroek, R. B. Stephan, and K. von Weissenberg, eds., Pudoc. Wageningen, Netherlands. 1982: 295-303.
- Miller, C. A. The analysis of fecundity in the unsprayed area. In: The dynamics of epidemic spruce budworm populations. R. F. Morris, ed. Mem. Entomol. Soc. Canada No. 31: 75-87; 1963.
- Miller, C. A. The feeding impact of spruce budworm on balsam fir. Can. J. For. Res. 7: 76-84; 1977.
- Palmer, S. R. Natural and induced variation in the nutrient and phenolic content of trembling aspen foliage. St. Paul: University of Minnesota; 1982. 50 p. Thesis.
- Radhakrishnan, M. R.; Sivaprasad, J. Tannin content of sorghum varieties and their role in iron bioavailability. J. Agric. Food Chem. 28: 55-57; 1980.

- Retnakaran, A. Spectrophotometric determination of larval ingestion rates in the spruce budworm (Lepidoptera: Tortricidae). Can. Ent 115: 31-40; 1983.
- Roy, S. N.; Mukherjee, S. Influence of food tannins on certain aspects of iron metabolism and excretion in normal and anemic rats. Indian J. of Biochem. and Biophys. 16: 93-98; 1979.
- Shaw, G. G.; Little, C. H. A. Effect of high urea fertilization of balsam fir trees on spruce budworm development. In: Insect and Mite Nutrition, J. D. Rodriguez, ed., North-Holland Publishing Co., Amsterdam. 1972: 589-597.
- Shaw, G. G.; Little, C. H. A. Natural variation in balsam fir foliar components of dietary importance to spruce budworm. Can. J. For. Res. 7: 47-53; 1977.
- Shaw, G. G.; Little, C. H. A.; Durzan, D. J. Effect of fertilization of balsam fir trees on spruce budworm nutrition and development. Can. J. For. Res. 8: 364-374; 1978.
- Shieh, T. R.; Mathews, E.; Wodzinski, R. J.; Ware, J. H. Effect of calcium and phosphate ions on the formation of soluble iron-gossypol complex. J.Agr. Food Chem. 16: 208-211; 1968.
- Singleton, V. L.; Kratzer, F. H. Toxicity and related physiological activity of phenolic substances of plant origin. J. Agr. Food Chem. 17: 497-511; 1969.
- Swinburne, T. R. Iron and iron chelating agents as factors in germination, infection, and aggression of fungal pathogens. In: Microbial Ecology of the Phylloplane, J. P. Blakeman, ed., Academic Press, NY. 1981: 227-243.
- von Rudloff, E. Chemosystematic studies in the genus <u>Picea</u> (Pinaceae). II. The leaf oil of <u>Picea galuca</u> and P. <u>mariana</u>. Can. J. Bot. 45: 1703-14; 1967.
- von Rudloff, E. Volatile leaf oil analysis in chemosystematic studies of North American conifers. Biochemical Syst. and Ecol. 2: 131-167; 1975.
- Wilkinson, R. C.; Hanover, J. W.; Wright, J. W.; Flake, R. H. Genetic variation in the monoterpene composition of white spruce. Forest Sci. 17: 83-90; 1971.
- Zavarin, E.; Snajberk, K. Geographical variability of monoterpenes from <u>Abies</u> balsamea and A. <u>fraseri</u>. Phytochemistry 11: 1407-1421; 1972.
- McGugan B. M. Needle mining habits of the Spruce budworm. Can. Ent. 86: 439-454; 1954.

65