



Review

Northern hardwood silviculture at a crossroads: Sustaining a valuable resource under future change

Nicole S. Rogers^{a,*}, Anthony W. D'Amato^b, Christel C. Kern^c, Steve Bédard^d^a School of Forest Resources, University of Maine, Orono, ME 04473, USA^b Rubenstein School of Environment and Natural Resources, University of Vermont, Burlington, VT 05405, USA^c USDA Forest Service, Northern Research Station, 1831 Hwy 169 East, Grand Rapids, MN 55744, USA^d Direction de la recherche forestière, Ministère des Forêts, de la Faune et des Parcs du Québec, 2700, rue Einstein, Québec, QC G10 3W8, Canada

ARTICLE INFO

Keywords:

Northern hardwood forest
Silviculture
Long-term management
Global climate change

ABSTRACT

Northern hardwoods are an economically, ecologically, and culturally important forest type spanning the upper latitudes of the United States and the lower latitudes of Canada. The prevalence and value of these forests have driven silviculture research for over a century. During this time, silvicultural approaches have varied widely, searching for scenarios to meet traditional commodity-based and diversifying ecological forestry objectives. To better understand this forest type and the spectrum of appropriate silvicultural options, we analyzed regional inventory data from the United States and Canada and synthesized decades of scientific studies. Calculated overstory tree (stems ≥ 12.5 cm diameter at breast height) metrics show common structural conditions across mature northern hardwood forests and dominance of sugar maple (*Acer saccharum*). However, density and composition metrics for established reproduction (saplings 2.5 to 12 cm dbh) emphasize challenges for establishing and maintaining economically and ecologically valued trees species broadly and regionally. Our work underscores the variation in northern hardwoods within and across its distribution, driven by characteristics like disturbance regimes, land use history, and ownership patterns. We conclude maintaining this important forest type amid climate uncertainty and associated effects, like proliferation of exotic insects and diseases, requires recalibration of historically applied silvicultural systems and application of emerging tools.

1. Introduction

The northern hardwoods are a wide-ranging forest type, covering approximately 20 million hectares across the northern United States and southern Canada (Leak et al., 1987; Rowe, 1972). Broadly, the northern hardwood forest (NHF) is defined by dominance of three temperate, deciduous species, sugar maple (*Acer saccharum*), yellow birch (*Betula alleghaniensis*), and American beech (*Fagus grandifolia*) (Eyre, 1980; Halliday, 1937). Compositional variability within this larger forest classification, however, is well documented (Beaudoin et al., 2017). Abundance of other species within the beech-birch-maple forest, as the NHF is also known (Gawler and Cutko, 2018), is driven by regional and site-level factors including land-use history (Foster, 1992; Orwig and Abrams, 1994; Schulte et al., 2007), species silvics (Burns and Honkala, 1990; Tubbs et al., 1983), wildlife dynamics (Horsley and Marquis, 1983; Sage et al., 2003), disturbance regimes (Lorimer and White, 2003), and local site attributes (Barnes et al., 1982; Leak, 1978; Nichols,

1935).

The NHF provides a spectrum of ecological, social, and economic benefits, although specifics vary by region, species abundance, or structure. Broadly the NHF plays a critical role in forest nitrogen and carbon cycles, provides vital habitat for mammal (Jensen et al., 2012), birds (Doyon et al., 2005), and amphibians (Hocking et al., 2013), as well as enhanced water quality (Kellison and Young, 1997; Zipper et al., 2011). Colorful fall foliage draws in tourists each year, contributing to local economies, as do specialty products like maple syrup and chaga (*Inonotus obliquus*) (Brydon-Williams et al., 2021; Matthews and Iverson, 2017). The NHF is used for traditional forest products, including timber and pulpwood. In the context of global climate change, the NHF is important for carbon storage (Ford and Keeton, 2017) and area of exploration for carbon markets (Russell-Roy et al., 2014).

Silviculture, the art and science of managing forests for diverse, human defined goals, is an essential tool for maintaining the NHF and the wide array of products and associated values. Decades of research

* Corresponding author.

E-mail address: Nicole.s.rogers@maine.edu (N.S. Rogers).




Fig. 1. Location of regions within the larger northern hardwood forest (NHF) and locations of forest inventory plots used in the Present-Day NHF analysis described below.




Fig. 2. Boxplots of average overstory tree (≥ 12.5 cm dbh) attributes by region A: quadratic mean diameter (cm), B: overstory density (trees per hectare), C: overstory basal area (m² per hectare). LS = Lake States United States, NB = New Brunswick Canada, NE = Northeast United States, and Quebec = Quebec Canada.




Fig. 3. Species dominance for overstory trees (≥ 12.5 cm dbh) by region. Dominance is measured by mean species importance; error bars represent standard deviation. A: New Brunswick, B: Quebec, C: Northeast United States, D: Lake States United States. AB = American beech, SM = sugar maple, YB = yellow birch, oHW = other hardwood species, SW = softwood species. For the mathematical calculation of species importance values see Appendix 1.

have explored how silvicultural systems shift species abundance and structure, generate sustained wood supplies, preserve forest health, and address regeneration challenges. However, the NHF continues to develop with changing forest ownership patterns (Butler et al., 2016; Sass et al., 2020) and transforming environmental conditions (Halofsky et al., 2018), warranting a detailed look at silvicultural tools for maintaining this important forest type. In this review, we define the NHF, broadly and regionally, consider how current species abundance and structural conditions reflect underlying site characteristics, disturbances, and present and past ownership patterns. We also examine the history of silviculture in the NHF and the management options in the face of novel and evolving challenges.

2. Northern hardwood regions

The northern hardwoods are a temperate forest type, occurring in humid and cool locations with average annual precipitation between 80 and 130 cm and range in elevation from 150 to 460 m (Seymour, 1994; Stearns, 1997). Warmer summers are balanced by winters cold enough to maintain a continuous snowpack. Proximity to large bodies of water, including the Great Lakes, the Atlantic Ocean and the St. Lawrence River, moderate temperatures and increase precipitation locally (Stearns, 1997).

2.1. Regional definitions

Across the range of the NHF are distinct regions, defined by unique abiotic and biotic conditions. For this work, we have designated four regions using geopolitical boundaries: the Lake States of the United States (Minnesota, Wisconsin, and Michigan), the Northeast United States (New York, Vermont, New Hampshire, and Maine), and the Canadian province of Quebec and the Canadian province of New Brunswick (Fig. 1). We acknowledge the NHF occurs in states or provinces adjacent to those listed but confined our geographic focus to complete

geopolitical units. For example, the NHF is a component of forests in Pennsylvania, but acreage is limited relative to New York, approximately 1 million vs. 5.6 million hectares respectively (Albright, 2018, 2017). For each region, we consider site characteristics, disturbance regimes/agents, and ownership patterns/land-use history. Although there are additional characteristics specific to each region, we believe these three are essential for understanding inherent species abundance and structure, and appropriate silvicultural options. Beyond the regions we have defined, extensions of the NHF occur in the central and southern portion of the United States, known as Appalachia and the transition to mixedwood forests (80%-20% softwood and hardwood mixtures) in the eastern US and Canada. However, we have excluded these forests from our review as they contain southern hardwood or northern conifer species, respectively, beyond this work.

2.2. Present-day northern hardwood forest

Before considering the attributes of each region mentioned above, we evaluated the structure and species abundance of the current NHF using available forest inventory data (<https://fia.fs.fed.us>, <https://www2.snb.ca/content/gnb/en/departments/erd.html>, <https://mffp.gouv.qc.ca/>; see Appendix 1 for methods). Trends are clear across the broader NHF and by region in the overstory (trees ≥ 12.5 cm dbh) and established reproduction layers (saplings 2.5 to 12 cm dbh). Calculated overstory structural attributes (quadratic mean diameter [QMD, cm] and density [trees per hectare and m^2 per hectare of basal area]), are similar among the defined regions, based on overlapping box-and-whisker plots and comparable median values (Fig. 2). Median values of QMD, approximately 10 cm (Fig. 2A), basal area, approximately $20 m^2 ha^{-1}$ (Fig. 2B), and density, approximately 375 trees ha^{-1} (Fig. 2C), were not visibly different by region. However, New Brunswick has a broader range of structural conditions relative to other regions (Fig. 2). Compositionally, the overstory of the NHF is dominated by sugar maple, evidenced by the high species importance values (IV, %)




Fig. 4. Boxplots of established reproduction (saplings 2.5 to 12.4 cm dbh) structural metrics by region: A. density (trees per hectare) and B. basal area (m² per hectare). LS = Lake States United States, NB = New Brunswick Canada, NE = Northeast United States, and Quebec = Quebec Canada.

compared to other species (Fig. 3). Though, dominance of sugar maple compared to other species varies by region. Importance of the other two defining northern hardwood species, yellow birch, and American beech, varies as well by region (Fig. 3). Considering established reproduction (saplings 2.5 to 12 cm dbh) at the same locations, regional trends emerge. Although abundant, the distributions of sapling density (m² per hectare of basal area and trees per hectare) are wide (Fig. 4). In eastern regions, median densities exceeded 1000 stems per hectare and 4 m²ha⁻¹ of basal area (Fig. 4). Median densities were lowest in the Lake States (Fig. 4A). Sugar maple is the dominant regenerating tree species in the Lake States, but not in other regions where American beech was more abundant based on species importance values (Fig. 5). Yellow birch had the lowest importance value of all regenerating tree species assessed across each region. Each region also had a substantial component of softwood established reproduction (Fig. 5). Median values of sapling density (trees per hectare) meet or surpass regional regeneration guidelines for the NHF (Leak et al., 1987; Marquis, 1975; Tubbs, 1977), but species abundance results emphasize the challenge of regenerating more economically valued species i.e. sugar maple and yellow birch, relative to American beech. This challenge is highlighted in the compositional disconnect between the overstory and established reproduction in stands across the NHF (Fig. 3 and Fig. 5). In the following sections, we consider the role of site characteristics, disturbance regimes/agents, and forest ownership/land-use history in creating these current conditions regionally and broadly in the NHF.

2.3. Site quality & species abundance

Site quality across the NHF is variable but tied closely to soil characteristics and landscape features. Soils in the NHF are classified as podzols (Lull, 1968; Robitaille and Saucier, 1998; Seymour, 1994) developing after the retreat of the last glaciers approximately 10,000 years ago (Stearns, 1997). The glaciated landscape fluctuates from

poorly drained depressions to well-drained topographic features (Stearns, 1997). Divergence in soil characteristics regionally and from stand-to-stand impact species abundance and growth potential. The calcareous parent material in the Lake States generates more productive soils than in eastern regions of the NHF (Schaeztl et al., 2012). Granitic parent material in the east creates rockier, less fertile soils, although bands of calcareous bedrock exist in portions of New York, Vermont, Northern Maine, southwestern Quebec, and eastern New Brunswick (Fahmy et al., 2010; Lull, 1968; Stearns, 1997). Changes in soil quality, specifically nutrient availability, is closely tied to species abundance in the NHF (Johnson et al., 1987; Schmoltdt et al., 1985). Sugar maple, a preferred species for its high economic and social value, increases in abundance on enriched sites with well drained soils (Lindsey et al., 1965; Nyland, 1999). However, the high nutrient requirements for sugar maple often limit the occurrence of pure stands without direct management intervention (Godman et al., 1990). In our overstory and established reproduction analysis, elevated importance of sugar maple in the Lake States (Fig. 2, Fig. 4) likely reflect the higher site quality of the region (Godman et al., 1990) combined with management that favors more shade tolerant species (Crow et al., 2002; Neuendorff et al., 2007) (See Section 3). Prevalence of American beech in eastern regions (Fig. 2, Fig. 4) reflects the underlying range of the species, which stretches from eastern Wisconsin, across southeastern Canada, and the eastern United States (Stephanson and Coe, 2017). The current range of American beech is seemingly controlled by environmental conditions (Woods and Davis, 1989). Within this context, the abundance of American beech in eastern regions likely reflects lower site quality too, which makes site demanding species, like sugar maple, less competitive (Tubbs and Houston, 1990; Ullah and Moore, 2009). The history of beech bark disease (BBD), which triggers root suckering (see Section 2.3), in the eastern regions is also suspected to influence prevalence of American beech.

Assessing and understanding site quality is a fundamental




Fig. 5. Species dominance for established reproduction (saplings 2.5 to 12.4 cm dbh) by region. Dominance is measured by mean species importance; error bars represent standard deviation. A: New Brunswick, B: Quebec, C: Northeast United States, D: Lake States United States. AB = American beech, SM = sugar maple, YB = yellow birch, oHW = other hardwood species, SW = softwood species. For the mathematical calculation of species importance values see Appendix 1.

underpinning of silviculture (Ashton and Kelty, 2018; Smith et al., 1997). Site quality, specifically soil fertility and drainage, are recognized as key controls on the species establishment and stand-level productivity in the NHF (Henry et al., 2021; Post, 1962). Further, silvicultural systems, when linked with site characteristic, in addition to other important attributes, are more likely to achieve species compositional goals (Willis et al., 2016). For example, in the eastern United States, sites on washed glacial till managed with group selection retained yellow and paper birch (*Betula papyrifera*) even with a component of American beech, while sugar maple was best maintained with single-tree selection, but only on enriched sites (Leak, 1980). Careful consideration of the site has been an integral part of selecting appropriate silvicultural systems within the NHF for nearly half a century (Baker, 1934; Leak, 1978; Marquis, 1975).

2.4. Disturbance regimes and disturbance agents

We propose three primary disturbance factors related to the results from our overstory and established reproduction analysis: disturbance regime, ungulate browsing, and beech bark disease. Similar to site quality and associated characteristics, natural disturbance regimes and disturbance agents influence the structural and compositional dynamics of the NHF (Bormann and Likens, 1979; Oliver, 1981). Commonly, natural disturbance events in the NHF are defined by size, severity, and frequency (Lorimer and White, 2003). For example, disturbances may be classified as local, low-intensity and frequent to regional, severe, and infrequent (Lorimer and White, 2003) although disturbances beyond this gradient are also important for successional and developmental dynamics (Franklin et al., 2002; Frelich and Lorimer, 1991). Small-scale disturbances, occurring every 50–200 years, establish canopy gaps in the forest between 4 and 1,135 m² in size (Seymour et al., 2002). These small openings favor growth of shade tolerant species like sugar maple,

American beech, and eastern hemlock (*Tsuga canadensis*), a common associate in the NHF (Canham, 2010). Sugar maple and beech are highly shade tolerant, persisting under low light conditions for decades. Larger stand replacing events occur every 800 to 9,000 years, disrupting 1 to > 80,000 ha (Seymour et al., 2002). The scale and frequency of meso-scale disturbances are intermediate relative to small and large-scale events, playing an equally important role in development of these forests (Jenkins, 1995; Wood et al., 2009). In the NHF, a spectrum of disturbance types occur across the scale of disturbance size, intensity, and frequency (Lorimer and White, 2003). For example, ice storms are associated with small-scale, insect outbreaks or drought with meso-scale, and fire with stand-replacing events (Brewer and Merritt, 1978; Krasny and Digregorio, 2001; Pederson et al., 2014; Rustad and Campbell, 2012). Wind disturbances, while often a small-scale, can also be a meso-scale, or even stand-replacing (Bormann and Likens, 1979; Rentch et al., 2010).

Increased canopy openings are important for establishing shade intolerant and mid-tolerant species such as yellow birch, paper birch, and white ash (*Fraxinus americana*) (Archambault et al., 1998; Godman et al., 1990; Tubbs, 1977). The low importance of yellow birch in the sapling size classes (Fig. 5) could signal greater prevalence of small-scale disturbances, both naturally occurring and human induced (management), across the NHF (Godman and Krefting, 1960; Leak, 1980), as well as differences in litter quality (Hupperts et al., 2020). The structural similarities in the overstory between regions (Fig. 2, Fig. 4) are likely explained by overarching disturbance regimes and similar management histories (Brown et al., 2018).

In addition to the broader disturbance regimes associated with the NHF, distinct disturbance agents are important to regional dynamics. Browse by white-tailed deer (*Odocoileus virginianus*) is one such example. Although influence of heavy deer browsing can be negative and long-lasting across the NHF (Boucher et al., 2004; Nuttle et al., 2013;

Table 1
Plot count by dataset used to analyze the current northern hardwood resource.

Dataset	# of Plots
United States	4042
New Brunswick	434
Quebec	1046

Tanentzap et al., 2011), the impact of over-browsing is especially clear in the Lake States where deer populations can exceed 17 per km² (Walters et al., 2016). Significant changes in forest composition, overall abundance of reproduction, and increased susceptibility to invasive species have been observed in forests with deer densities above 4–6 per km² (Alverson et al., 1988; Horsley et al., 2003; Walters et al., 2016). Many studies show the importance of controlling deer populations to ensure forest regeneration, but management can be difficult (Miller and Graefe, 2001; Royo and Stout, 2017; Witmer and DeCalesta, 1991). Local deer populations vary with winter severity, hunting policies, and ranges of matriarchal groups (Nesslage et al., 2001; Sage et al., 2003). Further, deer management is a topic of interest to a wide range of stakeholders whose objectives do not always align (Woolf and Roseberry, 1998). We suspect lower sapling densities observed in the Lake States compared to other regions of the NHF are partially because of this browse phenomena (Fig. 2, Fig. 4). In the eastern regions, where BBD triggers prolific root sucking in smaller size classes (see below), competitive interactions between American beech and more economically desired species (sugar maple and yellow birch) are further exacerbated by heavy browse pressure as American beech is less palatable to deer (Tierson et al., 1966). The combined dynamics of BBD and browse presumably affected the elevated abundance of established American beech reproduction (Fig. 5).

Beech bark disease first arrived in the eastern United States and Canada at the turn of the last century, permanently altering the NHF (Houston, 1975; Shigo, 1972). Today, the BBD complex (scale insect *Cryptococcus fagisuga* and fungus *Nectria coccinea* var. *faginata*) is widespread, although the impact of the disease is not consistent from region to region. In the Lakes States as well as northern portions of Quebec, BBD is a more recent arrival, creating different American beech dynamics relative to locations where the disease has been active for nearly a century, including the northeastern U.S. (McCullough and Wiererich, 2015). Three stages of BBD are recognized: the advancing front, killing front, and aftermath zone (Houston, 1994). The advancing front represents the earliest stages of the disease complex with the scale insect visible on infected trees, but with limited presence of the fungus. The killing front is associated with widespread mortality of American beech, particularly on large diameter trees, whereas the aftermath zone is defined by mortality of most large-diameter American beech, some remaining resistant trees, and numerous small, root-origin sprouts (Houston, 1994; Twery and Patterson, 1983). Much of northeastern North America is in the aftermath zone, while central and far northern locations are in the advancing front (Giencke et al., 2014). The shade tolerant nature of American beech and the ability to prolifically sucker following harvest or injury complicates management of hardwood forests, especially in systems where sugar maple and yellow birch are preferred (Jones and Raynal, 1988; Tubbs and Houston, 1990). The high importance of American beech in the established reproduction size classes, particularly in the eastern United States and southeastern Canada, is attributed to the long history of BBD in those regions (Fig. 3, Fig. 5).

In addition to BBD, the spruce budworm (*Choristoneura fumiferana*) is another disturbance agent that has shaped the structure and composition of eastern NHFs, particularly in Quebec and New Brunswick. Outbreaks of spruce budworm, a native insect, and subsequent mortality of softwoods, namely balsam fir (*Abies balsamea*), shifted composition of mixed conifer-hardwood forests (mixedwoods) to more hardwood

dominance (Danneyrolles et al., 2016; Dupuis et al., 2020). Compositional relics from previous disturbances likely contribute to the disconnect between current overstory and established reproduction species abundance (Fig. 3, Fig. 5).

Beyond the disturbance agents we have identified, numerous others exist, regionally and broadly. We acknowledge these disturbances have impacted the current structure and species abundance of the NHF. However, we have focused on the disturbance agents above as we see clear connections between them and the results from our overstory and established reproduction analysis.

2.5. Forest ownership and land-use history

Like site characteristics and disturbance agents, the influences of present and past ownership types, patterns, and objectives are seen throughout the current NHF. In the United States, forests are largely privately owned and within that classification, predominantly family owned (Sass et al., 2020). Property size within the private, family-owned classification ranges from 0.4 to 3.6 ha (Butler et al., 2021). However, ownership patterns for remaining forestland differ between the Lake States and Northeast. In the Lake States, the dominant land ownership group, family ownership, is followed by federally- and state-owned forests, while, in the Northeast, private corporations follow family ownership (Sass et al., 2020). Small, private individual and families often see timber extraction as a secondary goal (Leak et al., 2014; Rickenbach and Kittredge, 2009). Rather, these small landowners manage their woodlands for wildlife habitat and aesthetics (Kelty et al., 2003). For larger private landowners, including real estate investment trusts (REITs) or timber management organizations (TMOs), the management emphasis is on traditional forest products and return for investors (Mendell, 2016). Across Canadian forestland, public ownership is most prevalent, although ownership patterns differ between Quebec and New Brunswick too. Ninety percent of Quebec's forests are Crown land, while 50% of New Brunswick is Crown land (The State of Canada's Forests Annual Report, 2019). The additional forestland land in New Brunswick is divided among private ownership and industrial free holdings, 29 and 18% respectively. The high percentage of public ownership in Canada can cause unique management opportunities and challenges. Integration of societal expectations in natural resources management can be found across many forest ownership (Moffat et al., 2016). However, balancing the social license to practice forestry with the legal license is especially true in Canada where forest managers are stewards of a publicly owned resource (Beckley, 1998; Chambers and Beckley, 2003). An example of public input changing forestry practices in Canada is the banning of chemical herbicide on Crown lands in Quebec (Thiffault and Roy, 2011). In other regions of the NHF chemical herbicide is still used as a tool for controlling competing vegetation, namely American beech (Kochenderfer et al., 2004; Nyland et al., 2006).

Present day ownerships are part of a long history of management in the NHF, beginning with Indigenous Peoples. Early use of forest resources varied by region and forest type, but documentation of periodic burning, manipulation of wildlife, and harvesting for traditional and non-traditional forest products exists (Bromley, 1935; Emery, 2002; Patterson and Sassaman, 1988). In southern New England for example hardwood species such as oak (*Quercus* sp.) and American chestnut (*Castanea dentata*) were favored after fires, often set for agriculture purposes, (Abrams, 2000; Orwig and Abrams, 1994) while sugar maple, yellow birch, and American beech were relegated to smaller areas because of their low fire tolerance (Fuller et al., 1998). Canadian forests saw similar manipulation of fire regimes by First Nations people, although documentation is limited (Dupuis et al., 2020). The northern hardwoods have further been influenced by detrimental government policies during the 1800 s and 1900 s that limited land ownership by Indigenous Peoples. An exception to these policies is the Menominee tribe of Wisconsin (Mausel et al., 2017; Trosper, 2007). The Menominee have practiced sustainable forest management through thoughtful

removal of mature trees and tending to the residual forest stand for over 150 years (Kern et al., 2017; Pecore, 1992). Today, their forestland offers a sharp contrast to other forests across the Lake States with histories of more degrading forest practices (Sands and Abrams, 2011).

The arrival of European settlers to North America in the early 1600s triggered wide-spread land clearing, for construction and agriculture, shifting the age and species abundance dynamics of the NHF (Cogbill et al., 2002; Hermy and Verheyen, 2007). Heavy partial harvesting of red spruce (*Picea rubens.*), balsam fir, and eastern white pine (*Pinus strobus*) across temperate forests in the late 19th and early 20th century, moving composition from mixedwoods to more hardwood dominance (Bryant, 1917; Dupuis et al., 2011; Linn, 1918; Pinto et al., 2008; Westveld, 1949). In pure hardwood stands, high quality sawlogs were selectively removed, leaving unmerchantable trees of poor quality and vigor (Blum and Filip, 1963; Hall et al., 2002). Structure of the NHF changed again in the 1980s and 1990s with increased demand for hardwood pulp favoring removal of smaller stems (Luppold and Sendak, 2004). Today, these land use changes have resulted in large areas of even-aged northern hardwoods in regions recovering from agricultural abandonment (eastern regions) and clearcutting (Lake States) of the past centuries and irregular, often poor stand conditions in other regions where exploitative partial cuts prevailed (Kenefic and Nyland, 2006). Forest conditions created by past ownership and historical objectives, presumably create the compositional disconnect between overstory and established reproduction highlighted in our present-day analysis of NHF conditions (Fig. 3 and Fig. 5).

3. Silviculture history in the northern hardwood forest

Silvicultural systems for the NHF have long been governed by compositional and structural goals (Eyre and Neetzel, 1937; Jensen, 1943; Leak et al., 1969). However, effective management also incorporates the underlying attributes driving the current forest condition. Early research explored management systems for sustained yield of timber; a common guiding paradigm for research in many forest types in the early 20th century (Seymour, 2004; Stout et al., 2006). Foundational studies assessed harvest intensities of different silvicultural systems and subsequent effects on forest composition and structure (Eyre and Zillgitt, 1953; Gilbert and Jensen, 1958). These early works established silvicultural recommendations that continue to be used in the NHF and inspired silvicultural guides for other reaches of the northern hardwood range (Arbogast Jr, 1953; Leak et al., 1987; Majcen et al., 1984; Pond et al., 2014). Successive research has investigated species response to habitat (Barnes et al., 1982; Carmean, 1999), interference of competing vegetation with tree regeneration (Nyland et al., 2001; Willis et al., 2015), and how to emulate natural disturbances (Coates and Burton, 1997; Franklin et al., 2007; Hanson and Lorimer, 2007) in an effort to generate consistent outcomes and effective silvicultural systems. But forest conditions continue to advance as do management objectives (Butler and Leatherberry, 2004). This advancement includes management for global climate change (Bolton and D'Amato, 2011; Parker et al., 2000; Webster et al., 2018), improved emulation of underlying ecological principles (D'Amato and Palik, 2021; Kern et al., 2019) and broader ecosystem services in general (O'Hara, 2016). Given the increasingly complex nature of northern hardwood forests and the evolving challenges to management, silvicultural responses require careful evaluation of all available options.

Throughout the history of northern hardwood silviculture, use of and preference for different systems has oscillated between extremes (O'Hara, 2002; Smith, 1972). Unfortunately, such oscillation may leave valuable tools and management strategies out of consideration (Kern et al., 2014; Pond et al., 2014). In the context of the NHF, numerous studies have shown successful development of these forests with a wide range of silvicultural systems (Burns, 1983) and conversely, inconsistent results when the same method is applied across different conditions or locations (Bédard and Majcen, 2003; Neuendorff et al., 2007). As such,

preference for silvicultural systems fluctuates between extremes of even and uneven-age approaches over time (Smith, 1972).

3.1. Even-age systems

Even-age forests do occur in the NHF after natural, stand-replacing disturbances. However, this age structure is more frequently the product of historical, intensive land-use or more recent harvesting history. Clearcutting and uniform shelterwood systems have been applied in the region since the early period of formalized forest management to create even-age conditions (Leffelman and Hawley, 1925); however, these approaches have not experienced the wide-spread popularity of uneven-age silvicultural systems such as group and single-tree selection. In part, the limited application of even-age silviculture at a broad scale is due to historical and ongoing public reaction to these silvicultural systems relative to other, less intensive approaches (Hannah, 1988; Kelty et al., 2003). This is especially true of clearcutting. Public opposition to clearcutting is widespread with many associating the practice with environmental decline and detrimental management (Bliss, 2000). In some instances, public responses to clearcutting have initiated policy debates and reform in an effort to regulate use on federal (Fairfax and Achterman, 1977; Spurr, 1981) and private lands (Steelman and Ascher, 1997). Yet, despite constraints to these approaches, clearcutting has been demonstrated as an economical and efficient method for managing northern hardwoods, favoring more shade intolerant species (Hornbeck et al., 1986). In central New York, clearcutting increased species diversity in the newly developing cohort and resulted in stands with high stocking of commercially valuable species including sugar maple and yellow birch (Wang and Nyland, 1993). In New Brunswick, clearcutting promoted new germinates relative to advance reproduction, mostly yellow birch and pin cherry (*Prunus pensylvanica*) (Roberts et al., 1980). In the Lake States, 5-acre clearcuts successfully regenerated economically desired species in old-growth NHF, but not in second-growth, due to advance reproduction on site and diverse seed sources in the former and not the latter (Metzger and Schultz, 1984). Complete removal of the overstory in clearcuts also creates needed habitat for early successional bird species (Costello et al., 2000; Yamasaki et al., 2014) as well as mammals, reptiles, and amphibians that rely on young forest conditions (DeGraaf and Yamasaki, 2003; Moorman et al., 2011). Although benefits of clearcutting are clear for some conditions, ecological concerns exist as well, especially reduction in biodiversity of non-tree species and loss of mature forest legacies critical to sustaining certain species and processes in regenerating forests (Rudolphi et al., 2014). Increasing mature tree retention, specifically large biological legacies, is an increasingly common strategy to address these concerns (Gustafsson et al., 2010).

Shelterwood systems are another even-age method where public opinion or policy guidelines make widespread implementation difficult (Hannah, 1988). Outcomes vary by the number of entries, time of harvesting, and density of the overwood, but when these attributes are appropriately matched to stand conditions and objectives, shelterwood cutting is an effective method for regenerating northern hardwood forests in all NHF regions (Godman and Tubbs, 1973; Hannah, 1988). Research from the Adirondack region of New York showed ample regeneration of yellow birch, sugar maple, and white ash in deer exclosures following a two-stage shelterwood, outside exclosures regeneration was dominated by American beech and hobblebush (*Viburnum lantanoides*) (Curtis and Rushmore, 1958). Kelty and Nyland (1981) saw similarly abundant reproduction of desirable species using a two-stage shelterwood in combination with hunting to reduce populations of white-tailed deer and pretreatment mist blowing of herbicides to remove competing vegetation. In the Allegheny region of the United States, shelterwood cutting successfully regenerated commercial valuable northern hardwood species in combination with herbicide site preparation (Ristau et al., 2011). A crucial component of shelterwood systems is layout of the final harvest to remove the overwood. Jacobs (1974) observed 35% of established reproduction was damaged during

the removal cut in a northern hardwood stand, but overall reproduction was successful due to a major regeneration event following the initial cut. Yet, economic, and operational difficulties in removing the overwood (Leak et al., 2014) and landowner preference for less visible disturbance (Kelty et al., 2003) remain reasons uniform shelterwood systems have not been more widely adopted in northern hardwood forests of the northeast U.S. However, spatial, and temporal variations of the shelterwood system, including integration of reserve trees or more complex distribution of the overwood can be used to meet broader ecological goals (Ashton and Kelty, 2018; Palik et al., 2020).

3.2. Uneven-age systems

Preliminary guidelines were developed for single-tree selection based on theoretical balanced stand conditions aimed at sustaining a regular supply of timber over time and improving growth of the residual stand (Eyre and Zillgitt, 1953). Long-term research across the range of northern hardwoods has demonstrated successful regeneration and increased stand quality with careful application of single-tree selection, especially with efforts for American beech control and deer browse impacts (Jones et al., 1989). The implementation of single-tree selection has been most appropriate on sites where an increased abundance of tolerant species that yield high value products are desired results (Keyser and Loftis, 2013). Efforts to regenerate and maintain mid-tolerant species after repeated applications of single-tree selection have not been as successful (Johnson, 1984; Webster et al., 2018). The inherent favoring of large, shade tolerant trees have spurred extensions of traditional single-tree selection systems to generate late successional forest habitat (Keeton, 2006). Yet, for larger scale applications, single-tree selection requires a high skill level for marking implementation (Brockway et al., 2015). Further, the challenges of creating and maintaining a truly balanced stand structure, as well as questions about the propriateness, have increased the interest in alternative options for northern hardwood management (Hicks, 1998; Janowiak et al., 2008).

Group and patch selection are additional strategies for northern hardwood management, noted for maintaining uneven-age structure while recruiting mid-tolerant and tolerant species (Poznanovic et al., 2013). Long-term data from New England demonstrated that increased light in group selection relative to single-tree selection increased the component of desirable species, namely yellow birch and sugar maple, by providing a competitive advantage over beech (Leak and Filip, 1977). Early results in Ontario found similar compositional trends after group selection harvesting (Falk et al., 2010). This approach may also offer productivity benefits, as a study of northern hardwoods in Wisconsin showed increased growth in small group openings relative to single-tree selection openings with productivity reaching an asymptote as gaps size increased above 100 m² (Webster and Lorimer, 2005). While still considering structure of the stand, group selection offers increased flexibility in target diameter distributions where marking is driven by presence of advance reproduction or pockets of economically mature trees (Leak and Gottsacker, 1985). Patch selection, also a method of uneven-aged management, is similar to group selection but creates larger canopy openings, between 0.1 and 0.8 ha in size or greater than two tree lengths (Ashton and Kelty, 2018; Leak et al., 2014). Openings are created using stand area regulation, while removing single trees throughout the surrounding matrix (Leak et al., 2014; Nyland, 2005). Uneven-aged management with emphasis on greater canopy disturbance is important for regenerating and retaining more mid and intolerant hardwood species (D'Amato et al., 2015; Yamasaki et al., 2014). Additional and essential benefits of selection silviculture with larger openings include control of American beech and increased habitat for wildlife, specifically songbirds. In NHF's where a persistent understory of American beech exists, the larger canopy openings, relative to single-tree selection, remove beech advance reproduction and create an early competitive advantage for more commercially desirable species (Leak, 2005). Larger openings in group and patch selection create early seral

habitat too, especially in the first years following harvest (DeGraaf and Yamasaki, 2003). These stages of forest succession and development provide important structural and composition conditions for breeding songbirds (Costello et al., 2000). Generating this habitat with existing silvicultural options is particularly important as songbird populations decline (Bonnot et al., 2018).

Group and patch selection have become increasingly popular methods for managing northern hardwood forests, but as seen in other common silviculture systems, are not without their challenges. Balancing ecological benefits with operational efficiency is one such challenge, particularly when considering the layout of openings. Establishing groups or patches is a tool for increasing structural diversity in more homogenous stands (D'Amato and Palik, 2021), but arbitrary location of openings disregards the benefits of these treatments for accommodating spatial variability i.e. simultaneous removal of economically mature overstory trees and release of well stocked advance reproduction (Minckler, 1972; Murphy et al., 1993). Considering appropriate orientation and shape of openings is also necessary to ensure consistent results for goals of sustained timber supply and predictable species composition (Marquis, 1965). Over time, group and patch selection can increase the complexity of the forest by diversifying age structures and species abundance, which has strong ecological benefits (Hanson et al., 2012). However, successfully managing forests with group selection, like many silvicultural tools, requires careful consideration of multiple factors beyond opening size alone, including microsites, local disturbance regimes, and competing vegetation (Kern et al., 2017).

3.3. Hybrid systems

An alternative approach to more rigid even and uneven-aged systems is the irregular shelterwood system. Primarily in eastern Canada, but also in New England (Peterson and Maguire, 2004) and to some degree the Lake States (Helman et al., 2021), irregular shelterwoods have been proposed as a hybrid of traditional even and uneven-aged management for mixedwood and hardwood forests (Raymond et al., 2009). Irregular shelterwood systems are an approach to capture the ecological complexity, particularly species abundance and structure, of forests following mesoscale disturbances (Raymond and Bédard, 2017). The continuous cover and expanding gap versions of this system have increased spatial and temporal flexibility relative to traditional shelterwood, single-tree, and group selection methods to meet these objectives (Raymond et al., 2009). In Quebec, Canada continuous cover and extended irregular shelterwood systems have been employed to rehabilitate impoverished northern hardwood stands following repeated selective cutting (Bédard et al., 2014). For stands with a low percentage of acceptable growing stock and high component of American beech, Bédard et al. (2014) found irregular shelterwood systems may be more appropriate than single-tree selection if desired species are those that could be regenerated in groups; however, cutting cycle lengths would need to be greater than those used in single-tree selection given heavier volume removals at each entry. Initial results for irregular shelterwood studies are promising in terms of increasing species and structural diversity, metrics of increasing importance when assessing forests for changing climate and market conditions (Christel C Kern et al., 2017).

4. Northern hardwood silviculture, climate change, and options for the future

As we consider the prospects of the NHF, our look forward must be framed in the context of global climate change and the ambiguity of future forest conditions. Presently, forests store approximately 45% of terrestrial carbon, and have been highlighted as a critical component of climate change mitigation strategies (Bonan, 2008). These mitigation strategies include silvicultural recommendations for carbon storage and sequestration as well as new carbon markets and emphasis on longer-

lived forest products (Ameray et al., 2021; Landry et al., 2021; Winans et al., 2016). However, there remains uncertainty about the frequency of future forest disturbances, shifts in resource availability, and ultimately, movement of species ranges (Dale et al., 2001; Iverson et al., 2001; Sykes et al., 1996). It is this uncertainty, that requires careful consideration as we evaluate silvicultural options moving forward.

Field observations and model scenarios offer some insight as we contemplate the vulnerability of species or forest communities to changing environmental conditions (Iverson et al., 2008; Nagel et al., 2017). For sugar maple and yellow birch, two defining species of the NHF, abundance across their ranges is expected to decrease (Peters et al., 2020). This decrease is anticipated to be most severe for sugar maple where largest populations could be limited to zones of climate refugia (Oswald et al., 2018). Occurrence of American beech, conversely is increasing throughout portions of its range (Bose et al., 2017). Shifts in historical species composition is further exacerbated by alterations in natural disturbance regimes and invasive diseases and insects (Lovett et al., 2016). For example, non-native and invasive earthworms have no historical context in the NHF but are triggering declines in forest health through depletion of leaf litter and soil nutrients important for regenerating key species, like sugar maple (Burtelov et al., 1998; Callahan et al., 2006). Prevalence of ash species (*Fraxinus* spp.) in the NHF is also on the decline following fatal infestations of the invasive emerald ash borer (*Agrilus planipennis*) (DeSantis et al., 2013). As temperatures warm, the invasion range of the emerald ash borer is predicted to increase, directly threatening the ecological and cultural values associated with ash forests (Iverson et al., 2016; Looney et al., 2017). Beech bark disease and forest tent caterpillar (*Malacosoma disstria*), a native defoliator, are other important disturbance agents that may benefit from warmer climates, especially during winter months (Dukes et al., 2009). The intensity and frequency of natural disturbances are increasing too, including droughts (Dale et al., 2001), windstorms (Peterson, 2000), and ice storms (Rhoads et al., 2002). The cumulative impact of these changing conditions are large-scale shifts in the structural, compositional, and functional baselines that have long defined the NHF (Churchill et al., 2013). These shifts are not without challenges, but novel circumstances also create unique opportunities to reframe tools and strategies for evolving forests and management objectives.

Managing forests for adaptability and resiliency has been suggested as an approach that may allow forests and forest managers to buffer the uncertainty of future conditions (Millar et al., 2007; Spittlehouse and Stewart, 2003). Analysis of the outcomes of long-term silviculture studies in the Lake States region of the United States found multi-aged approaches, including irregular shelterwoods, may be the most effective at balancing mitigation and adaptation at the stand-level because they capture a wide range of historical disturbances and subsequent forest complexities (D'Amato et al., 2011; Hupperts et al., 2020). Integration of complexity into forest management frameworks can further support ecosystem resilience by retaining diverse species and functional niches, reducing fire, drought, and browse risk, and ultimately, more closely aligning human-managed systems with underlying natural drivers (D'Amato and Palik, 2021; Drever et al., 2006; Kern et al., 2019).

Within the NHF, specific activities such as adaptation planting, could be added to existing or evolving silvicultural systems for further resilience to climate change. Planting species outside their native range, as is the case with assisted migration, can be controversial and requires upfront financial investments (Messier et al., 2019). However, a proactive approach to changing species ranges may limit the mis-alignment between current species distributions and climate conditions (Williams and Dumroese, 2013). Studies within and beyond the NHF found assisted migration efforts, including adaptation plantings, maintained stand-level productivity, conserved endangered species, and increased overall species diversity (Boulanger and Puigdevall, 2020; Dumroese et al., 2015; Duveneck and Scheller, 2015). Clark et al. explored adaptation planting in the northeastern portion of the NHF and their recommendations were cautiously optimistic (2022). Yet, the authors were

clear, successful adaptation planting requires consideration of site attributes, like competing vegetation, and preparation for events like elevated summer droughts or unseasonable spring frosts which may be the new climate reality (Clark et al., 2022).

Utilization of slash walls are another example of creative problem solving within the NHF. Although this strategy is relatively novel, early results are promising. Smallidge et al. explored the use of slash walls to mitigate the effects of deer browsing in central New York (2021). Four years after harvesting, seedlings within the slash wall interior had higher height growth than surrounding areas. Additionally, slash wall construction was less expensive than traditional enclosures and provided a use for low-value or non-commercial species (Smallidge et al., 2021). Creating browsing refugia is especially important in the context of climate change, where regenerating preferred species could already be difficult or associated with planting costs (Champagne et al., 2021). The spectrum of changes anticipated for the NHF is wide and the magnitude of impact uncertain but careful modification of existing management strategies and integration of new tools provides a path forward (Keenan, 2015; Price et al., 2001; Spittlehouse and Stewart, 2003).

Flexibility in management is central to development of healthy and well-functioning forest under future uncertainty (Dukes et al., 2009). History in the NHF and other forest types reveals periods of preferences for one extreme of silvicultural options or another (O'Hara, 2002; Smith, 1972). Such actions were restrictive then and would be even more detrimental today. Rather forestry practitioners need to consider all options, including novel tools and modification of existing silvicultural systems, guided by management approaches best suited to their specific objectives and forest conditions (Millar et al., 2007). In the context of changing future conditions, an overriding objective for selection and application of any given approach should be the maintenance of a wide range of structural and functional conditions at the stand and landscape-level as a precautionary strategy for addressing the uncertainties associated with global change impacts (D'Amato et al., 2011; Puettmann, 2011).

5. Conclusions

The northern hardwoods are a geographically broad and highly valued forest type. For decades, silviculture research across the NHF has explored management strategies for balancing a range of objectives, including generating a sustainable yield of timber, restoring ecological diversity, creating wildlife habitat, and maintaining visual aesthetics. In addressing these objectives, silviculture in the NHF has historically fallen into two distinct categories aimed at achieving even-aged or uneven-aged stand structures. Our review of the existing literature, and subsequent analysis of regional inventory data, highlights the value and challenges associated with these silvicultural approaches. Further, our results emphasize structural, compositional, and functional variability of the NHF within and across its range. Silviculture that incorporates this variability is especially important as we look forward to an increasingly uncertain future because of climate change and the proliferation of non-native insects and diseases. Successfully establishing, growing, and maintaining northern hardwoods into the future will require a modification of traditional silvicultural approaches with new and creative tools, such as adaptation planting and irregular systems, to provide the range of options necessary and adapt with emerging management challenges.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We would like to thank Chris Woodall from the and Dale Wilson from the New Brunswick Department of Natural Resources for their help compiling regional forest inventory data. We also thank Hunter Stanke

and Lance Vickers for troubleshooting R code. Funding for this research was provided by the University of Vermont Rubenstein School of Environment and Natural Resources, Department of Interior Northeast Climate Adaptation Science Center, and .

Appendix 1. Northern hardwood data analysis

Our assessment utilized forest inventory data from the USDA Forest Service Forest Inventory Analysis (FIA) program (<https://fia.fs.fed.us>), the New Brunswick Department of Natural Resources (<https://www2.snb.ca/content/gnb/en/departments/erd.html>), the Quebec Ministry of Forests, Wildlife, and Parks (<https://mffp.gouv.qc.ca/>).

The USDA Forest Service Forest Inventory Analysis program is a national census comprised of a network of permanent sample plots established on a grid system across the United States. The inventory plot is constructed of four, circular 7.2 m radius subplots with nested micro-plots using a radius of 2 m. All trees ≥ 12.7 cm dbh are measured on subplots, while saplings (2.5 to 12.4 cm dbh) and seedlings (< 2.5 cm dbh, height ≥ 15.2 cm for softwood species, ≥ 30.4 cm for hardwood species) are measured on micro-plots. Field crews re-measure plots on a 5-year cycle with approximately 20% of plots measured each year (Burrill et al., 2018). Data included in this analysis were from the most recent and complete inventory of each state available at the time data acquisition in January 2020. We used the EVALID code 2018 to select all forested plots inventoried in each state between 2012 and 2018. Data were downloaded and extracted using the software R (version 3.6.1 “Action of Toes”, <https://www.r-project.org/>) and the package rFIA (Stanke et al., 2020).

Forest inventory data for the Canadian province of New Brunswick were obtained from the New Brunswick Department of Natural Resources’ Continuous Landscape Inventory (CLI). Data from the CLI are measured on circular fixed area plots with a nested design. Overstory trees ≥ 7.1 cm dbh are measured on 400 m² plots and trees < 7.1 cm dbh are measured on 50 m² plots. 25% of plots in the CLI network are permanent sample plots (PSPs) measured every five or ten years depending on plot designation. Trees measured on PSPs are tagged and numbered. Remaining CLI inventory data are collected every ten years, trees are not tagged or numbered. Data for this analysis are from the most recent inventory of each plot. Data are stored in excel data files available from the New Brunswick Department of Natural Resources.

The Quebec Ministry of Forests, Wildlife, and Parks created a permanent plot network across the Province of Quebec beginning in 1970. The network is intended for monitoring of forest dynamics over time and plots are randomly located within strata defined by species subzones. Sampling intensity the northern hardwood subzones is 1 plot per 26 km². Overstory trees, saplings, and seedlings are measured with a nested plot design. Trees (9.1 cm to 29 cm) dbh are recorded on a circular 11.3 m radius plots and trees > 29 cm are recorded on a circular 14.1 m plots, saplings, trees 1 to < 9 cm dbh, are recorded on a 3.6 m radius circular plot, and seedlings, stems > 60 cm in height and ≤ 1 cm dbh are tallied using two circular 1.13 m plots. Plots are remeasured every approximately every 10 years. For this analysis, we used data from the most recent inventory of each plot. Inventory data were extracted from a Microsoft Access database available through the Quebec Ministry of Forests, Wildlife, and Parks website.

For this analysis, overstory trees are those ≥ 12.5 cm dbh. Saplings are classified as trees 2.5 cm to 12.4 cm dbh.

A.1. Study area and plot selection

The study area for this work includes seven U.S. States (Maine, New Hampshire, Vermont, New York, Michigan, Wisconsin, and Minnesota) and two Canadian provinces (New Brunswick and Quebec). To discern regional trends, states were also aggregated into larger ecological blocks. We defined Maine to New York as the northeast region of the U.S., Michigan to Minnesota as the Lake States region. The number of plots per dataset are listed in Table 1.

Across all data sources, we selected a subset of inventory plots classified as northern hardwood based on dominance of three key species, American beech, sugar maple, and yellow birch. We considered a plot to be northern hardwood forest if species abundance by basal area, of each species singly or cumulatively, exceeded 50% for trees ≥ 2.5 cm dbh. To prevent selection of mixedwood forests we also limited softwood composition to $< 26\%$ by basal area (Vickers et al., 2021). We choose these criteria to capture the broad definition of the northern hardwood forest (Eyre, 1980). Basal area calculations and plot selection were determined using the R statistical package, version 4.0.5 (<https://r-project.org>). The location of each northern hardwood plot was mapped using QGIS (version 3.12, <https://qgis.org/>).

A.2. Data analysis

To assess the northern hardwood resource, we evaluated overstory trees and established reproduction for each state/province and region. We contained analysis to attributes we could calculate across all data sources. For overstory trees we considered four attributes, species importance values (IV), quadratic mean diameter (QMD [cm]), density (trees ha⁻¹), and basal area (m² ha⁻¹). We determined IV for five species/species groups, American beech, sugar maple, yellow birch, other hardwood, and softwood using the formula $IV = (RD_{den} + RD_{dom})/2$. RD_{den} is relative density and RD_{dom} is relative dominance where relative density is defined as the proportion of a given species/species group (tree ha⁻¹) to the total number of species a plot. Relative dominance is the basal area (m² ha⁻¹) of a given species/species group compared to total basal area per plot. Relative density was defined as the proportion of a given species (tree ha⁻¹) compared to the total number of species in a cohort. Relative dominance was defined as the basal area (m² ha⁻¹) of a given species compared to total basal area in a cohort. Relative density and dominance were expressed as percent of the total. We included QMD as an alternative measure of stage of stand structural development appropriate in forests that may have multiple age classes (Lorimer and Frelich, 1998). Boxplots were generated for each overstory attribute with state/province and region as the independent variable.

Two characteristics were calculated to represent regeneration, density (trees ha⁻¹) and species importance values, as described above. Species specific values were determined for American beech, sugar maple, yellow birch, other hardwood species, and softwood species. All values were calculated for the state/province and region. Analysis for overstory and regeneration data was completed using the R statistical package, version 4.0.5.

References

- Abrams, M.D., 2000. Fire and the ecological history of oak forests in the eastern United States. In: *Workshop on Fire, People, and the Central Hardwoods Landscape*. Richmond, KY, pp. 46–55.
- Albright, T.A., 2018. *Forests of New York 2017*. Newtown Square, PA.
- Albright, T.A., 2017. *Forests of Pennsylvania 2016*, United States Department of Agriculture: Resource Update FS-132. Northern Research Station, Newtown Square, PA.
- Alverson, W.S., Waller, D.M., Solheim, S.L., 1988. Forests too deer: edge effects in northern Wisconsin. *Conserv. Biol.* 2, 348–358.
- Ameray, A., Bergeron, Y., Valeria, O., Montoro Girona, M., Cavard, X., 2021. Forest Carbon Management: a Review of Silvicultural Practices and Management Strategies Across Boreal, Temperate and Tropical Forests. *Curr. For. Reports* 7, 245–266. <https://doi.org/10.1007/s40725-021-00151-w>.
- Arbogast Jr, C., 1953. Managing northern hardwoods in the Lake States. *United States Dep. Agric. For. Serv. Lake States For. Exp. Station*, p. 23.
- Archambault, L., Morissette, J., Bernier-Cardou, M., 1998. Forest succession over a 20-year period following clearcutting in balsam fir-yellow birch ecosystems of eastern Quebec. *Canada. For. Ecol. Manage.* 102, 61–74. [https://doi.org/10.1016/S0378-1127\(97\)00109-6](https://doi.org/10.1016/S0378-1127(97)00109-6).
- Ashton, M.S., Kelly, M.J., 2018. *The Practice of Silviculture: Applied Forest Ecology*. John Wiley & Sons Inc.
- Baker, F.S., 1934. *Theory and practice of silviculture*. McGraw-Hill Book Co., Inc., New York, NY.
- Barnes, B.V., Pregitzer, K.S., Spies, T.A., Spooner, V., 1982. Ecological Forest Site Classification. *J. For.* 80, 493–498.
- Beaudoin, A., Bernier, P.Y., Villemaire, P., Guindon, L., Guo, X.J., 2017. Species composition, forest properties and land cover types across Canada's forests at 250m resolution for 2001 and 2011. *Quebec Nat. Resour. Canada, Can. For. Serv. Laurentian For. Centre*.
- Beckley, T.M., 1998. Moving toward consensus-based forest management: A comparison of industrial, co-managed, community and small private forests in Canada. *For. Chron.* 74, 736–744. <https://doi.org/10.5558/tfc74736-5>.
- Bédard, S., Guillemette, F., Raymond, P., Tremblay, S., Larouche, C., DeBlois, J., 2014. Rehabilitation of Northern Hardwood Stands Using Multicohort Silvicultural Scenarios in Québec. *J. For.* 112, 276–286.
- Bédard, S., Majcen, Z., 2003. Growth following single-tree selection cutting in Québec northern hardwoods. *For. Chron.* 79, 898–906.
- Bliss, B.J.C., 2000. Public Perceptions of Clearcutting. *J. For.* 4–9.
- Blum, B.M. and Filip, S.M., 1963. A demonstration of Four Intensities of Management in Northern Hardwoods. *Res. Pap. NE-4*. 004.
- Bolton, N.W., D'Amato, A.W., 2011. Regeneration responses to gap size and coarse woody debris within natural disturbance-based silvicultural systems in northeastern Minnesota. *USA. For. Ecol. Manage.* 262, 1215–1222. <https://doi.org/10.1016/j.foreco.2011.06.019>.
- Bonan, G.B., 2008. Forests and climate change: forcings, feedbacks, and the climate benefits of forests. *Science (80-)* 320, 1444–1449.
- Bonnot, T.W., Cox, W.A., Thompson, F.R., Millsbaugh, J.J., 2018. Threat of climate change on a songbird population through its impacts on breeding. *Nat. Clim. Chang.* 8 <https://doi.org/10.1038/s41558-018-0232-8>.
- Bormann, F.H., Likens, G.E., 1979. Catastrophic disturbance and the steady-state in northern hardwood forests. *Am. Sci.* 67, 660–669.
- Bose, A.K., Weiskittel, A., Wagner, R.G., 2017. Occurrence, pattern of change, and factors associated with American beech-dominance in stands of the northeastern USA forest. *For. Ecol. Manage.* 392, 202–212. <https://doi.org/10.1016/j.foreco.2017.03.003>.
- Boucher, S., Crête, M., Ouellet, J.P., Daigle, C., Lesage, L., 2004. Large-scale trophic interactions: White-tailed deer growth and forest understorey. *Ecoscience* 11, 286–295. <https://doi.org/10.1080/11956860.2004.11682835>.
- Boulanger, Y., Puigdevall, P., 2020. Boreal forests will be more severely affected by projected anthropogenic climate forcing than mixedwood and northern hardwood forests in eastern Canada. *bioRxiv* In Press.
- Brewer, R., Merritt, P.G., 1978. Wind throw and tree replacement in a climax beech-maple forest. *Oikos* 149.
- Brockway, D.G., Loewenstein, E.F., Outcalt, K.W., 2015. Pro-B Selection Method for Uneven-Aged Management of Longleaf Pine Forests. *Proc. 17th Bienn. South. Silv. Res. Conf.* 534–544.
- Bromley, S.W., 1935. The original forest types of southern New England. *Ecol. Monogr.* 5, 61–89.
- Brown, M.L., Canham, C.D., Murphy, L., Donovan, T.M., 2018. Timber harvest as the predominant disturbance regime in northeastern U.S. forests: effects of harvest intensification. *Ecosphere* 9 (3). <https://doi.org/10.1002/ecs2.2062>.
- Bryant, R.C., 1917. Silviculture at Axton and in the Adirondacks Generally. *J. For.* 15, 801–810.
- Brydon-Williams, R., Munck, I.A., Asbjornsen, H., 2021. Incidence and ecology of the chaga fungus (*Inonotus obliquus*) in hardwood new england – acadian forests. *Can. J. For. Res.* 51, 122–131. <https://doi.org/10.1139/cjfr-2020-0144>.
- Burns, R.M., 1983. Silvicultural systems for the major forest types of the United States, U.S. Department of Agriculture, Forest Service. [https://doi.org/10.1016/0378-1127\(85\)90107-0](https://doi.org/10.1016/0378-1127(85)90107-0).
- Burns, R.M., Honkala, B.H., 1990. *Silvics of North America. Agric.* 654 (2), 877.
- Burrill, E.A., Wilson, A.M., Pugh, S.A., Menlove, J., Christiansen, G., Conkling, B.L., David, W., 2018. *The Forest Inventory and Analysis Database: database description and user guide version 8.0 for Phase 2*. U.S. Department of Agriculture Forest Service 946 9..
- Burtelow, A.E., Bohlen, P.J., Groffman, P.M., 1998. Influence of exotic earthworm invasion on soil organic matter, microbial biomass and denitrification potential in forest soils of the northeastern United States. *Appl. Soil Ecol.* 9, 197–202. [https://doi.org/10.1016/S0929-1393\(98\)00075-4](https://doi.org/10.1016/S0929-1393(98)00075-4).
- Butler, B.J., Butler, S.M., Caputo, J., Dias, J., Robillard, A., Sass, E.M., 2021. Family forest ownerships of the United States, 2018: results from the USDA Forest Service, National Woodland Owner Survey. *Gen. Tech. Rep. NRS-199*. Madison, WI U.S. Dep. Agric. For. Serv. North. Res. Station. 52 p. [plus 4 Append. 199.
- Butler, B.J., Hewes, J.H., Dickinson, B.J., Andrejczyk, K., Butler, S.M., Markowski-Lindsay, M., 2016. USDA Forest Service National Woodland Owner Survey. *Resour. Bull. NRS-99* 48.
- Butler, B.J., Leatherberry, E.C., 2004. America's Family Forest Owners. *J. For.* 102, 4–9.
- Callahan, M.A., González, G., Hale, C.M., Heneghan, L., Lachnicht, S.L., Zou, X., 2006. Policy and management responses to earthworm invasions in North America. *Biol. Invasions Belowgr. Earthworms as Invasive Species* 117–129. https://doi.org/10.1007/978-1-4020-5429-7_11.
- Canham, C.D., 2010. Different Responses to Gaps Among Shade-Tolerant Tree Species. *Ecology* 70, 548–550.
- Carnean, W.H., 1999. Site Quality and Site-Specific Management for White Birch, in: *Ecology and Management of White Birch*.
- Chambers, F.H., Beckley, T., 2003. Public involvement in sustainable boreal forests, in: *Towards Sustainable Management of the Boreal Forest*. p. 113.
- Champagne, E., Raymond, P., Royo, A.A., Speed, J.D.M., Tremblay, J.P., Côté, S.D., 2021. A Review of Ungulate Impacts on the Success of Climate-Adapted Forest Management Strategies. *Curr. For. Reports* 7, 305–320. <https://doi.org/10.1007/s40725-021-00148-5>.
- Churchill, D.J., Larson, A.J., Dahlgren, M.C., Franklin, J.F., Hessburg, P.F., Lutz, J.A., 2013. Restoring forest resilience: From reference spatial patterns to silvicultural prescriptions and monitoring. *For. Ecol. Manage.* 291, 442–457. <https://doi.org/10.1016/j.foreco.2012.11.007>.
- Clark, P.W., D'Amato, A.W., Evans, K.S., Schaberg, P.G., Woodall, C.W., 2022. Ecological memory and regional context influence performance of adaptation plantings in northeastern US temperate forests. *J. Appl. Ecol.* 59, 314–329. <https://doi.org/10.1111/1365-2664.14056>.
- Coates, K.D., Burton, P.J., 1997. A gap-based approach for development of silvicultural systems to address ecosystem management objectives. *For. Ecol. Manage.* 99, 337–354.
- Cogbill, C.V., Burk, J., Motzkin, G., 2002. The forests of presettlement New England, USA: Spatial and compositional patterns based on town proprietor surveys. *J. Biogeogr.* 29, 1279–1304. <https://doi.org/10.1046/j.1365-2699.2002.00757.x>.
- Costello, C.A., Yamasaki, M., Pekins, P.J., Leak, W.B., Neefus, C.D., 2000. Songbird response to group selection harvests and clearcuts in a New Hampshire northern hardwood forest. *For. Ecol. Manage.* 127, 41–54. [https://doi.org/10.1016/S0378-1127\(99\)00131-0](https://doi.org/10.1016/S0378-1127(99)00131-0).
- Crow, T.R., Buckley, D.S., Nauertz, E.A., Zasada, J.C., 2002. Effects of Management on Composition and Structure of Northern Hardwood Forests in Upper Michigan. *For. Sci.* 48, 129–145.
- Curtis, R., Rushmore, F., 1958. Some Effects of Stand Density and Deer Browsing on Reproduction in an Adirondack Hardwood Stand. *J. For.* 56, 116–121. <https://doi.org/10.1093/jof/56.2.116>.
- D'Amato, A.W., Bradford, J.B., Fraver, S., Palik, B.J., 2011. Forest management for mitigation and adaptation to climate change: Insights from long-term silviculture experiments. *For. Ecol. Manage.* 262, 803–816. <https://doi.org/10.1016/j.foreco.2011.05.014>.
- D'Amato, A.W., Catanzaro, P.F., Fletcher, L.S., 2015. Early regeneration and structural responses to patch selection and structural retention in second-growth northern hardwoods. *For. Sci.* 61, 183–189. <https://doi.org/10.5849/forsci.13-180>.
- D'Amato, A.W., Palik, B.J., 2021. Building on the last “new” thing: exploring the compatibility of ecological and adaptation silviculture. *Can. J. For. Res.* 51, 172–180.
- Dale, V.H., Joyce, L.A., McNulty, S., Neilson, R.P., Ayres, M.P., Flannigan, M.D., Michael Wotton, B., 2001. Climate Change and Forest Disturbances. *Bioscience* 51, 723–734.
- Dannevrolles, V., Arseneault, D., Bergeron, Y., 2016. Long-term compositional changes following partial disturbance revealed by the resurvey of logging concession limits in the northern temperate forest of eastern Canada. *Can. J. For. Res.* 46, 943–949. <https://doi.org/10.1139/cjfr-2016-0047>.
- DeGraaf, R.M., Yamasaki, M., 2003. Options for managing early-successional forest and shrubland bird habitats in the northeastern United States. *For. Ecol. Manage.* 185, 179–191. [https://doi.org/10.1016/S0378-1127\(03\)00254-8](https://doi.org/10.1016/S0378-1127(03)00254-8).
- DeSantis, R.D., Moser, W.K., Gormanson, D.D., Bartlett, M.G., Vermunt, B., 2013. Effects of climate on emerald ash borer mortality and the potential for ash survival in North America. *Agric. For. Meteorol.* 178–179, 120–128. <https://doi.org/10.1016/j.agrformet.2013.04.015>.
- Doyon, F., Gagnon, D., Giroux, J.F., 2005. Effects of strip and single-tree selection cutting on birds and their habitat in a southwestern Quebec northern hardwood forest. *For. Ecol. Manage.* 209, 101–116. <https://doi.org/10.1016/j.foreco.2005.01.005>.
- Drever, C.R., Peterson, G., Messier, C., Bergeron, Y., Flannigan, M., 2006. Can forest management based on natural disturbances maintain ecological resilience? *Can. J. For. Res.* 36, 2285–2299. <https://doi.org/10.1139/X06-132>.
- Dukes, J.S., Pontius, J., Orwig, D., Jeffrey, R.G., Vikki, L.R., Brazee, N., Cooke, B., Kathleen, A.T., Erik, E.S., Harrington, R., Ehrenfeld, J., Gurevitch, J., Lerdau, M., Stinson, K., Wick, R., Ayres, M., 2009. Responses of insect pests, pathogens, and invasive plant species to climate change in the forests of northeastern North America: What can we predict? *Can. J. For. Res.* 39, 231–248. <https://doi.org/10.1139/X08-171>.

- Dumroese, R.K., Williams, M.I., Stanturf, J.A., Clair, J.B.S., 2015. Considerations for restoring temperate forests of tomorrow: forest restoration, assisted migration, and bioengineering. *New For.* 46, 947–964. <https://doi.org/10.1007/s11056-015-9504-6>.
- Dupuis, S., Arseneault, D., Sirois, L., 2011. Change from pre-settlement to present-day forest composition reconstructed from early land survey records in eastern Québec. *Canada. J. Veg. Sci.* 22, 564–575. <https://doi.org/10.1111/j.1654-1103.2011.01282.x>.
- Dupuis, S., Danneyrolles, V., Laflamme, J., Boucher, Y., Arseneault, D., 2020. Forest Transformation Following European Settlement in the Saguenay-Lac-St-Jean Valley in Eastern Québec. *Canada. Front. Ecol. Evol.* 8 <https://doi.org/10.3389/fevo.2020.00257>.
- Duveneck, M.J., Scheller, R.M., 2015. Climate-suitable planting as a strategy for maintaining forest productivity and functional diversity Author (s): Matthew J. Duveneck and Robert M. Scheller Source : Ecological Applications , Vol. 25, No. 6 (September 2015), pp. 1653-1668 Publish. Ecol. Appl. 25, 1653–1668.
- Emery, M.R., 2002. Historical overview of nontimber forest product uses in the Northeastern United States. In: Jones, E.T., McLain, R.J., Weigand, J. (Eds.), *Nontimber Forest Products in the United States*. University Press of Kansas, Lawrence, KS, pp. 3–25.
- Eyre, F.H., 1980. Forest Cover Types, Forest CoverTypes of the United States and Canada. Washington, DC.
- Eyre, F.H., Neetzel, J.R., 1937. Applicability of the Selection Method in Northern Hardwoods. *J. For.* 35, 353–358(6).
- Eyre, F.H., Zillgitt, W.M., 1953. Partial cutting in northern hardwoods of the Lake States. *USDA For. Serv. Tech Bull.* 124 pp.
- Fahmy, S.H., Hann, S.W.R., Jiao, Y., 2010. Soils of New Brunswick: The Second Approximation. East. Canada Soil Water Conserv. Cent.
- Fairfax, S., Achterman, G., 1977. The Monongahela Controversy and the Political Process. *J. For.* 75, 485–487. <https://doi.org/10.1093/jof/75.8.485>.
- Falk, K.J., Elliott, K.A., Burke, D.M., Nol, E., 2010. Early seedling response to group selection harvesting in a northern hardwood forest. *For. Chron.* 86, 100–109. <https://doi.org/10.5558/tfc86100-1>.
- Ford, S.E., Keeton, W.S., 2017. Enhanced carbon storage through management for old-growth characteristics in northern hardwood-conifer forests. *Ecosphere* 8. <https://doi.org/10.1002/ecs2.1721>.
- Foster, D.R., 1992. *Land-Use History (1730–1990) and Vegetation Dynamics in Central New England, USA*. *Br. Ecol. Soc.* 80, 753–771.
- Franklin, J.F., Mitchell, R.J., Palik, B.J., 2007. Natural disturbance and stand development principles for ecological forestry. Newtown Square, PA. <https://doi.org/10.2737/NRS-GTR-19>.
- Franklin, J.F., Spies, T.A., Pelt, R. v., Carey, A.B., Thornburgh, D.A., Berg, D.R., Lindenmayer, D.B., Harmon, M.E., Keeton, W.S., Shaw, D.C., Bible, K., Chen, J., 2002. Disturbances and structural development of natural forest ecosystems with silvicultural implications, using Douglas-fir forests as an example. *For. Ecol. Manage.* 155, 399–423.
- Frelich, L.E., Lorimer, C.G., 1991. Natural Disturbance Regimes in Hemlock-Hardwood Forests of the Upper Great Lakes Region. *Ecol. Soc. Am.* 61, 145–164.
- Fuller, J.L., Foster, D.R., McLachlan, J.S., Drake, N., 1998. Impact of human activity on regional forest composition and dynamics in central New England. *Ecosystems* 1, 76–95. <https://doi.org/10.1007/s100219900007>.
- Gawler, S., Cutko, A.R., 2018. *Natural Landscapes of Maine: A Guide to Natural Communities and Ecosystems*, Revised an. ed. Maine Natural Areas Program, August, Maine.
- Giencke, L.M., Doviak, M., Mountrakis, G., Cale, J.A., Mitchell, M.J., 2014. Beech bark disease: spatial patterns of thicket formation and disease spread in an aftermath forest in the northeastern United States. *Can. J. For. Res.* 44, 1042–1050. <https://doi.org/10.1139/cjfr-2014-0038>.
- Gilbert, V.S., Jensen, A.M., 1958. A Management Guide for Northern Hardwoods in New England. U.S. Dep. Agric. For. Serv. Northeast. For. Exp. Stn.
- Godman, R.M., Krefting, L.W., 1960. Factors Important to Yellow Birch Establishment in Upper Michigan Author. *Ecology* 41, 18–28.
- Godman, R.M., Tubbs, C.H., 1973. Establishing even-age northern hardwood regeneration by the shelterwood method - a preliminary guide. *USDA For. Service Research P.* 13 pgs.
- Godman, R.M., Ywaney, H.W., Tubbs, C.H., 1990. *Acer saccharum* Marsh. Sugar maple., in: Burns, R.M., Honkala, B.H. (Eds.), *Silvics of North America: Volume 2*. Hardwoods. pp. 78–91.
- Gustafsson, L., Kouki, J., Sverdrup-Thygeson, A., 2010. Tree retention as a conservation measure in clear-cut forests of northern Europe: A review of ecological consequences. *Scand. J. For. Res.* 25, 295–308. <https://doi.org/10.1080/02827581.2010.497495>.
- Hall, B., Motzkin, G., Foster, D.R., Syfert, M., Burk, J., 2002. Three hundred years of forest and land-use change in Massachusetts. *USA. J. Biogeogr.* 29, 1319–1335. <https://doi.org/10.1046/j.1365-2699.2002.00790.x>.
- Halliday, E.D., 1937. *A Forest Classification For Canada*. Canada Dep. Mines Resour, Ottawa.
- Halofsky, J.E., Andrews-Key, S.A., Edwards, J.E., Johnston, M.H., Nelson, H.W., Peterson, D.L., Schmitt, K.M., Swanston, C.W., Williamson, T.B., 2018. Adapting forest management to climate change: The state of science and applications in Canada and the United States. *For. Ecol. Manage.* 421, 84–97. <https://doi.org/10.1016/j.foreco.2018.02.037>.
- Hannah, P.R., 1988. The Shelterwood Method in Northeastern Forest Types: a Literature Review. *North. J. Appl. For.* 5, 70–77.
- Hanson, J.J., Lorimer, C.G., 2007. *Forest Structure and Light Regimes Following Moderate Wind Storms : Implications for Multi-Cohort Management*. *Ecol. Soc. Am.* 17, 1325–1340.
- Hanson, J.J., Lorimer, C.G., Halpin, C.R., Palik, B.J., 2012. Ecological forestry in an uneven-aged, late-successional forest: Simulated effects of contrasting treatments on structure and yield. *For. Ecol. Manage.* 270, 94–107. <https://doi.org/10.1016/j.foreco.2012.01.017>.
- Helman, A.C., Kelly, M.C., Rouleau, M.D., Dickinson, Y.L., 2021. Preferences for Northern Hardwood Silviculture among Family Forest Owners in Michigan's Upper Peninsula. *J. For.* 119, 113–129. <https://doi.org/10.1093/jofore/fvaa051>.
- Henry, C.R., Walters, M.B., Finley, A.O., Roloff, G.J., Farinosi, E.J., 2021. Complex drivers of sugar maple (*Acer saccharum*) regeneration reveal challenges to long-term sustainability of managed northern hardwood forests. *For. Ecol. Manage.* 479, 118541 <https://doi.org/10.1016/j.foreco.2020.118541>.
- Herm, M., Verheyen, K., 2007. Legacies of the past in the present-day forest biodiversity: A review of past land-use effects on forest plant species composition and diversity. *Sustain. Divers. For. Ecosyst. An Interdiscip. Approach* 361–371. https://doi.org/10.1007/978-4-431-73238-9_1.
- Hicks, R.J., 1998. *Ecology and management of Central Hardwood forests*. Wiley, New York.
- Hocking, D.J., Babbitt, K.J., Yamasaki, M., 2013. Comparison of silvicultural and natural disturbance effects on terrestrial salamanders in northern hardwood forests. *Biol. Conserv.* 167, 194–202. <https://doi.org/10.1016/j.biocon.2013.08.006>.
- Hornbeck, J.W., Martin, C.W., Pierce, R.S., Bormann, F., Likens, G.E., Eaton, J.S., 1986. *The Northern Hardwood Forest Ecosystem: Ten Years of Recovery from Clearcutting*. *USDA For. Serv. North. Res. Stn. NE RP 596*, 34.
- Horsley, S.B., Marquis, D.A., 1983. Interference by weeds and deer with Allegheny hardwood reproduction. *Can. J. For. Res.* 13, 61–69. <https://doi.org/10.1139/x83-009>.
- Horsley, S.B., Stout, S.L., DeCalesta, D.S., 2003. White-Tailed Deer Impact on the Vegetation Dynamics of a Northern Hardwood Forest. *Ecol. Appl.* 13, 98–118.
- Houston, D.R., 1994. Major new tree disease epidemics: beech bark disease. *Annu. Rev. Phytopathol.* 32, 75–87.
- Houston, R., 1975. Beech Bark Disease: The Aftermath Forests Are Structured for a New Outbreak. *J. For.* 660–663 (4).
- Hupperts, S.F., Webster, C.R., Froese, R.E., Dickinson, Y.L., 2020. Forest Ecology and Management Seedling and sapling recruitment following novel silvicultural treatments in Great Lakes northern hardwoods. *For. Ecol. Manage.* 462, 117983 <https://doi.org/10.1016/j.foreco.2020.117983>.
- Iverson, L.R., Prasad, A.M., Low, 2001. Potential changes in tree species richness and forest community types following climate change. *Ecosystems* 4, 216–225. <https://doi.org/10.1007/s10021>.
- Iverson, L., Knight, K.S., Prasad, A., Herms, D.A., Matthews, S., Peters, M., Smith, A., Hartzler, D.M., Long, R., Almendinger, J., 2016. Potential Species Replacements for Black Ash (*Fraxinus nigra*) at the Confluence of Two Threats: Emerald Ash Borer and a Changing Climate. *Ecosystems* 19, 248–270. <https://doi.org/10.1007/s10021-015-9929-y>.
- Iverson, L., Prasad, A., Matthews, S., 2008. Modeling potential climate change impacts on the trees of the northeastern United States. *Mitig. Adapt. Strateg. Glob. Chang.* 13, 487–516. <https://doi.org/10.1007/s11027-007-9129-y>.
- Jacobs, R.D., 1974. Damage to northern hardwood reproduction during removal of shelterwood overstory. *J. For.* 72, 654–656.
- Janowiak, M.K., Nagel, L.M., Webster, C.R., 2008. *Spatial Scale and Stand Structure in Northern Hardwood Forests : Implications for Quantifying Diameter Distributions* 54, 497–506.
- Jenkins, J., 1995. Notes on the Adirondack Blowdown of July 15th , 1995, Wildlife Conservation Society Working Paper No. 5.
- Jensen, P.G., Demers, C.L., McNulty, S.A., Jakubas, W.J., Humphries, M.M., 2012. Marten and fisher responses to fluctuations in prey populations and mast crops in the northern hardwood forest. *J. Wildl. Manage.* 76, 489–502.
- Jensen, V.S., 1943. Suggestions for Management of Northern Hardwood Stands in the Northeast. *J. For.* 180–185.
- Johnson, A., 1984. Small-Woodlot Management by Single-Tree Selection : Results 1, 69–71 (3).
- Johnson, J.E., Haag, C.L., Bockheim, G.J., Erdmann, G.G., 1987. Soil-site relationships and soil characteristics associated with even-aged red maple (*Acer rubrum*) stands in Wisconsin and Michigan. *For. Ecol. Manage.* 21, 75–89.
- Jones, R.H., Nyland, R.D., Raynal, D.J., 1989. Response of American Beech Regeneration to Selection Cutting of Northern Hardwoods in New York. *North. J. Appl. For.* 6, 34–36(3).
- Jones, R.H., Raynal, D.J., 1988. Root Sprouting in American beech (*Fagus grandifolia*): Effects of Root Injury, Root Exposure, and Season. *For. Ecol. Manage.* 25, 79–90. [https://doi.org/10.1016/0378-1127\(88\)90121-1](https://doi.org/10.1016/0378-1127(88)90121-1).
- Keenan, R.J., 2015. Climate change impacts and adaptation in forest management : a review. *Ann. For. Sci.* 72, 145–167. <https://doi.org/10.1007/s13595-014-0446-5>.
- Keeton, W.S., 2006. Managing for late-successional/old-growth characteristics in northern hardwood-conifer forests. *For. Ecol. Manage.* 235, 129–142. <https://doi.org/10.1016/j.foreco.2006.08.005>.
- Kellison, R.C., Young, M.J., 1997. The bottomland hardwood forest of the southern United States. *For. Ecol. Manage.* 90, 101–115. [https://doi.org/10.1016/S0378-1127\(96\)03905-9](https://doi.org/10.1016/S0378-1127(96)03905-9).
- Kelty, M.J., Kittredge, D.B., Kyker-Snowman, T., Leighton, A.D., 2003. The Conversion of Even-Aged Stands to Uneven-Aged Structure in Southern New England. *North. J. Appl. For.* 20, 109–116.
- Kelty, M.J., Nyland, R.D., 1981. Regenerating Adirondack Northern Hardwoods By Shelterwood Cutting and Control of Deer Density. *J. For.* 79, 22–26.

- Kenefic, L.S., Nyland, R.D., 2006. Proceedings on the Conference on Diameter-Limit Cutting in Northeastern Forests, in: Kenefic, L.S. and R.D.N. (Ed.), Conference on Diameter-Limit Cutting in Northeastern Forests. USDA Forest Service, Northeastern Forest Experiment Station, Newtown Square, PA.
- Kern, Christel C, Burton, J.I., Raymond, P., D'Amato, A.W., Keeton, W.S., Royo, A.A., Walters, M.B., Webster, C.R., Willis, J.L., 2017. Challenges facing gap-based silviculture and possible solutions for mesic northern forests in North America. *For. Int. J. For. Res. For.* 1–14.
- Kern, C.C., G., E., L.S., K., Palik, B.J., Strong, T.F., 2014. Development of the selection system in northern hardwood forests of the Lake States: an 80-Year silviculture research legacy, in: USDA Forest Service Experimental Forests. Springer New York, pp. 631–650.
- Kern, C.C., Schoelch, M., Crocker, P., Fellman, D., Marsh, A., Mausel, D., Pecore, M., Phillippi, J., Waukau, R., Waupochick, A., 2017b. Group Opening Outcomes, Sustainable Forest Management, and the Menominee Nation Lands. *J. For.* 115, 416–424. <https://doi.org/10.5849/jof.2016.092>.
- Kern, C.C., Schwarzmann, J., Kabrick, J., Gerndt, K., Boyden, S., Stanovick, J.S., 2019. Mounds facilitate regeneration of light-seeded and browse-sensitive tree species after moderate-severity wind disturbance. *For. Ecol. Manage.* 437, 139–147. <https://doi.org/10.1016/j.foreco.2018.12.040>.
- Keyser, T.L., Loftis, D.L., 2013. Long-term effects of single-tree selection cutting on structure and composition in upland mixed-hardwood forests of the southern Appalachian Mountains. *Forestry* 86, 255–265. <https://doi.org/10.1093/forestry/cps083>.
- Kochenderfer, J.D., Kochenderfer, J.N., Warner, D.A., Miller, G.W., 2004. Preharvest manual herbicide treatments for controlling American Beech in Central West Virginia. *North. J. Appl. For.* 21, 40–49.
- Krasny, M.E., Digregorio, L.M., 2001. Gap dynamics in Allegheny northern hardwood forests in the presence of beech bark disease and gypsy moth disturbances. *For. Ecol. Manage.* 144, 265–274.
- Landry, G., Thiffault, E., Cyr, D., Moreau, L., Boulanger, Y., Dymond, C., 2021. Mitigation potential of ecosystem-based forest management under climate change: A case study in the Boreal-temperate forest ecotone. *Forests* 12. <https://doi.org/10.3390/f12121667>.
- Leak, W., Filip, S., 1977. Thirty-eight years of group selection in New England northern hardwoods. *J. For.* 641–643.
- Leak, W.B., 1980. Influence of Habitat on Silvicultural Prescriptions in New England. *Jour. For.*
- Leak, W.B., 1978. Relationship of Species and Site Index to Habitat in the White Mountains of New Hampshire. Broomall, PA.
- Leak, W.B., Gottsacker, J.H., 1985. New Approaches to Uneven-Age Management in New England. *North. J. Appl. For.* 2, 28–31.
- Leak, W.B., Solomon, D.S., DeBald, P.S., 1987. Silvicultural guide for Northern hardwood types in the Northeast (revised). United States Dep. Agric. For. Serv.
- Leak, W.B., Solomon, D.S., Filip, S.M., 1969. A Silvicultural Guide of Northern Hardwoods in the Northeast. USDA For. Serv. Res. Pap. NE-143.
- Leak, W.B., 2005. Effects of Small Patch Cutting on Sugar Maple Regeneration in New Hampshire Northern Hardwoods. *North. J. Appl. For.* 22, 68–70.
- Leak, W.B., Yamasaki, M., Holleran, R., 2014. Silvicultural Guide for Northern Hardwoods in the Northeast 46.
- Leffelman, L.J., Hawley, R.C., 1925. Studies of Connecticut hardwoods: the treatment of advance growth arising as a result of thinnings and shelterwood cuttings., Yale Univ. ed. Yale University, New Haven, CT.
- Lindsey, A.A., Crankshaw, W.B., Qadir, S.A., 1965. Soil Relations and Distribution Map of the Vegetation of Presettlement Indiana. *Am. J. Sociol.* 118, 676–727.
- Linn, E.R., 1918. Silvical Systems in Spruce in Northern New Hampshire. *J. For.* 16, 897–908.
- Looney, C.E., D'Amato, A.W., Palik, B.J., Slesak, R.A., 2017. Canopy treatment influences growth of replacement tree species in *Fraxinus nigra* forests threatened by the emerald ash borer in Minnesota. *USA. Can. J. For. Res.* 47, 183–192. <https://doi.org/10.1139/cjfr-2016-0369>.
- Lorimer, C.G., White, A.S., 2003. Scale and frequency of natural disturbances in the northeastern US: Implications for early successional forest habitats and regional age distributions. *For. Ecol. Manage.* 185, 41–64. [https://doi.org/10.1016/S0378-1127\(03\)00245-7](https://doi.org/10.1016/S0378-1127(03)00245-7).
- Lovett, G.M., Weiss, M., Liebhold, A.M., Homes, T.P., 2016. Nonnative forest insects and pathogens in the United States: Impacts and policy options. *Ecol. Appl.* 26, 1437–1455.
- Lull, H.W., 1968. A forest Atlas of the Northeast. Northeast. For. Exp. Stn. Upper Darb, 46pp.
- Luppold, W.G., Sendak, P.E., 2004. Analysis of the interaction between timber markets and the forest resources of Maine. *North. J. Appl. For.* 21, 135–143. <https://doi.org/10.1093/njaf/21.3.135>.
- Majcen, Z., Richard, Y., Ménard, M., 1984. Écologie et dendrométrie dans le sud-ouest du Québec.
- Marquis, D.A., 1975. Interim guide to regeneration of Allegheny hardwoods.
- Marquis, D.A., 1965. Regeneration of Birch and Associated Hardwood after Patch Cutting. U.S. Forest Service Research Paper NE-32, 15.
- Matthews, S.N., Iverson, L.R., 2017. Managing for delicious ecosystem service under climate change: Can United States sugar maple (*acer saccharum*) syrup production be maintained in a warming climate? *Int. J. Biodivers. Sci. Ecosyst. Serv. Manag.* 13, 40–52. <https://doi.org/10.1080/21513732.2017.1285815>.
- Mausel, D.L., Waupochick, A., Pecore, M., 2017. Menominee Forestry: Past, Present. *Future. Jour. For.* 115, 336–369.
- McCullough, D.G., Wiefelich, J.B., 2015. Beech bark disease in Michigan: spread of the advancing front and stand-level impacts., in: otter, KM, and B.C. (Ed.), Forest Health Monitoring: National Status, Trends and Analysis, 2014. General Technical Report SRS-209. Asheville, North Carolina, pp. 125–132.
- Mendell, B., 2016. From Cigar Tax to Timberland Trusts: A Short History of Timber REITs and TIMOS. *For. Hist.* today.
- Messier, C., Bauhus, J., Doyon, F., Maure, F., Sousa-Silva, R., Nolet, P., Mina, M., Aquilue, N., Fortin, M.J., Puettmann, K., 2019. The functional complex network approach to foster forest resilience to global changes. *Forest Ecosystem* 6. <https://doi.org/10.1186/s40663-019-0166-2>.
- Metzger, F., Schultz, J., 1984. Understorey Response to 50 Years of Management of a Northern Hardwood Forest in Upper Michigan. *Am. Midl. Nat.* 112, 209–223.
- Millar, C., Stephenson, N., Stephens, S., 2007. Climate Change and Forests of the Future: Managing in the Face of Uncertainty. *Ecol. Appl.* 17, 2145–2151.
- Miller, C.A., Graefe, A.R., 2001. Effect of harvest success on hunter attitudes toward white-tailed deer management in Pennsylvania. *Hum. Dimens. Wildl.* 6, 189–203. <https://doi.org/10.1080/108712001753461284>.
- Minckler, L.S., 1972. Hardwood Silviculture for Modern Needs. *J. For.* 70, 10–17.
- Moffat, K., Lacey, J., Zhang, A., Leopold, S., 2016. The social licence to operate: A critical review. *Forestry* 89, 477–488. <https://doi.org/10.1093/forestry/cpv044>.
- Murphy, P.A., Shelton, M.G., Graney, D.L., 1993. Group selection- problems and possibilities for the more shade-intolerant species, in: 9th Central Hardwood Forest Conference. West Lafayette, IN, pp. 229–247.
- Moorman, C.E., Russell, K.R., Greenberg, C.H., 2011. Reptile and amphibian response to hardwood forest management and early successional habitats. *Sustaining Young Forest Communities* 191–208.
- Nagel, L.M., Palik, B.J., Battaglia, M.A., Amato, A.W.D., Guldin, J.M., Swanston, C.W., Janowiak, M.K., Powers, M.P., Joyce, L.A., Millar, C.I., Peterson, D.L., Ganio, L.M., Kirschbaum, C., Roske, M.R., 2017. Adaptive Silviculture for Climate Change: A National Experiment in Manager-Scientist Partnerships to Apply an Adaptation Framework. *J. For.* 115, 167–178.
- Nesslage, Genevieve M, Porter, W.F., Nesslage, Genevieve M, Porter, W.F., 2001. Linked references are available on JSTOR for this article: A geostatistical analysis of deer harvest in the Adirondack Park, 1954-1997 29, 787–794.
- Neuendorff, J.K., Nagel, L.M., Webster, C.R., Janowiak, M.K., 2007. Stand structure and composition in a northern hardwood forest after 40 years of single-tree selection. *North. J. Appl. For.* 24, 197–202.
- Nichols, G.E., 1935. The Hemlock-White Pine-Northern Hardwood Region of Eastern North America. *Ecology* 16, 403–422.
- Nuttle, T., Royo, A.A., Adams, M.B., Carson, W.P., 2013. Historic disturbance regimes promote tree diversity only under low browsing regimes in eastern deciduous forest. *Ecol. Monogr.* 83, 3–17. <https://doi.org/10.1890/11-2263.1>.
- Nyland, R.D., 1999. Sugar Maple: Its Characteristics and Potentials. Horsley, Stephen B.; Long, Robert P., eds. Sugar maple Ecology. *Proc. an Int. Symp.* 1998 June 2-4; Warren, PA. Gen. Tech. Rep. NE-261. Radn. PA U.S. Dep. Agric. For. Serv. Northeast. Res. 1–13.
- Nyland, R.D., 2005. Diameter-Limit Cutting and Silviculture: A Comparison of Long-Term Yields and Values for Uneven-Aged Sugar Maple Stands. *Northern of Applied Forestry* 22, 111–116.
- Nyland, R.D., Bashant, A.L., Bohn, K.K., Verostek, J.M., 2006. Interference to hardwood regeneration in northeastern North America: ecological characteristics of American beech, striped maple, and hobblesh. *North. J. Appl. For.* 23, 53–61.
- Nyland, R.D., Bashant, A.L., Bohn, K.K., Verostek, J.M., 2001. Interference to Hardwood Regeneration in Northeastern North America: Controlling Effects of American Beech, Striped Maple, and Hobblesh Managing the Aftermath Forest.
- O'Hara, K.L., 2002. The historical development of uneven-aged silviculture in North America. *Forestry* 75, 339–346. <https://doi.org/10.1093/forestry/75.4.339>.
- O'Hara, K.L., 2016. What is close-to-nature silviculture in a changing world? *Forestry* 89, 1–6. <https://doi.org/10.1093/forestry/cpv043>.
- Oliver, C.D., 1981. Forest development in North America following major disturbances. *For. Ecol. Manage.* 3, 153–168. [https://doi.org/10.1016/0378-1127\(80\)90013-4](https://doi.org/10.1016/0378-1127(80)90013-4).
- Orwig, D.A., Abrams, M.D., 1994. Land-use history (1720–1992), composition, and dynamics of oak pine forests within the piedmont and coastal plain of northern Virginia. *Can. J. For. Res.* 24, 1216–1225. <https://doi.org/10.1139/x94-160>.
- Oswald, E.M., Pontius, J., Rayback, S.A., Schaberg, P.G., Wilmot, S.H., Dupigny-Giroux, L.A., 2018. The complex relationship between climate and sugar maple health: Climate change implications in Vermont for a key northern hardwood species. *For. Ecol. Manage.* 422, 303–312. <https://doi.org/10.1016/j.foreco.2018.04.014>.
- Palik, B.J., D'Amato, A.W., Franklin, J.F., Johnson, K.N., 2020. Ecological silviculture: Foundations and applications. Waveland Press.
- Parker, W.C., Colombo, S.J., Cherry, M.L., Flannigan, M.D., Greifenhagen, S., McAlpine, R.S., Papadopol, C., Scarr, T., 2000. Third millennium forestry: What climate change might mean to forests and forest management in Ontario. *For. Chron.* 76, 445–463. <https://doi.org/10.5558/tfc76445-3>.
- Patterson, W.A., Sassaman, K.E., 1988. Indian fires in the prehistory of New England. In: *Holocene Human Ecology in Northeastern North America*. Springer, Boston, MA, pp. 107–135.
- Pecore, M., 1992. Menominee sustained-yield management. *J. For.* 90, 12–16.
- Pederson, N., Dyer, J.M., Mcewan, R.W., Hessel, A.E., Cary, J., Orwig, D.A., Rieder, H.E., Cook, B.I., Pederson, N., Dyer, J.M., Mcewan, R.W., Hessel, A.E., Mock, C.J., Orwig, D.A., Rieder, H.E., Cook, B.I., 2014. The legacy of episodic climatic events in shaping temperate, broadleaf forests. *Ecol. Monogr.* 84, 599–620.
- Peters, M.P., Prasad, A.M., Matthews, S.N., Iverson, L.R., 2020. Climate change tree atlas, Version 4. Delaware, OH.
- Peterson, C.E., Maguire, D.A., 2004. Balancing ecosystem values: Innovative experiments for sustainable forestry - Foreword. *Balanc. Ecosyst. Values Innov. Exp. Sustain. For.* 635, 1. <https://doi.org/10.4067/S0717-65382006000300019>.

- Peterson, C.J., 2000. Catastrophic wind damage to North American forests and the potential impacts of climate change. *Sci. Total Environ.* 262.
- Pinto, F., Romaniuk, S., Ferguson, M., 2008. Changes to preindustrial forest tree composition in central and northeastern Ontario. *Canada. Can. J. For. Res.* 38, 1842–1854. <https://doi.org/10.1139/X08-034>.
- Pond, N.C., Froese, R.E., Nagel, L.M., 2014. Sustainability of Selection System in Northern Hardwood Forests 60, 374–381.
- Post, R.O.C. and B.W., 1962. Site indices. *Stn. Pap. No.* 171.
- Poznanovic, S.K., Webster, C.R., Bump, J.K., 2013. Maintaining mid-tolerant tree species with uneven-aged forest management: 9-year results from a novel group-selection experiment. *Forestry* 86, 555–567. <https://doi.org/10.1093/forestry/cpt025>.
- Price, D.T., Zimmermann, N.E., Van Der Meer, P.J., Lexer, M.J., Leadley, P., Jorritsma, I. M., Schaber, J., Clark, D.F., Lasch, P., McNulty, S., Wu, J., Smith, B., 2001. Regeneration in gap models: priority issues for studying forest responses to climate change. *Clim. Chang.* 51, 475–508.
- Puettmann, K.J., 2011. Silvicultural Challenges and Options in the Context of Global Change: “Simple” Fixes and Opportunities for New Management Approaches. *J. For.* 109, 321–331.
- Raymond, P., Bédard, S., 2017. The irregular shelterwood system as an alternative to clearcutting to achieve compositional and structural objectives in temperate mixedwood stands. *For. Ecol. Manage.* 398, 91–100. <https://doi.org/10.1016/j.foreco.2017.04.042>.
- Raymond, P., Bédard, S., Roy, V., Larouche, C., Tremblay, S., 2009. The irregular shelterwood system: review, classification, and potential application to forests affected by partial disturbances. *J. For.* 107, 405–413.
- Rentch, J.S., Schuler, T.M., Nowacki, G.J., Beane, N.R., Ford, W.M., 2010. Canopy gap dynamics of second-growth red spruce-northern hardwood stands in West Virginia. *For. Ecol. Manage.* 260, 1921–1929. <https://doi.org/10.1016/j.foreco.2010.08.043>.
- Rhoads, A.G., Hamburg, S.P., Fahey, T.J., Siccama, T.G., Hane, E.N., Battles, J., Cogbill, C., Randall, J., Wilson, G., 2002. Effects of an intense ice storm on the structure of a northern hardwood forest. *Can. J. For. Res.* 32, 1763–1775. <https://doi.org/10.1139/x02-089>.
- Rickenbach, M., Kittredge, D.B., 2009. Time and distance: Comparing motivations among forest landowners in New England, USA. *Small-scale For.* 8, 95–108. <https://doi.org/10.1007/s11842-008-9071-1>.
- Ristau, T.E., Stoleson, S.H., Horsley, S.B., deCalesta, D.S., 2011. Ten-year response of the herbaceous layer to an operational herbicide-shelterwood treatment in a northern hardwood forest. *For. Ecol. Manage.* 262, 970–979. <https://doi.org/10.1016/j.foreco.2011.05.031>.
- Roberts, M.R., Powell, G.R., MacDonald, J.E., 1980. Regeneration after clearcutting in the northern hardwood portion of the Nashwaq Experimental Watershed. *New Brunswick. Gen. Tech. Rep. NE 6*, 85.
- Robitaille, A., Saucier, J.P., 1998. Paysage régionaux du Québec Méridional. *Gouv. Québec, Ministère des Ressources Nat. Dir. la Gest. des Stock. For. Dir. des relations publiques.* Québec 213 p.
- Rowe, 1972. Forest Regions of Canada. *Can. For. Serv. Publ.* 1300, 172 pgs.
- Royo, A.A., Stout, S.L., 2017. REcognition, response, and recovery: Deer impact research in Allegheny hardwood forests., in: US Forest Service, N.R.S. (Ed.), SILVAH: 50 Years of Science -Management Cooperation. Proceedings of the Allegheny Society of American Foresters Training Session. Clarion, PA, pp. 26–36.
- Rudolphi, J., Jönsson, M.T., Gustafsson, L., 2014. Biological legacies buffer local species extinction after logging. *J. Appl. Ecol.* 51, 53–62. <https://doi.org/10.1111/1365-2664.12187>.
- Russell-Roy, E.T., Keeton, W.S., Pontius, J.A., Kerchner, C.D., 2014. Rehabilitation forestry and carbon market access on high-graded northern hardwood forests. *Can. J. For. Res.* 44, 614–627. <https://doi.org/10.1139/cjfr-2013-0437>.
- Rustad, L.E., Campbell, J.L., 2012. A novel ice storm manipulation experiment in a northern hardwood forest. *Can. J. For. Res.* 42, 1810–1818. <https://doi.org/10.1139/X2012-120>.
- Sage, R.W., Porter, W.F., Underwood, H.B., 2003. Windows of opportunity: white-tailed deer and the dynamics of northern hardwood forests of the northeastern US. *J. Nat. Conserv.* 10, 213–220. <https://doi.org/10.1078/1617-1381-00021>.
- Sands, B.A., Abrams, M.D., 2011. A 183-year history of fire and recent fire suppression impacts in select pine and oak forest stands of the Menominee Indian reservation. *Wisconsin. Am. Midl. Nat.* 166, 325–338.
- Sass, E.M., Butler, B.J., Markowski-Lindsay, M., 2020. Estimated Distribution of Forest Ownerships Across the Conterminous United States 2017, 2.
- Schaetzl, R.J., Frank, J.K., Miller, B.A., 2012. A taxonomically based ordinal estimate of soil productivity for landscape-scale analyses. *Soil Sci.* 177, 288–299.
- Schmoldt, D.L., Martin, G.L., Bockheim, J.G., 1985. Yield-based measures of northern hardwood site quality and their correlation with soil-site factors. *For. Sci.* 31 (1), 209–219.
- Schulte, L.A., Mladenoff, D.J., Crow, T.R., Merrick, L.C., Cleland, D.T., 2007. Homogenization of northern U.S. Great Lakes forests due to land use. *Landscape Ecol.* 22, 1105. <https://doi.org/10.1007/s10980-007-9112-8>.
- Seymour, R.S., 2004. Silviculture: Lessons from our past, thoughts about the future, in: B. P., L. L. (Eds.), Great Lake Silviculture Summit. St. Paul, MN, pp. 3–14.
- Seymour, R.S., 1994. The Northeastern Region. In: Barrett, J.W. (Ed.), *Regional Silviculture of the United States*. John Wiley & Sons Inc.
- Seymour, R.S., White, A.S., deMaynadier, P.G., 2002. Natural disturbance regimes in northeastern North America—evaluating silvicultural systems using natural scales and frequencies. *For. Ecol. Manage.* 155, 357–367. [https://doi.org/10.1016/S0378-1127\(01\)00572-2](https://doi.org/10.1016/S0378-1127(01)00572-2).
- Shigo, A., 1972. The Beech Bark Disease Today In The Northeastern U.S. *J. For.* 286–289.
- Smallidge, P.J., Chedzoy, B., Curtis, P., Sims, K., 2021. Evaluating the construction and effectiveness of slash walls at the perimeter of regeneration harvests to exclude deer. *For. Ecol. Manage.* 497, 119529. <https://doi.org/10.1016/j.foreco.2021.119529>.
- Smith, D.M., 1972. The Continuing Evolution of Silvicultural Practice. *J. For.* 70, 89–92.
- Smith, D.M., Larson, B.C., Kelty, M.J., Ashton, P.M.S., 1997. The practice of silviculture. John Wiley & Sons Inc.
- Spittlehouse, D.L., Stewart, R.B., 2003. Adaptation to climate change in forest management. *J. Ecosyst. Manag.* 4, 1–11.
- Spurr, S.H., 1981. Clearcutting on national forests. *Nat. Resour. J.* 21, 223–243.
- Stearns, F.W., 1997. Physical environment supporting Lake State forests. *Lake States Reg. For. Resour. Assess. Tech. Pap. USDA Fores 1–7*.
- Steelman, T.A., Ascher, W., 1997. Public Involvement Methods in Natural Resource Policy Making: Advantages, Disadvantages and Trade-Offs. *Policy Science* 30, 71–90. <https://doi.org/10.1023/A>.
- Stephanson, C.A., Coe, N.R., 2017. Impacts of beech bark disease and climate change on American beech. *Forests* 8. <https://doi.org/10.3390/f8050155>.
- Stout, S.L., Hoover, C.M., Ristau, T.E., 2006. Listening to Old Beech and Young Cherry Trees-Long-term in the Alleghenies. *Warren PA*.
- Sykes, M.T., Prentice, I.C., Cramer, W., Journal, S., Mar, N., Sykes, M.T., Prentice, I.C., Cramer, W., Systems, G., 1996. A Bioclimatic Model for the Potential Distributions of North European Tree Species Under Present and Future Climates. *J. Biogeogr.* 23, 203–233.
- Tanentzap, A.J., Bazely, D.R., Koh, S., Timciska, M., Haggith, E.G., Carleton, T.J., Coomes, D.A., 2011. Seeing the forest for the deer: Do reductions in deer-disturbance lead to forest recovery? *Biol. Conserv.* 144, 376–382. <https://doi.org/10.1016/j.biocon.2010.09.015>.
- The State of Canada's Forests Annual Report, 2019.
- Thiffault, N., Roy, V., 2011. Living without herbicides in Québec (Canada): Historical context, current strategy, research and challenges in forest vegetation management. *Eur. J. For. Res.* 130, 117–133. <https://doi.org/10.1007/s10342-010-0373-4>.
- Tierson, W.C., Patric, E.F., Behrend, D.F., 1966. Influence of White-Tailed Deer on the Logged Northern Hardwood Forest. *J. For.* 1, 801–805.
- Trosper, R.L., 2007. Indigenous influence on forest management on the Menominee Indian Reservation. *For. Ecol. Manage.* 249, 134–139. <https://doi.org/10.1016/j.foreco.2007.04.037>.
- Tabbs, C.H., 1977. Manager's Handbook for Northern Hardwoods in the North Central States. *USDA-Forest-Service-General-Technical-Report,-North-Central-Forest-Experiment-Station.* 1977, No. NC-39, 29 pp.; 2 pl.; 9 ref.
- Tabbs, C.H., Houston, D.R., 1990. *Fagus grandifolia* Ehrh. American beech, in: *Silvics of North America* 2. pp. 325–332.
- Tabbs, C.H., Jacobs, R.D., Cutler, D., 1983. Northern Hardwoods, in: Burns, R.M. (Ed.), *Silvicultural Systems for Major Forest Types of the United States*. U.S. Department of Agriculture Handbook No. 445, Washington, DC, pp. 121–127.
- Twery, M., Patterson, W.A.I., 1983. Effects of species composition and site factors on the severity of beech bark disease in western Massachusetts and the White Mountains of New Hampshire: a preliminary report, in: Proceedings, I.U.F.R.O. Beech Bark Disease Working Party Conference; 1982 September 26-October 8; Hamden, CT. Gen. Tech. Rep. WO-37. USDA Forest Service, Northeastern Forest Experiment Station, Washington, DC, pp. 127–133.
- Ullah, S., Moore, T.R., 2009. Soil drainage and vegetation controls of nitrogen transformation rates in forest soils, southern Quebec. *J. Geophys. Res. Biogeosciences* 114.
- Walters, B.F., Woodall, C.W., Russell, M.B., 2016. White-tailed deer density estimates across the eastern United States, 2008. <https://doi.org/https://doi.org/10.13020/D6G014>.
- Wang, Z., Nyland, R.D., 1993. Tree species richness increased by clearcutting northern hardwoods in central New York. *For. Ecol. Manage.* 57, 71–84.
- Webster, C.R., Dickinson, Y.L., Burton, J.I., Frelich, L.E., Jenkins, M.A., Kern, C.C., Raymond, P., Saunders, M.R., Walters, M.B., Willis, J.L., 2018. Promoting and maintaining diversity in contemporary hardwood forests: Confronting contemporary drivers of change and the loss of ecological memory. *For. Ecol. Manage.* 421, 98–108. <https://doi.org/10.1016/j.foreco.2018.01.010>.
- Webster, C.R., Lorimer, C.G., 2005. Minimum opening sizes for canopy recruitment of midtolerant tree species: A retrospective approach. *Ecol. Appl.* 15, 1245–1262. <https://doi.org/10.1890/04-0763>.
- Westveld, R.H., 1949. *Applied Silviculture in the United States*, 2nd ed. *Edward Brothers Inc., Ann Arbor, MI*.
- Williams, M.I., Dumroese, R.K., 2013. Preparing for climate change: Forestry and assisted migration. *J. For.* 111, 287–297. <https://doi.org/10.5849/jof.13-016>.
- Willis, J.L., Walters, M., Farinosi, E., 2016. Local seed source availability limits young seedling populations for some species more than other factors in northern hardwood forests. *For. Sci.* 62, 440–448. <https://doi.org/10.5849/forsci.15-143>.
- Willis, J.L., Walters, M.B., Gottschalk, K.W., 2015. Scarification and gap size have interacting effects on northern temperate seedling establishment. *For. Ecol. Manage.* 347, 237–247. <https://doi.org/10.1016/j.foreco.2015.02.026>.
- Winans, K.S., Whalen, J.K., Rivest, D., Cogliastro, A., Bradley, R.L., 2016. Carbon sequestration and carbon markets for tree-based intercropping systems in southern Quebec, Canada. *Atmosphere (Basel)* 7, 1–13. <https://doi.org/10.3390/atmos7020017>.
- Witmer, G.W., DeCalesta, D.S., 1991. The Need and Difficulty of Bringing the Pennsylvania Deer Herd Under Control. *Fifth East. Wildl. Damage Control Conf.* 130–137.
- Wood, D., Yanai, R., Allen, D., Wilmot, S., 2009. Sugar maple decline after defoliation by forest tent caterpillar. *J. For.* 107, 29–37. <https://doi.org/10.1093/jof/107.1.29>.
- Woods, K.D., Davis, M.B., 1989. Paleocology of Range Limits : Beech in the Upper Peninsula of Michigan. *Ecology* 70, 681–696.

Woolf, A., Roseberry, J.L., 1998. Deer Management : Our Profession's Symbol of Success or Failure? *Wildl. Soc. Bull.* 26, 515–521.

Yamasaki, M., Costello, C.A., Leak, W.B., 2014. Effects of Clearcutting , Patch Cutting , and Low-density Shelterwoods on Breeding Birds and Tree Regeneration in New Hampshire Northern Hardwoods. Res. Pap. NRS 26.

Zipper, C.E., Burger, J.A., Skousen, J.G., Angel, P.N., Barton, C.D., Davis, V., Franklin, J. A., 2011. Restoring forests and associated ecosystem services on Appalachian coal surface mines. *Environ. Manage.* 47, 751–765. <https://doi.org/10.1007/s00267-011-9670-z>.