

Fabaceae—Pea family

## *Delonix regia* (Bojer ex Hook.) Raf. flamboyan

J. A. Vozzo

Dr. Vozzo retired from the USDA Forest Service's Southern Research Station

**Other common names.** royal poinciana, flametree.

**Occurrence and growth habit.** A popular ornamental throughout the tropics, flamboyan—*Delonix regia* (Bojer ex Hook.) Raf.—is a small to medium-sized tree, typically 7 to 16 m high and up to 60 cm in diameter (Little and Wadsworth 1964). The champion Puerto Rican flamboyan, however, is 32 m high and 105 cm in diameter (Francis 1994). It grows well in moist soil derived from limestone, where it is common and reproduces well, but it is also tolerant of well-drained and somewhat droughty conditions (Francis and Liogier 1991). The species is briefly deciduous. Flamboyan has prominent buttresses and a broad, flat crown when grown in full sun. Its shallow but spreading root system limits the sites where it may be planted. The tree is susceptible to termites, shoot borers, and heart rot (Webb and others 1984). Although the genus is reported to have 3 species, flamboyan is the most cosmopolitan. A native of Madagascar, it has been planted in nearly every country in frost-free areas and is perhaps the most important flowering ornamental tropical tree of the world (Meninger 1962).

**Use.** This is a beautiful tree in form, shade, and flower. The flowers are predominantly red, although yellow and orange forms are cultivated; they are relatively short-lived as cut flowers. Trees remain in flower for several weeks, however. They are often seen planted along roadsides as living fence posts or as shade trees on both sides of the road that arch over the entire road. The wood is yellow-brown, weak, brittle, and soft, with a specific gravity of about 0.3. Although the species is not a good timber source, the wood is widely used as firewood. The legume (pod) is edible (Little and Wadsworth 1964; Meninger 1962; Webb and others 1984).

**Flowering and fruiting.** Showy flowers follow a dry season when the tree is almost leafless. The 5-pointed calyx is hairy and borne on racemes 15 to 25 cm long. Flowers are commonly red but may be white, yellow, orange, or yellow and vary from 8 to 25 cm across. Although flowers form after the dry season and during the wet season, they persist


during leaf emergence so that the crown appears feathery green while the colorful flowers are dominant. The hard legumes are 35 to 50 cm long, 6 cm wide, and 5 mm thick, and they hang tenuously on trees year-round. When mature, the legumes split into 2 parts lengthwise and are dark brown to black (Little and Wadsworth 1964); seeds (figures 1 and 2) are shed at that time. There are about 4,500 seeds/kg (2,040/lb) from Puerto Rican sources (Marrero 1949), whereas Colombian sources report only 2,000 to 3,000 seeds/kg (900 to 1,360/lb) (Navarette nd).

**Collection, extraction, and storage.** Pruning poles should be used to collect dark brown to black legumes. Legumes open naturally on trees after about 6 months. If unopened legumes are collected, they should be dried in the sun for 1 month; then the woody legumes should be forced open and the seeds removed. Seeds are relatively loosely attached in lateral grooves inside the legume. Dry seeds store very well in either open or closed containers and do not require refrigerated storage (Francis 1994). Seeds stored for 12 months at 26 °C germinated at 60% (Marrero 1949). Webb and others (1984) reported viability after 4 years storage but do not give germination rate or percentage germination.

**Figure 1**—*Delonix regia*, flamboyan: mature seed.




**Figure 2**—*Delonix regia*, flamboyan: longitudinal section through a seed.



**Germination.** Scarification—with either hot water, sulfuric acid, or abrasion—is required for germination. Millat-E-Mustafa (1989) recommends 90 °C water for 10 seconds followed by 24 hours of imbibition. A concentrated sulfuric acid soak for 0.5 to 5 hours improved germination for Duarte (1974), whereas a hot-wire scarification proved superior to other means described by Sandiford (1988). Seeds subjected to the various scarification treatments reported here had germination values superior to those of their respective controls. Within 8 days of fresh collections, expect 76% germination after 9 weeks.

**Nursery practice.** Seedlings (figure 3) are ready for outplanting after 3 to 4 months of growth in plastic nursery bags during the wet season. Saplings are also grown to 2 m, then “balled and burlaped” for large ornamental potted plants. Mature flowering and fruiting trees may be grown in 3 to 5 years in good sites (Francis 1994).

**Figure 3**—*Delonix regia*, flamboyan: seedling at 10 days after germination.



#### References

- Duarte O. 1974. Improving royal poinciana seed germination. *Plant Propagator* 20(1): 15–16.
- Francis JK. 1994. Personal communication. Rio Piedras, PR: USDA Forest Service, International Institute of Tropical Forestry.
- Francis JK, Liogier HA. 1991. Naturalized exotic tree species in Puerto Rico. Gen. Tech. Rep. SO-82. New Orleans: USDA Forest Service, Southern Forest Experiment Station. 12 p.
- Little EL Jr, Wadsworth FH. 1964. Common trees of Puerto Rico and the Virgin Islands. *Agric. Handbk.* 249. Washington, DC: USDA Forest Service: 176–177.
- Marrero J. 1949. Tree seed data from Puerto Rico. *Caribbean Forester* 10: 11–30.
- Menninger EA. 1962. *Flowering trees of the world*. New York: Hearthsides Press. 336 p.
- Millat-E-Mustafa M. 1989. Effect of hot water treatment on the germination of seed of *Albizia lebbek* and *Delonix regia*. *Bano-Biggyan-Patrika* 18(1/2): 63–64.
- Navarette EJ. [no date]. *Informacion basica y tratamientos pregerminativos en semillas forestales*. Colombia: Ministerio de Agricultura, Estacion Forestal la Florida. 28 p.
- Sandiford M. 1988. Burnt offerings: an evaluation of the hot-wire seed scarifier. *Commonwealth Forestry Review* 67(3): 285–292.
- Webb DB, Wood PJ, Smith JP, Henman GS. 1984. *A guide to species selection for tropical and sub-tropical plantations*. Trop. For. Pap. 15, 2nd ed. Oxford, UK: University of Oxford, Commonwealth Forestry Institute: 256 p.

Papaveraceae—Poppy family

## ***Dendromecon* Benth.**

### bushpoppy

W. Gary Johnson and Donald L. Neal

Mr. Johnson retired from the USDA Forest Service's National Seed Laboratory; Dr. Neal retired from the USDA Forest Service's Pacific Southwest Forest and Range Experiment Station


**Growth habit, occurrence, and use.** Bushpoppies (also known as treepoppies) are openly branched, evergreen shrubs from 0.6 to 2.5 m high, sometimes to 6 m. They have a woody base with gray or white shreddy-barked stems. The 2.5- to 10-cm-long leaves are mostly lanceolate, 3 to 8 times longer than wide (LHBH 1976). Environmental factors and shoot growth pattern affect leaf size (Bullock 1989). The 2 species considered here grow on dry chaparral slopes, ridges, and washes below 1,830 m. One species is found in California's Channel Islands and the other in the coastal range, from Sonoma County to the Sierra San Pedro Martir, Baja California, Mexico, and in the west foothills of the Sierra Nevada, from Shasta County to Tulare County (table 1). Bush-poppies rely on seed production to propagate. No lignotuber is formed on sprouts that appear after burning, so regrowth after fire is rare (Bullock 1989). The genus is useful for watershed protection (Sampson and Jespersen 1963) and for forage. Goats are especially fond of bushpoppies, and deer (*Odocoileus* spp.) and sheep eat the sprouts after fire.

**Flowering and fruiting.** Flowers are bisexual, yellow, showy, and solitary on stalks. At several locations, the shrubs first flowered in their second spring (Bullock 1989). Flowers appear in April through June and sometimes into August (Munz and Keck 1959). Bullock (1989) reports that the shrubs flower profusely from February through April in the Santa Monica Mountains. Several populations produce a

few flowers throughout the year. Fruits are linear, grooved capsules measuring 5 to 10 cm long, with 2 valves that separate incompletely at maturity. Ripe fruits (those that explode when grasped) may be collected in May, June, and July (Neal 1974). Fruits are dehiscent, scattering the seeds (figure 1) up to several meters from the shrub, and ants disperse the seeds, some below and others above the ground. Concentrations of seeds can be found around the entrances of harvester ant—*Pogonomyrmex* and *Veromessor* spp.—nests (Bullock 1989).

**Collection, cleaning, and storage.** The black seeds are almost spherical, 2 to 4 mm in diameter, with a slightly

**Figure 1**—*Dendromecon harfordii*, island bushpoppy: seeds.



**Table 1**—*Dendromecon*, bush-poppy: nomenclature and occurrence

Scientific name & synonym(s)	Common name(s)	Occurrence
<b><i>D. harfordii</i> Kellogg</b> <i>Dendromecon rigida</i> ssp. <i>harfordii</i> (Kellogg) Raven <i>D. rigida</i> ssp. <i>rhamnoides</i> (Greene) Thorne	<b>island bushpoppy,</b> Harford tree-poppy	Channel Islands, California
<b><i>D. rigida</i> Benth.</b>	<b>stiff bushpoppy,</b> tree-poppy	Central California to N Baja California

**Source:** Munz and Keck (1959).

pitted, hard, brittle testa. The seeds are dispersed by ants; the prominent caruncle is removed and used by the ants for food. The endosperm is oily, and the minute embryo rudimentary (Berg 1966) (figure 2). The mean number of seeds per fruit ranged from 2.9 to 10.7 in 14 populations (Bullock 1989). In 2 samples of cleaned seeds, purity was 77% and soundness was 97%. There were 92,400 to 114,400 seeds/kg (42,000 to 52,000/lb) (Neal 1974). Four other samples had purities of 99.4 to 99.9%, with an average of 99.4%, and 100,300 to 106,300 seeds/kg, with an average of 103,200/kg (45,600 to 48,300/lb, average 46,900/lb) (Vivrette 1996). Bullock found that seed weights varied greatly among the 14 populations studied, ranging from 10.1 to 15.8 mg (Bullock 1989). Vivrette reported seed weights ranging from 9.38 to 9.90 mg, average 9.70 mg, on 4 samples (Vivrette 1996). Bullock's slightly heavier fresh seeds may have had attached caruncles or a higher moisture content than Vivrette's laboratory samples.

There are no reports of seed storage of these species, but they likely can be stored at low moisture contents and near-freezing temperatures.

**Germination pretreatments.** Bushpoppy seeds have been sown in a moist medium at temperatures alternating diurnally from 4.5 °C (night) to 21 °C (day). Germination started after 50 days at these temperatures and reached 21% at 102 days after sowing (Mirov and Krabel 1939; Neal

1974). Vivrette reported no germination in of 9 samples tested for 21 days at 15 °C. A few seeds germinated on blotters moistened with 400 ppm GA<sub>3</sub> (gibberellic acid). Total viable seeds as determined by staining in tetrazolium chloride ranged from 11 to 50%, average 27% (Vivrette 1996).


Emery recommended fire treatment or 1 1/2 to 2 months of stratification and stated that 3 months of stratification with a diurnal fluctuation from 8 to 21 °C may improve germination (Emery 1988).

**Nursery practice.** Fire-treated bushpoppy seeds give the most reliable germination in nurseries (Emery 1988; Everett 1957). Seeds to be fire-treated should be sown in the fall in a slightly moist nurserybed. The seeds should be then covered with a layer of milled peat or sand 1 to 2 times as thick as the seeds' diameter and not watered. Then, a 10- to 15-cm (4- to 6-in) layer of dry pine needles or excelsior should be placed over the bed and burned. The seedbed should be watered after it has cooled. If wooden flats are being used, 2 layers of aluminum foil will protect the wood during the burning (Emery 1988).

## References

- Berg RY. 1966. Seed dispersal of *Dendromecon*: its ecologic, evolutionary and taxonomic significance. *American Journal of Botany* 53: 61-73.
- Bullock SH. 1989. Life history and seed dispersal of the short-lived chaparral shrub *Dendromecon rigida* (Papaveraceae). *American Journal of Botany* 76(10): 1506-1517.
- Emery D. 1988. Seed propagation of native California plants. Santa Barbara, CA: Santa Barbara Botanical Garden. 115 p.
- Everett PC. 1957. A summary of the culture of California plants at the Rancho Santa Ana Botanic Garden 1727-1950. Claremont, CA: Rancho Santa Ana Botanical Gardens. 223 p.
- LHBH [Liberty Hyde Bailey Hortorium]. 1976. *Hortus third: a concise dictionary of plants cultivated in the United States*. New York: Macmillan. 1290 p.
- Mirov NT, Kraebel CJ. 1939. Collecting and handling seeds of wild plants. For. Pub. 5. Washington, DC: USDA Civilian Conservation Corps. 42 p.
- Munz PA, Keck DD. 1959. *A California flora*. Berkeley: University of California Press. 1681 p.
- Neal DL. 1974. *Dendromecon rigida* Benth., stiff bushpoppy. In: Schopmeyer CS, tech. coord. *Seeds of woody plants in the United States*. Agric. Handbk. 450. Washington, DC: USDA Forest Service: 372.
- Sampson AW, Jespersen BS. 1963. *California range brushlands and browse plants*. Manual 33. University of California Extension Service, California Agricultural Experiment Station. 162 p.
- Vivrette NJ. 1996. Personal communication. Carpinteria, CA: Ransom Seed Laboratory.

**Figure 2**—*Dendromecon harfordii*, island bushpoppy: longitudinal section through a seed.



Ebenaceae—Ebony family

## *Diospyros* L.

### persimmon

David F. Olson, Jr., R. L. Barnes, and W. Gary Johnson

Drs. Olson and Barnes retired from the USDA Forest Service's Southeastern Forest Experiment Station;  
Mr. Johnson retired from the USDA Forest Service's National Seed Laboratory

**Growth habit, occurrence, and use.** Nearly 200 species of persimmons—*Diospyros* L.—are widely distributed, mostly in tropical regions. Only 2 persimmons—common persimmon and Texas persimmon—are native to the 48 contiguous states. Two others are grown in mild regions: Japanese persimmon for fruit and date-plum for root stock (table 1). Other persimmons are native to the tropical regions of the United States—*Diospyros hillebrandii* (Seem.) Fosberg and *D. sandwicensis* (A. DC.) Fosberg in Hawaii (Little and Skolman 1989) and *D. revoluta* Poir. and *D. sintonisii* (Krug & Urban) Standl. in Puerto Rico (Little and others 1974).

Common persimmon is a small to medium-sized deciduous tree, normally attaining a height of 9 to 18 m at maturity (Sargent 1965). It occurs in open woods and as an invader of old fields from Connecticut west through southern Ohio to eastern Kansas, and south to Florida and Texas (Sargent 1965). Common persimmon develops best in the rich bottom lands along the Mississippi River and its tributaries and in coastal river valleys. In these optimum habitats, common persimmon trees often attain heights of 21 to 24 m and diameters of 51 to 61 cm (Morris 1965).

In past years, persimmon wood was used extensively for weaver's shuttles, golf club heads, and other products requir-

ing hard, smooth-wearing wood (Olson and Barnes 1974). At present, such uses have diminished because of the use of laminates and other substitute materials.

The fruits are exceedingly astringent when green, but delicious when thoroughly ripe (Harlow and Harrar 1958); they are eaten by humans, animals, and birds. The common persimmon is a valuable honey plant and has been cultivated for its handsome foliage and fruit since 1629. Several varieties have been developed for fruit production (Harlow and Harrar 1958).

Texas persimmon is a shrub or small tree of south Texas and northeast Mexico, usually 1.8 to 3 m tall but sometimes reaching 12 m, with 4-cm-long leaves (Everitt 1984; LHBH 1976). The fruits are important wildlife food, but the shrub is considered as undesirable in rangelands of the Southwest (Everitt 1984).

Japanese persimmon (*kaki*) and date-plum are small persimmons from Asia grown commercially in the milder regions of the United States. Japanese persimmon grows to 14 m, with 18-cm-long leaves and large delicious fruit; many varieties are listed. Date-plum grows to 14 m, with leaves 13 cm long; it is often used as a rootstock for Japanese persimmon (LHBH 1976).

**Table 1**—*Diospyros*, persimmon: nomenclature and occurrence

Scientific name & synonym(s)	Common name(s)	Occurrence
<b><i>D. kaki</i> L.f.</b> <i>D. chinensis</i> Blume	<b>Japanese persimmon,</b> <i>kaki</i> , keg fir, date-plum	NE Asia & Japan
<b><i>D. lotus</i> L.</b> <i>D. japonica</i> Siebold & Zucc.	<b>date-plum</b>	NE Asia & Japan
<b><i>D. texana</i> Scheele</b>	<b>Texas persimmon,</b> black persimmon	SE Texas to central & trans-Pecos Texas
<b><i>D. virginiana</i> L.</b> <i>D. mosieri</i> Small	<b>common persimmon,</b> eastern persimmon	S Connecticut to SE Nebraska, S to Gulf of Mexico

Sources: LHBH (1976), Sargent (1965).


**Flowering and fruiting.** Male and female flowers are borne on different plants, but a few plants have bisexual flowers. The female flowers are solitary, with 4 to many staminodes. The male flowers are in cymes or clusters with 4 to many stamens. The fruits are juicy, 1- to 10-seeded berries with enlarged, persistent calyxes at the base (LHBH 1976).

Common persimmon has small, dioecious, axillary flowers borne after the leaves from March to mid-June, depending on the latitude (Little and Delisle 1962; Morris 1965; Olson and Barnes 1974; Radford and others 1964). Flowers are most common in April and May and are pollinated by insects.


The fruits are green before ripening and may vary in color when ripe from green, yellow, orange, and yellowish brown to dark reddish purple and black (Olson and Barnes 1974; Sargent 1965) (figure 1). The fruit is a 2- to 5-cm plumlike berry, glaucous and with a conspicuous, persistent calyx, that contains 3 to 8 seeds (Olson and Barnes 1974; Sargent 1965). The fruits ripen from September to November; the flat, brown seeds, about 15 mm long, are dispersed from the time of ripening until late winter (Little and Delisle 1962; Olson and Barnes 1974; Morris 1965; Radford and others 1964) (figures 1 and 2). The seeds are disseminated by birds and animals that feed on the fruits, and to some extent, by overflow water in low bottom lands (Morris 1965).

Seed bearing may begin at age 10, but the optimum seed-bearing age is 25 to 50 years (Little and Delisle 1962; Morris 1965; Olson and Barnes 1974). Good seedcrops are borne about every 2 years, with light crops in intervening years (Olson and Barnes 1974).

**Figure 1**—*Diospyros virginiana*, common persimmon: mature fruit and a single seed.



**Figure 2**—*Diospyros virginiana*, common persimmon: longitudinal section through a seed.



The small Texas persimmon flowers have 5 lobes. The black, globe-shaped fruits are 2.5 cm in diameter and have dark flesh (LHBH 1976). Animals and birds eat the fruits and disseminate the seeds (Everitt 1984).

Japanese persimmon flowers are yellowish white, about 2 cm long. Male flowers have 16 to 24 stamens and female flowers have 8 staminodes. The orange to reddish fruits are variable in shape, to 7.6 cm in diameter, with orange flesh (LHBH 1976).

Date-plum flowers are reddish to greenish, 7.5 mm long, with 4 lobes. Male flowers have 16 stamens. The small, yellow, globe-shaped fruits are 12.5 mm in diameter and turn blackish as they ripen (LHBH 1976).

**Collection of fruits; extraction and storage of seeds.** The fruits of common persimmon may be collected by picking them or shaking them from the trees as soon as they are ripe and soft in texture. They may also be picked from the ground after natural fall. If the fruits have started to dry, they should be softened by soaking in water (Myatt and others 1988). The seeds are easily removed by running the fruits with water through a macerator and allowing the pulp to float away or by rubbing and washing the pulp through 6.4-mm ( $1/4$ -in) mesh hardware cloth (Olson and Barnes 1974). For small quantities, ripe fruits can be placed in plastic bags and left until the pulp turns to juice, which can then be drained away before drying the seeds (Dirr and Heuser 1987).

After being cleaned, the seeds should be spread out to dry for a day or two. Spreading the seeds on screens to dry is common (Myatt and others 1988). Prolonged storage requires thorough drying. After the seeds are dried, they should be passed over a 9.9-mm (#25) screen on an air-



screen cleaner to remove trash and twigs. Use of a gravity table with high air may also be necessary (Myatt and others 1988). The seeds can then be safely stored in sealed dry containers at 5 °C (Engstrom and Stoeckler 1941).

One hundred kilograms (220 pounds) of fruit of the common persimmon will yield 10 to 30 kg (22 to 66 lbs) of cleaned seeds (Olson and Barnes 1974); the number of seeds per weight ranges from 1,460 to 3,880/kg (665 to 1,764/lb), with an average of 2,640 seeds/kg (1,200/lb) (table 2) (Aroeira 1962; Engstrom and Stoeckler 1941; Olson and Barnes 1974). Seedlots of 96% purity and 90% soundness have been obtained (Olson and Barnes 1974).

Japanese persimmon has about 3,400 seeds/kg (1,550/lb). Seeds stored at 0 °C at 45% moisture content retained the greatest viability after 18 months. Viability decreased rapidly as the seeds were dried, regardless of the speed of drying, with almost no germination at moisture contents below 10% (Kotobuki 1978). Date-plum has about 8,910 seeds/kg (4,040/lb).

**Pregermination treatments.** Natural germination of common persimmon usually occurs in April or May, but 2- to 3-year delays have been observed (Blomquist 1922; Olson and Barnes 1974). The main cause of the delay is the seed covering, which caps the radical, restricts the embryo, and causes a decrease in water absorption (Blomquist 1922). After removal of this cap, 100% germination was secured with mature seeds collected in the autumn (Blomquist 1922). Seed dormancy also can be broken by stratification in sand or peat for 60 to 90 days at 3 to 10 °C (Aroeira 1962; Crocker 1930; Olson and Barnes 1974; Thornhill 1968). Sulfuric acid scarification for 2 hours proved to be less effective in breaking dormancy than did stratification (Aroeira 1962).

Japanese persimmon does not have strong dormancy. Oh and others (1988) have shown that, although stratification was not essential, it improved germination. Rate of germination of date-plum seeds increased as the stratification length

increased to 10 weeks (Oh and others 1988). No pretreatment is needed to germinate Texas persimmon seeds (Vora 1989).

**Germination tests.** Germination of stratified common persimmon seeds was tested in sand or peat flats at diurnally alternating temperatures of 20 to 30 °C. Germinative energies ranging from 54 to 94% were obtained in 20 to 34 days; and germinative capacities at 60 days varied from 62 to 100% (Olson and Barnes 1974). Payne achieved 90% uniform germination on common persimmon and date-plum by stratifying the seeds for 60 to 90 days in wet vermiculite after lightly dusting them with a fungicide. Scratching the seedcoat can shorten the stratification period (Payne 1996).

Fresh Japanese persimmon seeds taken from ripe fruits and sown immediately germinate best. Germination ranged from 20 to 77% in a study of 18 cultivars with fresh seeds sown immediately (Dirr and Heuser 1987). Date-plum seeds germinated best without light at alternating 18 to 30 °C with 10 weeks stratification at 5 °C. Germination of seeds stratified for 2 weeks was increased by treating them with 500 ppm gibberellin (GA<sub>3</sub>) (Oh and others 1988). Fresh Texas persimmon seeds sown immediately after extraction germinated 33%. Germination was reduced with all other treatments (Dirr and Heuser 1987).

The tetrazolium chloride staining test is often used to estimate the viability of common persimmon and date-plum seeds due to the long stratification period needed to overcome dormancy. Clipping the radicle end of a seed with toenail clippers and soaking the seed for several days in water or 500 ppm GA<sub>3</sub> will soften it. Then it should be cut lengthwise to expose the embryo and storage tissue for staining.

**Nursery practice.** Common persimmon seeds may be fall-sown or stratified and sown in the spring. In Missouri, fall-sowing at a depth of 5 cm (2 in) is the normal practice, and seedbeds are mulched. Steavenson (Olson and Barnes 1974) reported a tree percent of 50%; an average tree percent of 25 to 33% is easily attainable. Seedlings of this

**Table 2**—*Diospyros*, persimmon: seed yield data


Species	Cleaned seeds/weight				Samples
	Range		Average		
	/kg	/lb	/kg	/lb	
<i>D. kaki</i>	3,015–3,790	1,370–1,720	1,550	3,400	2
<i>D. lotus</i>	—	—	8,910	4,040	1
<i>D. virginiana</i> *	1,460–3,880	665–1,765	2,640	1,200	—

Source: Olson and Barnes (1974).

\* Seed weight to fruit weight ratio (in kilograms/100 kg or pounds/100 lb) = 10 to 30.

species have a strong taproot (figure 3) and should be field-planted at the end of the first season. Root wrenching will cause the seedlings to form a compact, fibrous root system (Myatt and others 1988).

**Figure 3—*Diospyros virginiana*, common persimmon: seedling development at 4, 6, and 8 days after germination.**



### References

- Aroeira JS. 1962. Dormencia e conservacao de sementes de algumas plantas frutiferas. *Experientiae* 2: 541–609.
- Blomquist HL. 1922. Dormancy in seeds of persimmon. *Elisha Mitchell Scientific Society Journal* 38: 14.
- Crocker W. 1930. Harvesting, storage, and stratification of seeds in relation to nursery practice. Boyce Thompson Institute Professional Paper 1: 114–120.
- Dirr M, Heuser C Jr. 1987. The reference manual of woody plant propagation: from seed to tissue culture. Athens, GA: Varsity Press. 239 p.
- Engstrom HE, Stoeckler JH. 1941. Nursery practice for trees and shrubs suitable for planting on the prairie-plains. Misc. Pub. 434. Washington, DC: USDA. 159 p.
- Everitt JH. 1984. Germination of Texas persimmon seed. *Journal of Range Management* 37: 189–192.
- Harlow WM, Harrar ES. 1958. Textbook on dendrology. New York: McGraw-Hill. 561 p.
- Kotobuki K. 1978. Seed storage of Japanese persimmon, *Diospyros kaki*. In: Long term preservation of favourable germ plasm in arboreal crops. Ibaraki-Ken, Japan: Fruit Tree Research Station, Ministry of Agriculture and Forestry: 36–42.
- LHBH [Liberty Hyde Bailey Hortorium]. 1976. Hortus third: a concise dictionary of plants cultivated in the United States and Canada. New York: Macmillan. 1290 p.
- Little EL Jr, Delisle AL. 1962. Time periods in development: forest trees, North America. In: Biological handbook on growth. Washington, DC: Federation of American Societies for Experimental Biology: table 104.
- Little EL Jr, Skolmen RG. 1989. Common forest trees of Hawaii (native and introduced). Agric. Handbk. 679. Washington, DC: USDA Forest Service. 321 p.
- Little EL Jr, Woodbury RO, Wadsworth FH. 1974. Trees of Puerto Rico and the Virgin Islands. 2nd volume. Agric. Handbk. 449. Washington, DC: USDA Forest Service. 1024 p.
- Morris RC. 1965. Common persimmon (*Diospyros virginiana* L.). In: Silvics of forest trees of the United States. Agric. Handbk. 271. Washington, DC: USDA Forest Service: 168–170.
- Myatt A, Huffman G, Odell J Jr. 1988. Seed processing manual. Washington, OK: Oklahoma Department of Agriculture, Forestry Division.
- Oh JH, Kim SK, Ahn HK. 1988. Studies on seed germination of *Diospyros* species. *Journal of the Korean Society for Horticultural Science* 29(4): 297–303.
- Olson DF Jr, Barnes RL. 1974. *Diospyros virginiana* L., common persimmon. In: Schopmeyer CS, tech. coord. Seeds of woody plants in the United States. Agric. Handbk. 450. Washington, DC: USDA Forest Service: 373–375.
- Payne JA. 1996. Personal communication. Byron, GA: USDA Agricultural Research Service.
- Radford AE, Ahles HE, Bell CR. 1964. Guide to the vascular flora of the Carolinas. Chapel Hill: University of North Carolina Book Exchange.
- Sargent CS. 1965. Manual of the trees of North America. 2nd ed., corrected and reprinted. New York: Dover: 934 p.
- Thornhill HB. 1968. Propagation of woody ornamentals by seeds. *American Nurseryman* 127(6): 18, 86–89.
- Vora RS. 1989. Seed germination characteristics of selected native plants of the lower Rio Grande Valley, Texas. *Journal of Range Management* 42(1): 36–40.



Thymelaeaceae—Mezereum family

## *Dirca palustris* L.

### eastern leatherwood

John C. Zasada, David S. Buckley, Elizabeth Ann Nauertz, and Colleen F. Matula

Dr. Zasada retired from the USDA Forest Service's North Central Research Station; Dr. Buckley is an ecologist and Ms. Nauertz is a biological technician at the USDA Forest Service's North Central Research Station, Grand Rapids, Minnesota; Ms. Matula is a botanist at the USDA Forest Service's Ottawa National Forest, Bessemer Ranger District, Bessemer, Michigan


**Growth habit, occurrence, and use.** Eastern leatherwood—*Dirca palustris* L.—is also known as moosewood, rope-bark, and wicopy. Its natural distribution extends from New Brunswick to Ontario in the north and from northern Florida to Louisiana in the south (Fernald 1950). Within this range, the distribution is restricted to very specific site conditions. It is found almost exclusively in mesic, relatively rich hardwood forests or mixed conifer–hardwood forests (Alban and others 1991; Curtis 1959; Fernald 1950; Ferrari 1993; Jones 2000; Kotar and others 1988; Meeker and others 1993; Neveling 1962; Rooney 1996; Soper and Heimberger 1982; Voss 1985; Weir-Koetter 1996). In aspen ecosystems across the upper Great Lakes region, leatherwood is present in stands with a relatively high aspen site index and a significant northern hardwood component (Alban and others 1991). In Ontario, the northern limit of distribution is similar to that of beech (*Fagus grandifolia* Ehrh.) and sugar maple (*Acer saccharum* Marsh.) (Soper and Heimberger 1982). The distribution of plants on a particular site can vary from apparently random to aggregated (Jones 2000). Forests in which leatherwood is common are characterized by a dense overstory that permits relatively little light to reach the forest floor during the growing season. It is often the only true understory shrub in these stands; the other woody understory species are tolerant to mid-tolerant trees—for example, sugar maple, ironwood (*Ostrya virginiana* (Mill.) K. Koch.), white ash (*Fraxinus americana* L.), eastern hemlock (*Tsuga canadensis* (L.) Carr.), and balsam fir (*Abies balsamea* (L.) Mill.)—having the capacity to grow into the overstory (Alban and others 1991; Buckley 1996; Ferrari 1993; Jones 2000).

Western leatherwood—*D. occidentalis* Gray—is very similar to eastern leatherwood (Neveling 1962). Its distribution is limited to the wooded hills of the San Francisco Bay region (Vogelman 1953). Flower descriptions and morphological comparisons of the 2 species are provided by Vogelmann (1953). A related species in the Thymelaeaceae—*Daphne mezereum* L.—is an introduced species that has

become established in some areas. The information presented here is for eastern leatherwood; some of it may also apply to western leatherwood.

In its natural habitat, eastern leatherwood reaches a height of 3 to 4 m and basal diameters of 5 to 10 cm. Crown width and depth of larger plants can be as much as 2 to 3 m; the largest crown volumes that we have measured are in the range of 15 to 25 m<sup>3</sup>. Crown architecture can be fairly complex, with frequent branching and numerous apical growing points (figure 1). The largest individuals that we have observed are in old-growth northern hardwood forests where logging is prohibited and in older hardwood stands managed under a single-tree selection system. The maximum age attained by leatherwood is not known, but 30- to 50-year-old plants occur in older hardwood forests.

**Figure 1**—*Dirca palustris*, eastern leatherwood: mature forest-grown plant in full flower, with plant about 1.3 m tall.



Annual height growth varies considerably (Jones 2000). On mature plants, elongation of an individual apical meristem ranges from 1 to 25 cm, but cumulative annual growth over the many apical meristems comprising the crown may be 0.5 to 1 m or more. Complete removal or reduction of canopy cover to less than 50% seems to reduce the frequency of leatherwood, but more work is needed to understand effects of disturbance on the survival, growth, and reproduction of leatherwood. This reduction, however, may be more the result of physical damage during harvesting than to the change in the physical environment resulting from harvesting. In plants that have had branches completely or partially separated, callus growth covers the wound relatively quickly, giving wounded stems a distinct appearance. The flexible nature of the stem and branches is the result of a relatively low level of lignification in the wood (Neveling 1962). The specific gravity of the wood is 0.41, ranking it among the least dense woods of deciduous broadleaved species, comparable to poplar and basswood species (Alden 1995; Neveling 1962).

There is poor sprouting in leatherwood after the main stem is cut or broken. Layering of branches has been observed, but usually it does not occur, even on branches in good contact with an apparently suitable substrate layer. Seedling regeneration seems to be the most common way that the species is maintained in forests.


The only current documented use of eastern leatherwood is for landscaping. Even for this, it is not used to the extent possible, particularly in more northern areas (for example, in the northern Great Lakes region and northern New England), where the choice of plants is limited by climate. Although leatherwood provides a very early flowering, medium-sized shrub for these northern areas (Esson 1949), unfortunately its leaves are often infected by a rust and leaf miners in mid-summer, turning yellow prematurely and falling early. It can be planted and does best in moist, shaded areas. If the plant is naturally present in areas where development is planned, efforts should be made to protect it and provide conditions that favor growth, as older plants provide interesting form and structure to managed landscapes such as yards and gardens (del Tredici 1991; Dirr 1990). The plant appears to be browsed very little by deer (*Odocoileus* spp.), even in forests where other woody plants are repeatedly and severely browsed throughout the year (Weir-Koetter 1996). The lack of browsing could be due to the plants' diuretic qualities (Meeker and others 1993); the stem also contains large quantities of calcium oxalate crystals (Holm 1921). Ramsewak and others (1999) have described novel phenolic glycosides in leatherwood. The

strong pliable bark (the source of the common name) was used by Ojibwa for making bowstrings, baskets, and fishing lines (Holm 1921; Meeker and others 1993; Weir-Koetter 1996). The wood is very easy to slice with a sharp knife.

**Flowering and fruiting.** Leatherwood is monoecious. The pale yellow, fragrant flowers are perfect and borne in clusters of 2 to 7 (figures 1 and 2) (Neveling 1962; Soper and Heimberger 1982; Vogelmann 1953; Zasada and others 1996). The buds from which flowers develop are small and conical, with 4 distinct dark, silky scales that persist after flowering (figure 2). Clusters of 3 flowers are most common in northern Wisconsin–Michigan; clusters of 4 are somewhat less common, and clusters of 5 to 7 uncommon or rare. Mature plants commonly produce 300 to 1,500 flowers; the greatest number we observed on a plant was about 4,500 flowers (Zasada and others 1996). [Note: in the following discussion, we refer to unpublished data on fruits and seeds collected on or in the vicinity of the Ottawa National Forest in Michigan's Upper Peninsula and the Nicolet and Chequamegon National Forests in northern Wisconsin.]

Flowering occurs in April–May, 2 to 3 weeks before the overstory leaves out and generally before the spring ephemeral species flower. In 1994 and 1995 in northern Wisconsin–Michigan, fairly average years in terms of spring weather, pollination was mostly completed by May 11–15. In 1996, a relatively cold, wet spring, flowering began on May 14 on warm, south-facing aspects and several days later on north aspects. Flower buds opened as late as May 25 and some

**Figure 2**—*Dirca palustris*, eastern leatherwood: 3-flower cluster, subtending structures are silky bud scales.



anthers still contained pollen in early June. Flower parts drop quickly if pollination/fertilization is not successful but remain attached to developing fruits for a longer period (figure 3). Fruits ripen in June and July, with one report of ripening as late as September–October (Vogelman 1953). About 1 month (mid-June) after the first flowers appear in northern Wisconsin–Michigan, 75% of seeds contained embryos that filled 80% or more of the seed; 5% of the seeds were less than 50% filled. When fully ripe, the endosperm is a minor component of the seed (figure 4) (Neveling 1962; Zasada and others 1996). The outermost fleshy portion of the fruit cannot be separated easily from the seed coat until mid-late June.

Immature fruits are green and change to a very light green; some fruits are almost white when they fall from the plant. There are some reports that fruits turn reddish when mature (McVaugh 1941; Meeker and others 1993; Neveling 1962). However, McVaugh (1941) summarized the literature and concluded that although the reddish fruit color can be observed in dried herbarium specimens, his own and other's observations led to the conclusion that mature fruits are light to yellowish green. We found no evidence in the 8 stands studied in Wisconsin and Michigan that fruits were reddish in color when mature. The fleshy outer fruit wall (figure 5) of naturally dispersed fruits turns black within about 24 hours in some fruits, but in others it remains light green for several days.

Each flower has the potential to produce 1 single-seeded fruit, and hence fruits can be in clusters of 3 to 7 if all flow-

**Figure 3**—*Dirca palustris*, eastern leatherwood: fully developed but immature fruits. Flower parts are still attached to some fruits. Fruit length varies from 6.5 to 15 mm in the northern Wisconsin–Michigan area where fruits were collected.



**Figure 4**—*Dirca palustris*, eastern leatherwood: ripe fruits collected shortly after dispersal.



ers produce fruits. Fruits with 2 seeds were observed, but they were very rare. Clusters with 1 to 3 fruits were most common, as many flowers do not produce fruits. Clusters of more than 4 fruits are uncommon or rare. Number of flowers, fruit set, and number of fruits per cluster varies annually and among stands in the same year (table 1).

The fruit (figure 3) is a drupe and described as “bilaterally symmetrical, somewhat spindle-shaped....circular in cross-section at the widest point...and (having) a narrow, slightly elevated ridge...from the base of the style down the whole length of the fruit” (McVaugh 1941). Fruits are reported as 9 to 12 mm long for Ontario populations (Soper and Heimberger 1982), 12 to 15 mm long by 7 mm wide (Vogelman 1953) for Michigan and Indiana populations, and 12.5 to 15 mm long (McVaugh 1941) for New York populations. Average dimensions for fruits from 6 northern Wisconsin–Michigan populations were 8.5 to 9.5 mm long by 5.5 to 6.5 mm wide. Range in length was from 6.5 to 15.0 mm and width 4.5 to 7.5 mm for these latter populations (Zasada and others 1996). The fresh weight of individual fruits containing fully developed seeds varied from 0.08 to 0.23 g. Moisture content of whole fruits was 100 to 175% (dry weight basis) for dispersed fruits and those about to be dispersed.

Individual seeds, with their fleshy coats removed (figure 4), were 5.5 to 8.5 mm long and 3.5 to 5.0 mm wide for northern Wisconsin–Michigan populations. Fresh seed weight varied from 0.04 to 0.08 g and percentage moisture content was 40 to 55% (dry weight basis; seeds dried to a constant weight at 65 °C) for seeds collected from the ground shortly after dispersal; individual seeds on plants

**Table 1**—*Dirca palustris*, eastern leatherwood: flowering and fruit production in 2 forest types in northern Wisconsin


Stand type*	No. 3-flower clusters/shrub†		% cluster w/1–3 fruits	% of all clusters with fruits			Fruits/plant	
	Mean	Range		1 fpc	2 fpc	3 fpc	Mean	Range
Hardwood forest								
1995	153	5–515	23	22	48	30	40	0–386
1995	198	21–695	—	—	—	—	194	13–840
1996	—	—	55	20	41	19	—	—
Pine forest								
1995	59	0–255	2	0	77	23	3	0–45
1995	85	0–325	—	14	57	29	—	—
1996	—	—	23	—	—	—	36	0–260

**Note:** fpc = fruits per cluster; these populations did not have 4-flower or 4-fruit clusters in the 2 years of observation.

\* Based on 15 randomly selected shrubs in each stand.

† To obtain total number of flowers, multiply by 3.

**Figure 5**—*Dirca palustris*, eastern leatherwood: seeds with fleshy outer fruit wall removed. All seeds have a light-colored area along which the ovular trace is located; the 2 seeds in the center are positioned to show this area.



from which seeds were being dispersed but still firmly attached to the peduncle had moisture contents of 100 to 125% (Zasada and others 1996). Mature seeds are dark brown-black with a well-developed lighter longitudinal strip (figure 5). The strip is the point of attachment of the seed to the fruit wall in the area of the elevated ridge, which is a noticeable aspect of the shape of the fruit; the ovular trace is attached to the seed in this strip (figures 3 to 6) (McVaugh 1941; Neveling 1962).

Embryo length varied from 4 to 6 mm and from 2 to 4 mm in width in the Wisconsin–Michigan populations. At maturity, the embryo fills 95% or more of the seed; a small cavity develops at both poles of the seed (figure 6). Multiple embryos, all poorly developed, occurred in less than 0.5% of

the seeds (Zasada and others 1996).


The general anatomical features of a seed are illustrated in figure 6. The ovular trace and the pore through which it passes are an interesting feature of the seed (Neveling 1962). The pore appears filled with a fibrous material in seeds that have fallen from the plant. The black, stony seed-coat does not appear to completely seal the pore, even at maturity.

#### Collection of fruits; extraction and storage of seeds.

Fruits ripen in June and July. McVaugh (1941) observed that dispersal for plants in New York was completed by the following dates during a 3-year period—June 20, July 1–3, and later than July 7. In northern Wisconsin–Michigan populations, fruits have been observed on plants as late as July 16, but as observed by McVaugh (1941), dispersal was completed by July 1 in some years. The seeds disappear fairly quickly once they are fully mature but are fairly obvious on the soil surface for several days after dispersal (Durr and Heuser 1987; McVaugh 1941; Neveling 1962; Soper and Heimberger 1982). During windy periods, deposition rates were as high as 27 fruits/m<sup>2</sup>/hr under 1 shrub. Timing of fruit abscission and fruit drop vary among plants in a stand, among branches within a plant and among fruits in a cluster (Zasada and others 1996). In an area where dispersal was followed on a daily basis, seeds from the entire population were dispersed over about a 2-week period; some plants shed all of their seeds in 2 to 3 days, whereas others dispersed seeds over a 6- to 8-day period.

Birds do not seem to be a critical factor in seed removal, but they do consume some seeds and may be more important than our observations suggest. They may remove entire fruits, but, in some cases, they remove only the seed, leaving the fleshy fruit coat attached to the plant. Although the level of fruit use by rodents after dispersal is not known, it seems that this might be an important way in which seeds are

**Figure 6**—*Dirca palustris*, eastern leatherwood: generalized longitudinal section of mature seed, based on Neveling (1962) and Buckley (1996).



removed from the seed pool. Remnants of the black, stony seedcoat are fairly common under plants about 1 month after dispersal.

If seeds are needed, we recommend keeping a close watch on shrubs with fruits and collecting the fruits soon as they are ripe. Once some fruits fall naturally, all fruits have fully developed seeds. Embryo development is easily checked by cutting seeds longitudinally; those that are fully developed will appear as in figure 6.

Fruits can be picked by hand from the plant. However, when fully ripened, they readily fall when the plant is shaken and could be collected from the ground. Because each fruit contains only 1 seed, the number of seedlings desired (plus additional seeds as insurance against poor germination) will determine the number of fruits required. Based on cutting tests, 90 to 100% of developed fruits contained seeds with apparently viable embryos.

The pulp can be removed by hand or mechanically. When the seeds are fully mature, a cavity, with the exception of the attachment between the ovular trace and fruit

wall, develops between the fleshy fruit wall and the hard inner seed coat, making it fairly easy to hand-clean small quantities of seeds. The stony seedcoat is easily broken with the pressure of a fingernail and the seed can be squashed by squeezing between the thumb and forefinger with moderate pressure. Consequently, any type of mechanical cleaning must be done with care.

No information was found on the best ways to handle fruits or store seeds. Seeds remain viable in the forest floor from the time of dispersal until they germinate in the spring (del Tredici 1984), suggesting that storage for at least 8 to 10 months is possible. Seeds are exposed to a fairly wide range of temperature and moisture conditions between dispersal and germination.

**Germination.** Detailed information on the effects of environmental conditions on germination was not found. del Tredici (1984) used a number of standard methods to stimulate germination, but untreated seeds planted in a nursery bed soon after they were collected were the only ones that produced seedlings (67% germination). Dirr and Heuser (1987) reported that both cleaned and uncleaned (fleshy fruit wall removed) seeds produced seedlings. In controlled environment studies, a seedlot was observed to produce germinants over at least a 3-year period (Zasada and others 1996).

**Nursery.** Based on the limited information available, we recommend planting seeds soon after collection with and without the fleshy fruit wall in order to provide the range of conditions under which seeds appear to germinate naturally; seeds sown this way will germinate the next spring (del Tredici 1984; Dirr and Heuser 1987). Dirr and Heuser (1987) reported finding a number of young plants under a mature plant growing in a landscaped area, indicating that it might be possible to obtain some small seedlings from these situations. The growth rate of seedlings under open conditions is not documented. In its natural habitat, seedlings grow to a height of 20 to 30 cm in 5 to 10 years.

There has been little or no success in stimulating rooting in stem cuttings (Dirr and Heuser 1987). Layering occurs under natural conditions, suggesting that air-layering is a potential option for propagating leatherwood. However, Hendricks (1985) reported that air-layered stems did not produce roots or callus during an 8-week period. Our observations of layering of branches and the main stem under natural conditions suggest that it might take longer than 8 weeks for rooting to occur.

## References

- Alban DH, Perala DA, Jurgensen MF, Ostry ME, Probst JR. 1991. Aspen ecosystems properties in the upper Great Lakes. Res. Pap. NC-300. St. Paul: USDA Forest Service, North Central Forest and Range Experiment Station. 47 p.
- Alden HA. 1995. Hardwoods of North America. Gen. Tech. Rep. FPL-83. Madison, WI: USDA Forest Service, Forest Products Laboratory. 136 p.
- Curtis JT. 1959. The vegetation of Wisconsin. Madison: University of Wisconsin Press. 657 p.
- Buckley DS. 1996. Unpublished data. Rhinelander, WI: USDA Forest Service, North Central Research, Forestry Sciences Laboratory.
- del Tredici P. 1991. Propagating leatherwood: a lesson in humility. *Arnoldia* 51 (4): 62–66.
- Dirr MA, Heuser CW Jr. 1987. The reference manual of woody plant propagation: from seed to tissue culture. Athens, GA: Varsity Press. 239 p.
- Dirr MA. 1990. Manual of woody landscape plants: their identification, ornamental characteristics, culture, propagation, and uses. Champaign, IL: Stipes Publishing. 1007 p.
- Esson JG. 1949. Leatherwood for early spring bloom. *Journal of the New York Botanical Gardens* 50: 57–59.
- Fernald ML. 1943. The fruit of *Dirca palustris*. *Rhodora* 45: 117–119.
- Fernald ML. 1950. Gray's manual of botany. 8th ed. New York: American Book Co. 1632 p.
- Ferrari JB. 1993. Spatial patterns of litterfall, nitrogen cycling, and understory vegetation in a hemlock-hardwood forest [PhD thesis]. St. Paul: University of Minnesota. 180 p.
- Hendricks DR. 1985. Air layering native woody plants. *Proceedings of the International Plant Propagators' Society* 34: 528–531.
- Holm T. 1921. *Dirca palustris* L.: a morphological study. *American Journal of Science* 2: 177–182.
- Jones C. 2000. Growth and establishment of eastern leatherwood (*Dirca palustris*) in relation to forest management history [MS thesis]. Edwardsville: Southern Illinois University. 75 p.
- Kotar J, Kovach JA, Locey CT. 1988. Field guide to forest habitat types of northern Wisconsin. Madison: University of Wisconsin, Department of Natural Resources.
- McVaugh R. 1941. The fruit of eastern leatherwood. *Castanea* 6: 83–86.
- Meeker JE, Elias JE, Heim JA. 1993. Plants used by the Great Lakes Ojibway. Odanah, WI: Great Lakes Indian and Wildlife Commission. 440 p.
- Neveling LI. 1962. The Thymelaeaceae in the southeastern United States. *Journal of the Arnold Arboretum* 43: 428–434.
- Ramsewak RS, Nair MG, DeWitt DL, Mattson WG, Zasada JC. 1999. Phenolic glycosides from *Dirca palustris*. *Journal of Natural Products* 62: 1558–1561.
- Rooney SC. 1996. Personal communication. Houlton, ME: Consulting botanist.
- Soper JH, Heimberger ML. 1982. Shrubs of Ontario. Toronto: Royal Ontario Museum. 495 p.
- Vogelman H. 1953. A comparison of *Dirca palustris* and *Dirca occidentalis* (Thymelaeaceae). *Asa Gray Bulletin* 2(1): 77–82.
- Voss EG. 1985. Michigan flora. Part 2, Dicots. Bull. 59. Ann Arbor: University of Michigan, Cranbrook Institute of Science. 724 p.
- Weir-Koetter C. 1996. Of moosewood and blue beech. *Better Forests* (Winter 1996): 20–21.
- Zasada JC, Buckley D, Nauertz E. 1996. Unpublished data. Rhinelander, WI: USDA Forest Service, North Central Research Experiment Station, Forestry Sciences Laboratory.



Fabaceae—Pea family

## *Ebenopsis ebano* (Berl.) Barneby & Grimes

### Texas-ebony

Franklin T. Bonner

Dr. Bonner is a scientist emeritus at the USDA Forest Service's Southern Research Station, Mississippi State, Mississippi

**Synonyms.** *Pithecellobium flexicaule* (Benth.) Coult., *P. ebano* (Berl.) C.H. Muller, *Mimosa ebano* Berl., *Acacia flexicaulis* Benth.

**Other common names.** ebony blackbead, ape's earring.

**Growth habit, occurrence, and use.** *Ebenopsis* is a small genus, with only 2 species found in the United States. Texas-ebony occurs in Mexico and southern Texas and is the most valuable tree in the Rio Grande Valley. The species was formerly placed in the genus *Pithecellobium*; nomenclature of these species is discussed briefly in the *Pithecellobium* chapter of this book. The wood of Texas-ebony is used for furniture and fence posts, and the seeds can be used as a coffee substitute (they are boiled when green or roasted when ripe) (Vines 1960).

**Flowering and fruiting.** Texas-ebony flowers are yellow or cream-colored umbels about 4 cm in length borne in panicle clusters on the end of twigs. They appear from June to August (Vines 1960). The legumes (pods) turn from green to dark brown or black as they mature in the fall. They are flat, about 13 cm long, and 2.5 cm wide (figure 1). The legumes are also indehiscent and may remain on the trees for a year or more. The seeds are reddish brown, bean-shaped, and about 1.5 cm long (figures 2 and 3). Weights range from 1,550 to 1,990 seeds/kg (700 to 900/lb) (Walters and others 1974).

**Collection, extraction, and storage.** Legumes are usually picked by hand from the trees and air-dried in the sun. Seeds can be extracted by hand-flailing or by using mechanical macerators. Legume fragments can be removed with screens. There are no long-term storage data for Texas-ebony, but it is a typical hardseeded legume with orthodox storage behavior. Storing seeds for several years should be easy at low moisture contents (<10%) and temperatures of 2 to 5 °C (Walters and others 1974).

**Figure 1—** *Ebenopsis ebano*, Texas-ebony: legume.




**Figure 2—** *Ebenopsis ebano*, Texas-ebony: seeds.



**Germination.** The seedcoats of Texas-ebony are very hard, and few seeds will germinate without scarification. Soaking in sulfuric acid for 30 to 150 minutes has yielded germination of 78 to 88% (Alaniz and Everitt 1978; Vora 1989). This wide variation in soaking times suggests considerable variation in hardness of the seedcoats. In such cases, time trials should be carried out with small samples to choose the optimum soaking period for a given seedlot. Official seed testing organizations do not include Texas-ebony in their prescriptions for testing, but alternating temperatures of 15 and 30 °C have been quite successful following acid scarification (Alaniz and Everitt 1978).

**Figure 3**— *Ebenopsis ebano*, Texas-ebony: longitudinal section through a seed.



**Nursery practice.** There is little information on nursery practices for Texas-ebony. Nurserybed densities of 160 to 215/m<sup>2</sup> (15 to 20/ft<sup>2</sup>) appear to be suitable for raintree (*Albizia saman* (Jacq.) F. Muell.), a similar species (Walters and others 1974), and the same is suggested for Texas-ebony. Optimum planting depth in a greenhouse was reported to be 1 cm (0.4 in) (Alaniz and Everitt 1978). Direct seeding in old fields in Texas was improved by mulching the seeds with a commercial straw blanket (Vora and others 1988), and either mulching or shading would seem to be beneficial in nursery beds in that region.

#### References

- Alaniz MA, Everitt JH. 1978. Germination of Texas ebony seeds. *Journal of the Rio Grande Valley Horticultural Society* 32: 95–100. [Horticultural Abstracts 49(12): 9556; 1979].
- Little EL, Jr. 1979. Checklist of United States trees (native and naturalized). *Agric. Handbk.* 541. Washington, DC: USDA Forest Service. 375 p.
- Vines RA. 1960. Trees, shrubs, and woody vines of the Southwest. Austin: University of Texas Press. 1104 p.
- Vora RS. 1989. Seed germination characteristics of selected native plants of the lower Rio Grande Valley, Texas. *Journal of Range Management* 42: 36–40.
- Vora RS, Schumacher RW, Labus Z. 1988. Planting seeds of four south Texas native species under mulch cover. *Texas Journal of Science* 40: 456–458.
- Walters GA, Bonner FT, Petteys EQP. 1974. *Pithecellobium* Mart., blackbead. In: Schopmeyer CS, Tech. Cord. Seeds of woody plants in the United States. *Agric. Handbk.* 450. Washington, DC: USDA Forest Service: 639–640.

Elaeagnaceae—Oleaster family

## *Elaeagnus* L.

### elaeanus

David F. Olson, Jr., and Jill R. Barbour

Dr. Olson retired from the USDA Forest Service's Southeastern Forest Experiment Station; Ms. Barbour is a germination specialist at the USDA Forest Service's National Seed Laboratory, Dry Branch, Georgia

**Growth habit, occurrence, and use.** The genus *Elaeagnus* includes about 40 species of shrubs and trees, but there are only 3 species that are valuable for planting and for which reliable information is available (table 1). Although these deciduous trees and shrubs are grown often as ornamentals, they also produce edible fruits and serve as a source of wildlife food and as honey plants. Russian-olive is grown widely and has escaped from cultivation in many river lowland areas, particularly in the Great Plains, where it was extensively planted for shelterbelts (Olson 1974). In many areas, it has become invasive.

**Flowering and fruiting.** The fragrant, small, perfect flowers are borne in late spring (table 2) and are pollinated by insects (Mowry 1971). The fruit is a dry and indehiscent achene that is enveloped by a persistent fleshy perianth and hence is drupaceous (Jack 1969) (figures 1–3). The color of ripe fruit varies with the species (table 3). Seeds are often

distributed by birds following consumption of the ripe fruits (Turcek 1961).

**Collection of fruits; extraction and storage of seeds.** Ripe fruits are collected by picking them from the plants by hand or by beating or stripping them from the branches onto canvas or plastic sheets, usually from September to December (Olson 1974). Fruits may be spread out to dry or run through a macerator with water and the pulp floated off or screened out (Heit 1968; Olson 1974). Accordingly, commercial seedlots may consist of either dried fruits or cleaned stones. Dried fruits or cleaned stones at a moisture content from 6 to 14% can be stored successfully in sealed containers at 1 to 10 °C (Heit 1967; Mickelson 1968; Olson 1974; Peaslee 1969). Under ordinary storage conditions, seeds of silverberry remain viable for 1 to 2 years and those of Russian-olive up to 3 years (Olson 1974). The number of cleaned seeds (stones) per weight and other important yield

**Table 1**—*Elaeagnus*, elaeagnus: nomenclature and occurrence

Scientific name & synonym(s)	Common name(s)	Occurrence
<i>E. angustifolia</i> L. <i>E. hortensis</i> Bieb.	Russian-olive, oleaster, narrow-leafed oleaster	S Europe, W & Central Asia; Pacific Northwest to Minnesota, S through Great Plains to Mexico
<i>E. commutata</i> Bernh. ex Rydb. <i>E. argentea</i> Porsch, non Moench	silverberry, wolfberry	Quebec to Yukon, S to New Mexico, E to Nebraska
<i>E. umbellata</i> Thunb. <i>E. crispa</i> Thunb.	autumn-olive, autumn elaeagnus	China, Korea, & Japan; Maine to New Jersey & Pennsylvania, W to SW Minnesota, occasionally S to South Carolina


Sources: Fernald (1950), Harrington (1954), Olson (1974), Rehder (1940), Small (1933).

**Table 2**—*Elaeagnus*, elaeagnus: phenology of flowering and fruiting

Species	Location	Flowering	Fruit ripening	Seed dispersal	Seed size (mm)
<i>E. angustifolia</i>	—	June	Aug–Oct	All winter	12–13
<i>E. commutata</i>	Black Hills, South Dakota	June–July	Aug–Sep	Sep–Nov	8–9
<i>E. umbellata</i>	—	May–June	Aug–Oct	Sep–Nov	6–8

Source: Borell (1962), Dietz (1969), Hora (1981), McDermand (1969), Radford and others (1964), Rehder (1940).


**Figure 1**—*Elaeagnus angustifolia*, Russian-olive: fruit.



**Figure 2**—*Elaeagnus, elaeagnus*: achenes with fleshy perianth removed of *E. angustifolia*, Russian-olive (**left**) and *E. commutata*, silverberry (**right**).



**Figure 3**—*Elaeagnus angustifolia*, Russian-olive: longitudinal section through an achene enclosed in the fleshy perianth.



data are presented in table 4. From 4.5 kg (10 lb) of fruit, about 0.45 kg (1 lb) of cleaned seeds can be extracted. Fresh fruits of Russian-olive lost about 16 to 20% of their initial weight when air dried. The number of dried fruits per weight ranged from 3,970 to 9,900/kg (1,800 to 4,500/lb), with an average of 6,400/kg (2,900/lb). Purity of commercial seedlots for all 3 species has been high, ranging from 95 to 100% (Mickelson 1968; Olson 1974; Zarger 1968).

**Pregermination treatments.** Several pregermination treatments have been tested to overcome embryo dormancy in elaeagnus seeds. The most effective treatment is cold stratification at 1 to 10 °C for 10 to 90 days (Carroll 1971; Heit 1967, 1968; Lingquist and Cram 1967; Molberg 1969; Olson 1974). Stratification for less than 60 days is less effective than for longer periods (Carroll 1971). Intact autumn-olive seeds stratified at 5 °C from 2 to 6 weeks germinated less than 50% after 12 weeks at 25 °C, whereas seeds stratified for 10 to 14 weeks germinated completely in 12 weeks (Hamilton and Carpenter 1976). Allan and Steiner (1965) found that a 24-hour water soak followed by 45 days at 2 to 3 °C was sufficient to break dormancy in seeds of autumn-olive.

Russian-olive stones sometimes exhibit hard-seededness, and then should be soaked for 1/2 to 1 hour in sulfuric acid before germinating (Heit 1967). The optimum length of after-ripening for Russian-olive was reached at 12 weeks (Hogue and LaCroix 1970). Belcher and Karfalt (1979) found that snipping off 2 mm at the radicle end, after 7 days of water soaking, resulted in 96% germination. Snipping 2 mm at the cotyledon end only resulted in 50% germination. When 2 mm was snipped off both ends of the seeds, however, germination was 100%.

Seeds of Russian-olive that were not given a cold treatment but were soaked in Ethrel (2-chloroethyl phosphonic acid) germinated significantly better than seeds soaked in distilled water (Hamilton 1972). Concentrations of 300 and 600 ppm of Ethrel gave the maximum germination of 100 and 90%, respectively (Hamilton 1972). Germination was not further stimulated by giving the seeds 45 days of cold treatment before soaking in Ethrel (Hamilton 1972).

Gibberillic acid (GA<sub>3</sub>) applied to autumn-olive seeds at concentrations of 500 and 900 ppm decreased the time of cold stratification and increased the total germination percentage (Hamilton and Carpenter 1976). A coumarin-like inhibiting substance was found in all parts of the dormant and fully chilled seeds of Russian-olive (Hamilton and Carpenter 1976). Gibberillic acid at concentrations of 100 and 500 ppm and kinetin at 100 ppm appear to reverse the action of the inhibitor (Hamilton and Carpenter 1976).

Silverberry seeds, with endocarps removed, reached 85 to 100% germination within 10 days (Corns and Schraa

**Table 3**—*Elaeagnus, elaeagnus*: height; seed-bearing age, seedcrop frequency, and fruit ripeness criteria

Species	Height at maturity (m)	Year first cultivated	Minimum seed-bearing age (yr)	Years between large seedcrops	Fruit ripeness criteria	
					Preripe color	Ripe color
<i>E. angustifolia</i>	46–9	Long cultivated	≥3	3	Whitish to silvery l	Silver-gray outer; lemon-yellow inside
<i>E. commutata</i>	1.8–4.6	1813	—	1–2	Silvery green	Silver
<i>E. umbellata</i>	0.9–3.7	1830	6	—	Silvery, with	Red-pink brown scales

Sources: Borell (1962), Dietz (1969), Fernald (1950), Rehder (1940).

**Table 4**—*Elaeagnus, elaeagnus*: seed yield data

Species	Seed wt/fruit wt ratio	Cleaned seeds/weight				Samples
		Range		Average		
		/kg	/lb	/kg	/lb	
<i>E. angustifolia</i>	15–60	7,650–15,400	3,470–6,990	11,380	5,160	15
<i>E. commutata</i>	—	5,950–10,140	2,700–4,600	8,380	3,800	5
<i>E. umbellata</i>	5–10	46,525–84,670	21,100–38,400	62,180	28,200	30

Sources: Belcher and Washburn (1965), Carroll (1971), Harrington (1954), Heit (1970), Hinds (1967), McDermid (1969), Mickelson (1968), Molberg (1969), Mowry (1971), Olson (1974), Schumacher (1968), Zarger (1968).

1962). After intact seeds were stratified at 5 °C for periods of 40 to 110 days, the germination ranged from 23 to 75%, respectively (Corns and Schraa 1962). Supplemental treatments such as hot water soaks, gibberillic acid, and potassium nitrate (KNO<sub>3</sub>) soaks did not affect the germination of silverberry (Corns and Schraa 1962).

**Germination tests.** Some germination test results on stratified seeds are listed in table 5. Germination is epigeal. Silverberry had the best total germination (95 to 96%) and speed of germination after 60 to 90 days of stratification at 4 °C (Morgenson 1990). Seeds of silverberry used for strip mine reclamation yielded the highest germination (80%) after a 2-day warm (50 °C) water soak (Fung 1984). Results for autumn-olive seeds indicated that the optimum germination was achieved with cold stratification at 5 °C for 16 weeks and a night/day temperature of 10/20 °C (Fowler and Fowler 1987). Tests on excised embryos of Russian-olive have been completed in a very short time (Heit 1955). Belcher and Karrfalt (1979) found that it took 1 hour to completely excise the embryo from the seed and it resulted in 100% germination after 3 days incubation at 20 to 30 °C. Viability testing with 2,3,5-triphenyl tetrazolium chloride stain yielded 86% viable seeds for Russian-olive and 68 % viable seeds for autumn-olive (Olson 1974). Rules of the International Seed Testing Association (ISTA 1993) call for the use of tetrazolium staining for elaeagnus. Seeds should be soaked in water for 18 hours, then cut transversely at

both ends to open the embryo cavity. After 48 hours of soaking in 1% tetrazolium chloride, the seeds should be cut longitudinally to expose the embryos. The radicle tips and as much as one-third of the distal cotyledons can be unstained, and the seeds still considered viable. A secondary procedure calls for longitudinal cuts at the beginning.

**Nursery practice.** Seeds may be sown 13 to 25 mm (<sup>1</sup>/<sub>2</sub> to 1 in) deep in the late summer or fall without stratification, or in the spring after 10 to 90 days of cold stratification (Baker 1969; Growl 1968; Hinds 1967; Jack 1969; McDermid 1969; Mickelson 1968; Molberg 1969; Olson 1974; Zarger 1968). July seeding after 90 days of stratification gave excellent germination of Russian-olive in southeast Saskatchewan (Cram and Elliott 1966). In Michigan, autumn-olive is seeded by broadcasting 1.7 kg of fresh fruit/10 m<sup>2</sup> of bed area (1 lb/25 ft<sup>2</sup>) (Carroll 1971). At the Los Lunas Plant Material Center, Russian-olive is sown at a rate of 200 seeds, or 40 g (1.4 oz) of clean seeds/m, which yields 150 usable plants. In areas with a large population of mice, the pulp should be removed and cleaned seeds used for sowing (Carroll 1971). Russian-olive seedlings are susceptible to damage from rabbits and must be protected if these rodents are a problem.

Soil splash, which coats the pubescent leaves of newly emerged seedlings, is an important cause of mortality, and consequently, nursery beds should be mulched to cover the soil and prevent rain spattering (Carroll 1971; Growl 1968;

**Table 5**—*Elaeagnus*, *elaeanus*: germination test conditions and results for stratified seedlots\*

Species	Germination test conditions				Germinative energy		Germinative capacity	
	Medium	Temp (°C)		Days	Amt (%)	Days	Average	Samples
		Day	Night					
<i>E. angustifolia</i>	Sand	30	20	60	7–76	10–32	7–79	32
	Sand	—	—	21–40	—	—	54–90	11
	Moss	—	—	28	27	10	30	1
	Kimpak	30	20	28	—	—	68	19
<i>E. commutata</i>	Sand	30	20	50	52	13	60	1
<i>E. umbellata</i>	Kimpak	30	20	28	—	—	41	57
<i>E. umbellata</i>	—	30	10	—	—	—	93	—

Sources: Belcher and Washburn (1965), Heit (1968), Molberg (1969), Olson (1974).

\* Seeds were stratified for 10 to 90 days at 1.1 to 10 °C.

Hinds 1967; Mickelson 1968; Molberg 1969; Olson 1974; Zarger 1968). A seedling density of 130 to 320/m<sup>2</sup> (12 to 30/ft<sup>2</sup>) is desirable (Baker 1969; Molberg 1969; Zarger 1968). Stock usually is field planted as 1+0 or 2+0 seedlings, and grows well in most soils, including limestone or alkaline soils (Stoekeler 1946).

#### References

- Allan PF, Steiner WF. 1965. Autumn olive for wildlife and other conservation uses. Leaflet 458. Washington, DC: USDA.
- Baker LA. 1969. Correspondence. Elkton, OR: Dwight L. Phipps Forest Nursery.
- Belcher EV, Karrfalt RP. 1979. Improved methods for testing the viability of Russian olive seed. *Journal of Seed Technology* 4(1): 57–64.
- Borell AE. 1962. Russian-olive for wildlife and other conservation uses. Leaflet 517. Washington, DC: USDA. 8 p.
- Carroll DA. 1971. Personal communication. East Lansing, MI: USDA Soil Conservation Service, Rose Lake Plant Materials Center.
- Corns WG, Schraa RJ. 1962. Dormancy and germination of seeds of silverberry (*Elaeagnus commutata* Bernh.). *Canadian Journal of Botany* 40: 1051–1055.
- Cram WH, Thompson AC, Elliott T. 1966. Nursery cultural investigations: 1966 summary report for the tree nursery. Indian Head, SK: Canada Department of Agriculture, Prairie Farms Rehabilitation Administration. 35 p.
- Crowl JM. 1968. Personal communication. Earlinton, KY: Kentucky Reclamation Association.
- Fernald ML. 1950. *Gray's manual of botany*. 8th ed. New York: American Book Co. 1632 p.
- Fowler LJ, Fowler DK. 1987. Stratification and temperature requirements for germination of autumn olive (*Elaeagnus umbellata*) seed. *Tree Planters' Notes* 38(1): 14–17.
- Fung MYP. 1984. Silverberry seed pretreatment and germination techniques. *Tree Planters' Notes* 35(3): 32–33.
- Hamilton DF, Carpenter PL. 1976. Regulation of seed dormancy in *Elaeagnus angustifolia* by endogenous growth substances. *Canadian Journal of Botany* 54: 1068–1073.
- Hamilton DF, Carpenter PL. 1976. Factors influencing low-temperature after-ripening in seed of *Elaeagnus umbellata* Thunb. *Journal of the American Society of Horticultural Science* 10(2): 171–175.
- Harrington HD. 1954. *Manual of the plants of Colorado*. Denver: Saga Books. 666 p.
- Heit CE. 1955. The excised embryo method of testing germination quality of dormant seed. *Proceedings of the Association of Official Seed Analysts* 45: 108–117.
- Heit CE. 1967. Propagation from seed: 8. Fall planting of fruit and hardwood seeds. *American Nurseryman* 126(4): 12–13, 85–90.
- Heit CE. 1967. Propagation from seed: 11. Storage of deciduous tree and shrub seeds. *American Nurseryman* 126(10): 12–13, 86–94.
- Heit CE. 1968. Personal communication. Geneva, NY: New York State Agricultural Experiment Station.
- Heit CE. 1967. Propagation from seed: 6. Hard-seededness—a critical factor. *American Nurseryman* 125(10): 10–12, 88–96.
- Heit CE. 1968. Propagation from seed: 15. Fall planting of shrub seeds for successful seedling production. *American Nurseryman* 128 (4): 8–10, 70–80.
- Heit CE. 1970. Personal communication. Geneva, NY: Cornell University, New York State Agricultural Experiment Station.
- Hinds LW. 1967. Personal communication. Bismarck, ND: Lincoln-Oakes Nurseries.
- Hogue EJ, LaCroix LJ. 1970. Seed dormancy of Russian olive (*Elaeagnus angustifolia* L.). *Journal of the American Society of Horticultural Science* 95(4): 449–452.
- Hora B. 1981. *The Oxford encyclopedia of trees of the world*. Oxford: Oxford University Press. 288 p.
- ISTA [International Seed Testing Association]. 1993. Rules for testing seeds: rules 1993. *Seed Science and Technology* 21 (Suppl.): 1–259.
- Jack RA. 1969. Personal communication. Silverton, OR: Silver Falls Nursery.
- Lawrence GHM. 1951. *Taxonomy of vascular plants*. New York: Macmillan. 823 p.
- Lindquist CH, Cram WH. 1967. Propagation and disease investigations: summary report for the tree nursery. Indian Head, SK: Canadian Department of Agriculture, Prairie Farm Rehabilitation Administration. 42 p.
- McDermand J. 1969. Personal communication. Bismarck, ND: USDA Soil Conservation Service, Bismarck Plant Materials Center.
- Mickelson AS. 1968. Personal communication. Jonesboro, IL: Union State Tree Nursery.
- Molberg JM. 1969. Personal communication. Bottineau, ND: North Dakota School of Forestry.
- Morgenson G. 1990. Pregermination treatment and stratification of silverberry seed. *Tree Planters' Notes* 41(1): 24–25.
- Mowry CE. 1971. Personal communication. Elsberry, MO: USDA Soil Conservation Service.
- Müller H. 1883. *The fertilization of flowers*. Thompson DW, transl. & ed. London: Macmillan.
- Olson DF Jr. 1974. *Elaeagnus* L., *elaeanus*. In: Schopmeyer CS, tech. coord. *Seeds of woody plants in the United States*. Agric. Handbk. 450. Washington, DC: USDA Forest Service: 376–379.
- Peaslee AR. 1969. Personal communication. Lakin, WV: Clements State Tree Nursery.
- Radford AE, Ahles HE, Bell CR. 1964. *Guide to the vascular flora of the Carolinas*. Chapel Hill: University of North Carolina Book Exchange. 383 p.
- Rehder A. 1940. *Manual of cultivated trees and shrubs*. 2nd ed. New York: Macmillan. 996 p.
- Schumacher FW. 1968. Personal communication. Sandwich, MA.
- Small JK. 1933. *Manual of the southeastern flora*. New York: J.K. Small. 1554 p.
- Stoekeler JH. 1946. Alkali tolerance of drought-hardy trees and shrubs in the seed and seedling stage. *Minnesota Academy of Science Proceedings* 14: 79–83.
- Turcek FJ. 1961. *Ökologische Beziehungen der Vögel und Gehölze*. Bratislava: Verlag der Slowakische Akademie der Wissenschaften.
- Zarger TG. 1968. Personal communication. Norris, TN: Tennessee Valley Authority.



Asteraceae—Aster family

## *Encelia* Adans.

### encelia

Jane E. Rodgers and Carol Miller

Ms. Rodgers is at the USDI Point Reyes National Seashore, Point Reyes, California; Ms. Miller was a vegetation specialist at the USDI Joshua Tree National Park's Center for Arid Lands Restoration and Native Plants Nursery, Twentynine Palms, California

**Other common names.** brittlebush, bush-sunflower.

**Growth habit, occurrence and uses.** The brittlebush genus—*Encelia*—includes 14 species of low branching shrubs native to western America. The plants are suffrutescent, often with a pungent odor (Benson and Darrow 1954). Ray flowers (sometimes absent) are yellow, usually conspicuous when present, and produce neither pollen or fertile seeds. Disk flowers are yellow or purple (Benson and Darrow 1954). Species frequently hybridize, especially in disturbed areas. Species commonly found in the southwestern United States are listed in table 1.


The brittle wood secretes a clear resin used by Native Americans as a glue. In some parts of Mexico, the resin has been burned as incense for religious ceremonies (Benson and Darrow 1954). The Cahuilla of the southwestern United States have used gum from this plant as a medicine; the gum was heated and applied to the chest to relieve pain (Bean and Saubel 1972).

**Flowering and fruiting.** Flowering can begin in February and continue through July, weather conditions permitting. Most encelia flowerheads are yellow or a brown- or yellow-purple. The achenes are densely compressed, obovate or wedge-shaped, with edges that are long-ciliate and faces that are glabrous or short-hairy (figures 1 and 2) (Jepson 1993).

**Collection, extraction, and storage.** Timing of seed collection is critical, as the achenes are easily blown from

the plant after maturity (Kay and others 1977). Seeds may be hand-harvested and stored successfully for several years. Cleaning is difficult, for seeds are often mixed with dry flower and plant parts of similar size and weight. Studies on long-term storage of seeds of rayless encelia and Acton brittlebush showed good germination after 4 and 14 years, respectively (Kay and others 1988). Seeds of both species that were stored under 4 conditions (–15 °C, 4 °C, room temperature, and warehouse temperatures) also showed significantly poorer germination rates in the warehouse after 3 years. Storage conditions studied by Padgett and others (1999) showed that seedlots stored for 6 months in a stan-

**Figure 1**—*Encelia farinosa*, brittlebush: achenes.




**Table 1**—*Encelia*, encelia: nomenclature and occurrence

Scientific name	Common names	Occurrence
<i>E. californica</i> Nutt.	California brittlebush	Coastal California scrub to Baja California
<i>E. farinosa</i> Gray ex Torr.	brittlebush, incienso, goldenhills	Deserts of SW Utah, Arizona, & NW Mexico
<i>E. frutescens</i> (Gray) Gray	rayless encelia, green brittlebush	Deserts of S Nevada, W Arizona, & Baja California
<i>E. virginensis</i> A. Nels.	Virgin River encelia, brittlebush	E Mojave to SW Virgin River, Utah, & NW Arizona
<i>E. virginensis</i> var. <i>actonii</i> (Elmer) B.L. Turner	Acton brittlebush	SW California, SW Nevada to N Baja California

Source: Jepson (1993).

**Figure 2**— *Encelia farinosa*, brittlebush: longitudinal section through a seed.



dard refrigerator held at about 5 to 10 °C exhibited 2 to 3 times greater germination percentages than those stored at room temperature.

**Pregermination treatment and germination tests.**

No seed treatment is necessary (Emery 1988). Some research has been done to test dormancy in encelias, especially in brittlebush. At Joshua Tree National Park (JTNP), germination of brittlebush was tested with direct sowing and with the following seed treatments: 24 hours of cold water soaking, 6 hours of cold water soaking, and 24 hours of leaching, with all seeds sown on moist blotter paper (CALR 1993). Results showed very low germination rates (<1%). Research at the University of California at Riverside (UCR) has shown that the most significant cause of poor germination is lack of viable embryos in nearly half of the seeds tested for viability and germination behavior (Padgett and others 1999). Pre-soaking appears critical, and gibberellic acid (GA) has enhanced germination rates 2 to 3-fold (Padgett and others 1999). Treatment tests subjected seeds to warm water soaking for 30 minutes, followed by soaking in 100 ppm GA in water for 30 minutes. These treated seeds were then sown on or in 3 different media: UCR soil mix, vermiculite, and germination paper. All seeds were incubated in low light at 25 to 30 °C. Two general trends were observed: seed treatment with gibberellic acid for 30 minutes significantly increases germination rates, and sowing into vermiculite followed by transplantation into sterile potting medium appears to be the best method for seedling germination and survival. The vermiculite is pre-soaked, and misted every 2 days to avoid drying out. Seeds sown into the UCR soil mix had severe damping-off problems, and results

from the germination paper are thus far inconclusive. From the vermiculite, seedlings were successfully transplanted to larger containers and maintained in greenhouse conditions.

Work with Acton brittlebrush has correlated temperature and germination rates (Kay and others 1997):

<b>Temperature (°C)</b>	2	5	10	15	20	25	30	40
<b>Germination (%)</b>	0	1	47	65	55	5	7	0

Seeds were collected in late June 1973, yielding 1.96 kg of material of 24% purity. Cleaned seeds were 86% pure and had a pure fruit weight of 477,000 achenes/kg (216,721/lb). Emergence testing of encelia seedlots planted at 1-cm depth over a 10-day period showed a total emergence of 57%. Emergence at a 2-cm ( $\frac{3}{4}$  in.) depth was somewhat reduced and delayed, whereas no plants emerged from a 4-cm ( $\frac{1}{2}$  in.) depth (Kay 1975).

**Nursery practice.** Both cuttings and seedlings of brittlebush and Virgin River encelia have been successfully planted into 76-cm (30-in) tall and 15.2-cm (6-in) diameter tubes at JTNP using a mixture of sand, perlite, and mulch with a slow-release fertilizer. Both species were also grown in 3.8-liter (1-gal) pots, 15.1-liter (4-gal) pots, and plant bands, with the greatest outplanting success being the 76-cm (30-in) tubes or “tall pots” (CALR 1993). Plants in the nursery require a hardening-off period of 1 to 2 weeks and may be subject to aphid predation. Seedlings must be transplanted with care, for branches break off easily (CALR 1993).

**References**

- Bean LJ, Saubel KS. 1972. Temalpakh (from the earth): Cahuilla Indian knowledge and usage of plants. Morongo Indian Reservation, CA: Malki Museum Press. 225 p.
- Benson L, Darrow RA. 1954. The trees and shrubs of the southwestern deserts. Tuscon: University of Arizona Press. 437 p.
- CALR [Center for Arid Lands Restoration]. 1993. Seed test data filed 1989–1993. Twentynine Palms, CA: National Park Service, Joshua Tree National Park.
- Emery DE. 1988. Seed propagation of native California plants. Santa Barbara, CA: Santa Barbara Botanical Garden. 115 p.
- Jepson WL. 1993. The Jepson manual of higher plants of California. Hickman JC, ed. Berkeley: University of California Press. 1400 p.
- Kay BL. 1975. Test of seeds of Mojave Desert shrubs. Prog. Rep. BLM Contract 53500-CT4-2(N). 24 p.
- Kay BL, Brown CR, Graves WL. 1977. Virgin River encelia. Mojave Reveg. Notes 4. Davis: University of California Department of Agronomy and Range Science.
- Kay BL, Graves WL, Young JA. 1988. Long-term storage of desert shrub seed. Mojave Reveg. Notes 23. Davis: University of California Department of Agronomy and Range Science.
- Padgett P, Vazquez L, Allen EB. 1999. Seed viability and germination behavior of the desert shrub *Encelia farinosa*. Madroño 46(3): 126–133

Fabaceae—Pea family

## *Enterolobium cyclocarpum* (Jacq.) Griseb.

guanacaste or earpod-tree

John K. Francis

Dr. Francis retired from the USDA Forest Service's International Institute of Tropical Forestry

**Growth habit, occurrence, and use.** At maturity, the fast-growing *guanacaste* is a huge, spreading tree with feathery, bipinnately compound leaves. The trunks of open-grown trees are short and thick, tipped with an inverse cone-shaped crown; trunks of trees growing in closed stands have much longer boles. Guanacaste grows in both acid and alkaline soils (Bauer 1982) in forests and savannas from central Mexico (23°N) through Central America to about the Equator in northern Brazil (Little and others 1974; Pennington and Sarukhan 1968). However, the species has been widely planted in tropical and subtropical areas, including Puerto Rico, the U.S. Virgin Islands, Florida, and Hawaii (Francis 1988). Guanacaste is recommended for planting in areas that receive from 750 to 2,000 mm of mean annual precipitation (Bauer 1982; Fournier and Herrera 1977). Dry seasons of 1 to 6 months are normal in the native range (Bauer 1982; Janzen 1983). Guanacaste is principally used as an ornamental and shade tree in parks, estates, and broad avenues. It is also valued as a pasture shade tree, especially in Central America. Cattle, horses, and goats feed heavily on its sweet legumes (pods). The heartwood has a rich brown color and is in demand for cabinetry, furniture, crafts, and construction (Chudnoff 1984; Guridi 1980).

**Flowering and fruiting.** Small white flowers are borne in clusters or heads at the base of leaves (Little and others 1974; Pennington and Sarukhan 1968). Flowering takes place in March and April during the regrowth of new leaves after the leafless dry season (Hughes and Styles 1984; Janzen 1983). There is no indication in the literature as to the age at which flowers first appear; however, trees in a 26-year-old plantation in Puerto Rico had not yet flowered. The fruits are shiny, dark brown legumes that curve around one edge, giving them a shape that resembles a human ear (figure 1). Legumes are 7 to 12 cm in diameter and contain 8 to 18 seeds (Holdridge and Poveda 1975; Janzen 1983; Pennington and Sarukhan 1968); they mature the year they are formed and fall in March and April.


**Collection, cleaning, and storage.** Seeds can be collected in quantity by picking up the legumes after they have fallen to the ground. They can be separated by macerating the legumes and then washing them to remove the sticky syrup or by picking the seeds from the tough legumes with the point of a knife (Francis 1988). One thousand to a few thousand seeds are produced per tree. The 1.3- to 1.9-cm ( $\frac{1}{2}$ - to  $\frac{3}{4}$ -in) seeds (figure 2) number 1,100/kg (500/lb) (Janzen 1983; Neal 1965). The seeds store well according to Bauer (1982).

**Germination.** Without scarification, a moderate percentage of the seeds germinate over a span of several

**Figure 1**—*Enterolobium cyclocarpum*, guanacaste: seeds and legume.



**Figure 2**—*Enterolobium cyclocarpum*, guanacaste: longitudinal section through a seed.



months. Scarified seeds in most kinds of soil germinate in 3 to 7 days. Germination values of 79 and 84% were observed in tests in Puerto Rico (Francis and Rodríguez 1993) and Costa Rica (Salazar 1985). A seed can be scarified by nicking it on a grindstone, by cracking it with a pair of pliers, or by immersing it briefly in boiling water. In nature, seeds are scarified and disseminated primarily when the legumes are eaten by domestic and wild ungulates or when they are nicked by rodents (Janzen 1983). Germination is epigeal (Francis 1988).

**Nursery practice.** Seeds may be sown in nursery beds, germination trays, or directly in pots. They should be covered with about 1 cm (.4 in) of soil or potting mix. Seedlings develop rapidly in full sunlight, reaching plantable size of about 0.5 m (20 in) in about 6 months (Francis 1988). The seedlings are very drought hardy and generally, good survival at outplanting can be obtained with potted seedlings and stump plants (Bauer 1982). A test of container seedlings in southeastern Mexico yielded an average of 77% survival (Beroni and Juarez 1980).

#### References

- Bauer J. 1982. Especies con potencial para la reforestación en Honduras, resúmenes. Tegucigalpa, Honduras: Corporación Hondureña de Desarrollo Forestal. 42 p.
- Bertoni VR, Juarez VM. 1980. Comportamiento de nueve especies forestales tropicales plantadas en 1971 en el campo experimental "El Tormento". *Revista Ciencia Forestal* 25(5): 4-39.
- Chudnoff M. 1984. *Tropical timbers of the world*. Agric. Handbk. 607. Washington, DC: USDA Forest Service. 466 p.
- Fournier LA, Herrera de Fournier ME. 1977. La sucesión ecológica como un método eficaz para la recuperación del bosque de Costa Rica. *Agronomía Costarricense* 1(1): 23-29.
- Francis JK. 1988. *Enterolobium cyclocarpum* (Jacq.) Griseb., guanacaste, ear-pod tree. Res. Note SO-ITF-SM-15. New Orleans: USDA Forest Service, Southern Forest Experiment Station. 4 p.
- Francis JK, Rodríguez A. 1993. Seeds of Puerto Rican trees and shrubs: second installment. Res. Note SO-374. New Orleans: USDA Forest Service, Southern Forest Experiment Station. 5 p.
- Guridi LI. 1980. La madera en las artesanías del estado de Michoacana. *Boletín Divulgativo* 50. Distrito Federal, México: Instituto Nacional de Investigaciones Forestales. 128 p.
- Holdridge, LR, Poveda LR. 1975. *Arboles de Costa Rica*. Volume 1. San José, Costa Rica: Centro Científico Tropical. 546 p.
- Hughes CE, Styles BT. 1984. Exploration and seed collection of multiple-purpose dry zone trees in Central America. *International Tree Crops Journal* 3: 1-31.
- Janzen DH. 1983. *Costa Rican natural history*. Chicago: University of Chicago Press. 816 p.
- Little EL Jr, Woodbury RO, Wadsworth FH. 1974. *Trees of Puerto Rico and the Virgin Islands*, Volume 2. Agric. Handbk. 449. Washington, DC: USDA Forest Service. 1024 p.
- Neal MC. 1965. *In gardens of Hawaii*. Spec. Pub. 50. Honolulu: Bernice P. Bishop Museum. 924 p.
- Pennington TD, Sarukhan J. 1968. *Manual para la identificación de campo de los principales árboles tropicales de México*. Distrito Federal, México: Instituto Nacional de Investigaciones Forestales. 413 p.
- Salazar R. 1985. *Técnicas de producción de leña en fincas pequeñas*. Turrialba, Costa Rica: Centro Agronómico de Investigaciones y Enseñanza. 459 p.

Ephedraceae—Ephedra family

## *Ephedra* L.

### ephedra or Mormon-tea

Susan E. Meyer

Dr. Meyer is a research ecologist at the USDA Forest Service's Rocky Mountain Research Station, Shrub Sciences Laboratory, Provo Utah

**Growth habit, occurrence, and uses.** The genus *Ephedra*—known in much of North America as Mormon-tea—comprises about 40 shrubby species that are found throughout the arid and semiarid regions of the Northern Hemisphere. Ephedras are gymnosperms that are characterized by their greatly reduced, bractlike leaves and their evergreen, broomlike photosynthetic stems. They are common plants in the semiarid region of western North America (table 1) and are often locally codominant with creosotebush (*Larrea tridentata* (Sessé & Moc. ex DC.) Coville), blackbrush (*Coleogyne ramosissima* Torr.), shadscale saltbush (*Atriplex confertifolia* (Torr. & Frem.) S. Wats.), and various species of sagebrush (*Artemisia* spp.). Species of ephedra are often the dominant vegetation on sand hills at middle elevations, where they perform an important role as sandbinders. They provide a significant source of browse for domestic livestock, especially sheep, and for wild ungulates such as mule deer (*Odocoileus hemionus*) and pronghorn antelope (*Antilocapra americana*). The seeds provide food for rodents and birds. The twigs, especially those of green Mormon-tea, are used to make a reputedly refreshing tea, although ephedrine, the pharmaceutically active compound found in the Old World species *E. sinica* Stapf., has not been detected in any North American species. Ephedras are

attractive and interesting plants, with considerable potential for landscape use, and green Mormon-tea can now be readily obtained from commercial nurseries.

**Flowering and fruiting.** Ephedras are dioecious, with male and female cones occurring on separate plants. The cones are borne singly or in pairs or whorls at the branch nodes. The seeds are borne singly or in pairs in the axils of the female cone scales. The inner cone scales are modified to enclose the seed and form integuments that mimic the angiosperm pericarp. Flowering usually takes place in March through May, and seeds ripen from June through September, depending on elevation and species. The plants are wind-pollinated. Ephedra plants do not flower every year; their reproductive pattern could be described as mast fruiting, where most individuals in the population flower synchronously in a year with ample rainfall, and large quantities of seeds are produced. The population does not flower again for several years, whether or not a high-rainfall year occurs. The seedcrop may be damaged by late frosts, late spring drought, or infestations of pentatomid bugs.

The distribution of male and female ephedra plants is not random; individuals on dry slopes are overrepresented by males, whereas those growing on run-on surfaces are 4 times as likely to be females as males (Freeman and others

**Table 1**—*Ephedra*, ephedra: habit, habitat, and geographic distribution of some species used in revegetation

Scientific name	Common name(s)	Habit	Habitat	Distribution
<i>E. nevadensis</i> S. Wats.	Nevada Mormon-tea, gray Mormon-tea, gray ephedra, gray Nevada joint-fir	Sprawling, gray-green, leaves in pairs, bases	Creosote bush shrubland to pinyon-juniper woodlands	W US
<i>E. torreyana</i> S. Wats.	Torrey Mormon-tea, Torrey ephedra, Torrey's joint-fir	Sprawling, gray-green, leaves in whorls of 3, bases gray	Creosote bush shrub to pinyon-juniper woodlands	Colorado Plateau, Chihuahuan Desert
<i>E. viridis</i> Coville	green Mormon-tea, Brigham tea, green ephedra, Mormon-tea	Erect, broomlike, bright green, leaves in pairs, bases black	Blackbrush shrubland to mountain brush	W US


Sources: Welsh and others (1987).

1976). The genetic basis for sex differentiation in *Ephedra* is not known, but the spatial arrangement of males and females functions to maximize reproductive output, as it places males where their pollen can be easily wind-dispersed early in the season and females where they are more likely to have resources later in the season to ripen a seedcrop.

North American ephedra species fall into 2 groups characterized by differences in seed size and dispersal ecology (table 2). The large-seeded species (for example, green and Nevada Mormon-teas) are dispersed by scatter-hoarding rodents such as kangaroo rats (*Dipodomys* spp.), which deposit them in shallowly buried caches and later return to eat most of the seeds or sprouts. The cone scales in these species are small. In small-seeded species (for example, Torrey Mormon-tea) the outer cone scales are large and membranous, and the intact cones are often seen windrowed at some distance from adult plants. The seeds are apparently wind-dispersed, as they have long, awnlike points that probably make them unattractive to rodents. Cones with seeds intact may remain on the surface for many months.

**Fruit collection, cleaning, and storage.** Ephedra seeds (figures 1–3) are easily collected when fully ripe by beating the branches over a hopper or pan; in most years, large quantities can be collected in a short time. The collection window is narrow and crops must be watched carefully, as ripe seeds can be dislodged by wind or rain in a single day. Seed fill is usually high (table 2), but it is a good idea to check fill in the field before harvesting, as late drought can prevent filling of seeds that otherwise look normal. After the seeds are thoroughly dried, they may be broken free of the cone scales in a barley de-bearder if necessary and then cleaned in an air-screen cleaner (fanning mill). The seeds are usually long-lived in warehouse storage if initially of high quality, and storage times of 15 years may result in little viability loss (Stevens and others 1981). In a warehouse storage experiment, seedlots stored from 10 to 20 years had an average viability of 80% (n = 3), whereas seedlots stored from 20 to 30 years had an average viability of 31% (n = 12). The vigor of the older lots was low, however, as evi-

**Figure 1**—*Ephedra nevadensis*, Nevada Mormon-tea: seed.



denced by their low germination in response to chilling (<5% for lots >20 years old). It is doubtful whether seedlots >20 years old could be field-seeded successfully. The ability to remain highly viable for many years (orthodox storage behavior) facilitates stockpiling of ephedra seeds collected in most years for use over the period when few seeds are produced.

**Seed germination and testing.** Ephedra seeds are sometimes dormant at harvest, but this dormancy usually disappears through after-ripening after a few months in storage (Kay and others 1977). The dormancy also disappears after short periods (2 to 4 weeks) of chilling; this chilling also hastens the subsequent germination of nondormant seeds (Kay and others 1977; Meyer and others 1988). In experiments with 6-month-old seedlots, 7-day germination for unchilled seeds at 10 to 20 °C was 10% for 1 lot of green Mormon-tea, 49 to 54% for 2 lots of Nevada Mormon-tea, and 95 to 100% for 3 lots of Torrey Mormon-tea. The 7-day germination after 2 weeks of chilling at 1 °C was over 90% for all seedlots. Germination is generally highest at temperatures of 15 to 20 °C, except in more dormant lots, which show higher percentages of germination in temperature regimes that include a temperature in the chilling range (Young and others 1977). Germination is suppressed by higher temperatures, which probably prevent the


**Table 2**—*Ephedra*, ephedra: seed weight and viability data

Species	Seed weight				% viability	
	Range		Mean		Range	Mean
	/g	/oz	/g	/oz		
<i>E. nevadensis</i>	33–57	935–1,615	43	1,220	84–94	89
<i>E. torreyana</i>	108–128	3,060–3,630	118	3,345	76–91	83
<i>E. viridis</i>	28–62	790–1,760	47	1,330	46–100	89


**Sources:** Belcher (1985), Kay and others (1977), Meyer (1995), Meyer and others (1995), Monsen (1995).



**Figure 2**—*Ephedra viridis*, Torrey Mormon-tea: seeds (outer) and cone (center) with single seed.



**Figure 3**—*Ephedra nevadensis*, Nevada Mormon-tea: longitudinal section through a seed.



otherwise nondormant seeds from precocious summer germination. *Ephedra* seeds germinate readily during prolonged chilling. Kay and others (1977) reported 76% germination during a 30-day stratification at 2 °C for a Mojave desert collection of Nevada Mormon-tea. In chilling experiments with the 6 seedlots mentioned above, weeks to 50% germination at 1 °C varied from 6 to 7 weeks for the Torrey Mormon-tea collections and from 8 to 9 weeks for collections of the other 2 species. All viable seeds germinated during chilling within 12 weeks.

Official rules for testing green Mormon-tea call for a 4-week test at 15 °C, with the option of a 4-week prechill for

more dormant lots (AOSA 1993; Meyer and others 1988). Ungerminated seeds should be scored for viability using tetrazolium staining, which is also an acceptable substitute for a germination test. Seeds should be allowed to imbibe water, then clipped or slit on the cotyledon end and immersed in 1% tetrazolium solution for 8 hours. Seeds are then bisected longitudinally for evaluation.

**Nursery and field practice.** Large-seeded species of *ephedra* have been successfully established from direct seeding using drilling or with a seed dribbler or thimble seeder. If seeds are distributed aurally, a method for covering them with soil must be provided, as the seeds must be planted for successful establishment. Seedlings emerge and establish quickly and can withstand considerable drought once established. Emergence is best from depths of 1 to 2 cm (4/10 to 8/10 in) (Kay and others 1977). Emergence in green Mormon-tea is epigeal and the seedlings resemble those of conifers, whereas emergence in Nevada Mormon-tea is reported to be hypogeal (Kay and others 1977). Late-fall seedlings have been successful in the northern part of the range, where most effective precipitation comes in winter; whereas early-summer seedlings are recommended in the southern part of the range, where rainfall comes in the summer.

*Ephedra* planting stock may be produced either in bare-root or container culture. Plants do best in a coarse well-drained medium. Roots are fragile, so stock must be handled very carefully to avoid damage. The root systems of container stock are often too small to bind the root plug together, and those of bareroot stock are also usually poorly developed, resulting in low root–shoot ratios. Outplanting success rates are generally quite low (<50%).

#### References

- AOSA [Association of Official Seed Analysts]. 1993. Rules for testing seeds. *Journal of Seed Technology* 16(3): 11–13.
- Belcher E, ed. 1985. Handbook on seeds of browse shrubs and forbs. Tech. Pub. R8-TP8. Atlanta: USDA Forest Service, Southern Region. 246 p.
- Freeman DC, Klikoff LG, Harper KT. 1976. Differential resource utilization by the sexes of dioecious plants. *Science* 193: 597–599.
- Kay BL, Ross CM, Graves WL, Brown CR. 1977. Gray ephedra and green ephedra. *Mojave Reveg. Notes* 19. Davis: University of California, Department of Agronomy and Range Science. 8 p.
- Meyer SE. 1995. Unpublished data. Provo, UT: USDA Forest Service, Rocky Mountain Research Station.
- Meyer SE, Kitchen SG, Wilson GR, Stevens R. 1988. Proposed rule: *Ephedra viridis* green mormon tea. *Association of Official Seed Analysts Newsletter* 62(1): 18–19.
- Monsen SB. 1995. Unpublished data. Provo, UT: USDA Forest Service, Rocky Mountain Research Station.
- Stevens R, Jorgensen KR, Davis JN. 1981. Viability of seed from thirty-two shrub and forb species through fifteen years of warehouse storage. *Great Basin Naturalist* 41: 274–277.
- Welsh SL, Atwood ND, Higgins LC, Goodrich S. 1987. A Utah flora. *Great Basin Naturalist Memoirs*: 1–894.
- Young JA, Evans RA, Kay BL. 1977. *Ephedra* seed germination. *Agronomy Journal* 61: 209–211.

Ericaceae—Heath family

## *Epigaea repens* L.

### trailing-arbutus

Franklin T. Bonner

Dr. Bonner is a scientist emeritus at the USDA Forest Service's Southern Research Station, Mississippi State, Mississippi

**Other common names.** mayflower, ground-laurel, gravel weed, mountain-pink, winter-pink, crocus, gravel plant.



**Growth habit, distribution, and use.** Trailing-arbutus is an evergreen, prostrate, creeping shrub that grows in patches up to 60 cm in diameter (Bailey 1949). It is found growing in woodlands on acid, sandy soils from Florida to Mississippi, north to New England, southeast to New York, Pennsylvania, West Virginia, and Ohio. The variety *glabrifolia* Fern. ranges north from the higher parts of the Appalachian Mountains to Newfoundland, Nova Scotia, and Labrador and west to Saskatchewan (Fernald 1950). Although it is difficult to grow, trailing-arbutus has been planted as an ornamental since 1736 (Barrows 1936; Lemmon 1935). In some parts of its range it has become locally rare (Clay 1983). The blossoms are quite fragrant, and the fruits are sometimes eaten by small game. An infusion of the above-ground parts was used by the Cherokees to treat diarrhea in children (Jacobs and Burlage 1958).

**Flowering and fruiting.** The flowers are spicy smelling, pink to white in color, and bloom from March to May, although specimens have been known to bloom as early as January at low elevations in the southern part of its range (Stupka 1964). Flowering usually begins when plants are 3 years old (Steffek 1963). Flowering is normally dioecious, but perfect flowers may occasionally be found (Bailey 1949; Barrows 1936; Fernald 1950). Double-flowered forms and fall-blooming forms have been reported (Fernald 1950). The fruit is a 5-lobed, hairy, dehiscent capsule about 6 mm in diameter (Bailey 1949; Fernald 1950; Steffek 1963). The seeds are embedded in a sticky, white, fleshy pulp within the capsule (Barrows 1936; Clay 1983; Steffek 1963). A sample of 155 wild fruits contained an average of 241 (range: 29 to 415) tiny, shiny, brown, hard seeds per capsule (figures 1 and 2). In June and July, as the capsules ripen, the sutures split open and many of the seeds are ejected with some force (Blum and Krochmal 1974). As

the sutures begin splitting, ants will commonly enter the fruits and rapidly remove all seeds (Clay 1983).

**Collection of fruits; extraction and storage of seeds.** Capsules should be collected after they are mature and before they eject their seeds. Small collections of capsules can be air-dried in open containers until seeds are ejected. The empty capsules can be separated by screening. One sample of cleaned seeds contained 22,700 seeds/g (643,750/oz) (Blum and Krochmal 1974). Storing seeds for more than 1 year is not recommended, but short-term storage at room temperature or in a refrigerator is satisfactory (Barrows 1936).

**Figure 1**— *Epigaea repens*, trailing-arbutus: seeds (top) and longitudinal section through a seed (bottom).



**Germination tests.** Germination is epigeal and has been reported to require no pretreatment (Blum and Krochmal 1974). To secure complete germination on moist filter paper in petri dishes; however, Lincoln (1980) found it necessary to stratify seeds for 30 days at 5 to 8 °C and then germinate them at alternating temperatures of 15 to 25 °C or 20 to 30 °C with light at the higher temperature. This procedure yielded germination values of 92 and 90%, respectively.

**Nursery and field practice.** The seeds of trailing-arbutus are so small that sowing in small pots or trays filled with acid soil, sand, and peat moss or leaf mold mixtures is recommended (Blum and Krochmal 1974). The seeds should be scattered on top of the mixture, and the container should be covered with a glass plate or plastic bag to maintain a high humidity. With this method, germination takes place over a period of 22 to 66 days, with most germination occurring in 30 days (Barrows 1936). There are other reports of good germination within 3 to 5 weeks of time (Dirr and Heuser 1987).

When the seedlings have 3 to 5 leaves above the cotyledons, they may be transplanted to individual pots. High humidity should be maintained until the plants are well established (Barrows 1936). In 1 year, the plants develop into rosettes about 10 cm (4 in) in diameter (Blum and Krochmal 1974). Plants will tolerate a fairly wide range of acidity. Wild plants in Connecticut grew on soils ranging in pH from 7.67 to 4.65, but the larger plants occurred on the more acid soils (Barrows 1936; Coville 1911; Lemmon 1935; Steffek 1963).

Trailing-arbutus thrives best in association with mycorrhizal fungi. Including soil that was collected near healthy wild plants in soil mixtures will introduce the necessary fungus (Barrows 1936; Coville 1911, 1915). The mycorrhizal fungus also appears to be essential for propagation from cuttings (Barrows 1936). Stem cuttings taken in August have given 94% rooting in a sand-peat mixture without any treatment (Dirr and Heuser 1987).

#### References

- Bailey LH. 1949. Manual of cultivated plants. New York: Macmillan. 851 p.
- Barrows FL. 1936. Propagation of *Epigaea repens* L.: 1. Cuttings and seeds. Contributions of the Boyce Thompson Institute 8: 81-97.
- Blum BM, Krochmal A. 1974. *Epigaea repens* L., trailing-arbutus. In: Schopmeyer CS, tech. coord. Seeds of woody plants in the United States. Agric. Handbk. 450. Washington, DC: USDA Forest Service: 380-381.
- Clay K. 1983. Myrmecochory in the trailing arbutus (*Epigaea repens* L.). Torrey Botanical Club Bulletin 110: 166-169.
- Coville FV. 1911. The use of acid soil for raising seedlings of the mayflower, *Epigaea repens*. Science 33(853): 711-712.
- Coville FV. 1915. The cultivation of the mayflower. National Geographic Magazine 27: 518-519.
- Dirr MA, Heuser CW Jr. 1987. The reference manual of woody plant propagation. Athens, GA: Varsity Press. 239 p.
- Fernald ML. 1950. Gray's manual of botany. 8th ed. New York: American Book Co. 1632 p.
- Jacobs ML, Burlage HM. 1958. Index of plants of North Carolina with reputed medicinal uses. Chapel Hill, NC. 322 p.
- Lemmon RS. 1935. The trailing arbutus in home gardens. Horticulture 13: 101-102.
- Lincoln WC Jr. 1980. Laboratory germination of *Epigaea repens*, mayflower or trailing arbutus. AOSA Newsletter 54(1): 72-73.
- Steffek EF. 1963. Wild flowers and how to grow them. 5th ed. New York: Crown. 192 p.
- Stupka A. 1964. Trees, shrubs, and woody vines of the Great Smoky Mountains National Park. Knoxville: University of Tennessee Press. 186 p.

Asteraceae—Aster family

## *Ericameria parishii* (Greene) Hall

### Parish goldenweed

Raymond D. Ratliff and Franklin T. Bonner

Dr. Ratliff retired from the USDA Forest Service's Pacific Southwest Forest and Range Experiment Station; Dr. Bonner is a scientist emeritus at the Southern Research Station, Mississippi State, Mississippi

E

**Synonyms.** *Haplopappus parishii* (Greene) Blake, *Aplopappus parishii* (Greene) Blake, *Bigelovia parishii* Greene, *Chrysoma parishii* Greene.


**Other common names.** Parish goldenrod, Parish goldenbush, Parish heathgoldenrod.

**Growth habit, occurrence, and uses.** An erect shrub, Parish goldenweed has a mature height of 1 to 2.5 m (Jepson 1951). Plants 15 years old have attained heights of 2 m and crown spreads of 1.2 m (Everett 1957). Parish goldenweed occurs in the lower parts of the chaparral belt between 460 and 2,130 m of elevation in the mountains of southern California and Baja California (Munz and Keck 1959). Frequently, it is found on outwash fans and exposed hillsides. The primary value of this species is for erosion control on dry slopes (Ratliff 1974). Since the final writing of this manual, several sections of the genus *Chrysothamnus* (see table 1) have been transferred to the genus *Ericameria*.


**Flowering and fruiting.** Parish goldenweed will flower and bear seeds at 2 years of age and produce seeds each year thereafter (Everett 1957). Flowering takes place from July to October (Munz and Keck 1959), and ripe seeds may be collected in October and November (Mirov and Kraebel 1937). The fruit of Parish goldenweed is a single-seeded achene (figure 1) that is handled as a seed. The achenes are about 2 mm long (figure 2), and there are about 3,600 cleaned achenes/g (101,900/oz) (Mirov and Kraebel 1937).

**Collection, cleaning, and storage.** Achenes are usually collected by hand and separated from their bristles by rubbing and blowing (Ratliff 1974). There are no known studies of storage, but the seeds are probably orthodox and can be easily stored at low temperatures and moisture contents.

**Figure 1**—*Ericameria parishii*, Parish goldenweed: achene with pappus removed.



**Figure 2**—*Ericameria parishii*, Parish goldenweed: longitudinal section through an achene.



**Germination.** Parish goldenweed seeds are not dormant, and no pretreatments are required to stimulate germination (Emery 1964). Seeds sown on sand began germinating in 4 days, and a maximum of 95% was obtained (Mirov and Kraebel 1937). Germination is, however, usually much lower (about 20%) because of a high percentage of defective seeds (Ratliff 1974). Parish goldenweed may also be propagated by cuttings (Jepson 1951).

---

#### References

- Emery D. 1964. Seed propagation of native California plants. Santa Barbara Botanical Garden Leaflet 1(10): 81-96.
- Everett PC. 1957. A summary of the culture of California plants at the Rancho Santa Ana Botanic Garden 1927-1950. Claremont, CA: Rancho Santa Ana Botanic Garden. 223 p.
- Jepson WL. 1951. A manual of the flowering plants of California. Berkeley: University of California Press. 1025 p.
- Mirov NT, Kraebel CJ. 1937. Collecting and propagating the seeds of California wild plants. Res. Note 18. Berkeley, CA: USDA Forest Service, California Forest and Range Experiment Station. 27 p.
- Munz PA, Keck DD. 1959. A California flora. Berkeley: University of California Press. 1681 p.
- Ratliff RD. 1974. *Haplopappus parishii*, Parish goldenweed. In: Schopmeyer CS, tech. coord. Seeds of woody plants in the United States. Agric. Handbk. 450. Washington, DC: USDA Forest Service: 445.

Polygonaceae—Buckwheat family

## *Eriogonum* Michx. wild-buckwheat, buckwheatbrush

Susan E. Meyer

Dr. Meyer is a research ecologist at the USDA Forest Service's Rocky Mountain Research Station, Shrub Sciences Laboratory, Provo, Utah

**Growth habit, occurrence, and uses.** The North American genus *Eriogonum*—wild-buckwheat, also buckwheatbrush—is made up of about 200 species of annual and perennial herbs and shrubs, most of which are found in the West. About half are woody, at least at the base. The habit of the woody species may be either (a) truly shrubby, (b) subshrubby, with annual renewal of upper shoots, or (c) pulvinate (mat-forming), with the woody shoots condensed into an above-ground caudex. The usually evergreen leaves are borne alternately and may be predominantly basal or borne along the stems. There may be whorls of leaves on the flowering stalks. The leaves are usually tomentose, at least below, and the stem nodes are often tomentose as well. The often-flat-topped inflorescences are usually borne above the leafy part of the plant and are conspicuous and characteristic even after seed dispersal.

Most plant communities in the West contain at least 1 species of woody wild-buckwheat (table 1). Some species are widely distributed and of wide ecological amplitude (for example, sulfurflower buckwheat brush), whereas others are narrowly restricted geographically and often edaphically as well (for example, pretty buckwheat brush). Wild-buckwheat species are often important pioneer plants after natural disturbance, and their presence may facilitate the establishment of later-successional species. This makes them useful for erosion control and for revegetation of anthropogenically disturbed sites such as mined land and highway rights-of-way (Ratliff 1974; Zamora 1994). Some species are important as browse plants for wild ungulates, particularly in the early spring when their evergreen habit makes them more highly nutritive than many other spring browse species (Tiedemann and Driver 1983; Tiedemann and others 1997). Some wild-buckwheat species are important bee plants. In California, Mojave buckwheatbrush has been rated third in importance for honey production, exceeded only by 2 native *Salvia* species (Kay and others 1977). Many wild-buckwheat species also have tremendous potential as easily grown, drought-tolerant ornamentals. Their interesting forms and leaf textures combined with masses of showy,

long-lasting flowers make them excellent candidates for home xeriscapes. Named varieties that have been released are 'Sierra' sulfurflower wild-buckwheat (Stevens and others 1996) and 'Umatilla' snow wild-buckwheat (Tiedemann and others 1997).

**Flowering and fruiting.** The small, usually perfect flowers of wild-buckwheat are borne in clusters within cup-like or cylindrical involucre that are variously solitary or arrayed in capitate, cymose, or paniculate inflorescences. Each flower consists of a perianth with 9 stamens inserted at its base and a superior 1-celled and 1-seeded ovary. The perianth is made up of 6 fused segments in 2 whorls of 3. The ovary ripens in fruit into a usually 3-angled achene (figures 1 and 2). This achene is held more or less tightly within the perianth, depending on the species. For example, in snow wild-buckwheat the achenes fall free of the perianth at dispersal, whereas in Shockley wild-buckwheat the woolly perianth with the achene enclosed is the dispersal unit. The ovule within the seed is anatropous, so that the radicle end is pointing outward and upward. This makes it possible for germination and emergence to take place with the perianth still attached.

Wild-buckwheat species may flower at any time from early spring to fall, depending on species and habitat. Within a given habitat, species may bloom in succession. For example, at mid-elevation in central Utah, cushion wild-buckwheat blooms in spring, followed by James wild-buckwheat in early to midsummer, and finally by lace buckwheatbrush in late summer and fall. The bloom time for any species usually lasts well over a month, and the plants are almost as showy in fruit as in flower. The flowers are insect-pollinated.

**Seed collection, cleaning, and storage.** The window of opportunity for seed collection of wild-buckweats is often rather wide, as the fruits usually persist on the plant for 2 to 3 weeks after maturity (Stevens and others 1996). When achenes are mature, the perianths dry and often change color, turning brown or rusty. At this point, the achenes can be harvested by hand-stripping or by beating them into hop-



Table 1— *Eriogonum*, wild-buckwheat: habit, habitat, and geographic range

Species	Common name(s)*	Habitat	Range
<b>SHRUBS</b>			
<i>E. corymbosum</i> Benth.	lace buckwheatbrush, buckwheatbrush, crisp-leaf buckwheat	Desert shrub, pinyon juniper, mostly on shales	Colorado Plateau, Uinta Basin, & adjacent areas
<i>E. fasciculatum</i> Benth.	Mojave buckwheatbrush, California buckwheatbrush, flat-top buckwheatbrush	Warm desert shrub, coastal sage scrub, chaparral, pinyon-juniper	Mojave & Colorado Deserts & coastal & cismontane S California
<i>E. heermannii</i> Dur. & Hilg.	Heermann buckwheatbrush, molecule model plant	Warm desert shrub, mostly on rock outcrops	Mojave Desert
<b>SUBSHRUBS</b>			
<i>E. brevicaule</i> Nutt.	shortstem wild-buckwheat	Open, barren hills, mountain brush to alpine	Central Rocky Mtns of Wyoming, Utah & Idaho
<i>E. heracleoides</i> Nutt.	Wyeth wild-buckwheat, parsnipflower buckwheat	Sagebrush-grassland to aspen & Douglas-fir	N Rocky Mtns from BC to central Utah
<i>E. jamesii</i> Benth.	James wild-buckwheat	Desert shrub to mountain brush & ponderosa pine	S Rocky Mtns S into N Mexico
<i>E. niveum</i> Dougl. ex Benth.	snow wild-buckwheat, snow eriogonum	Sagebrush-grassland	Columbia River Plateau
<i>E. umbellatum</i> Torr.	sulfurflower wild-buckwheat, sulfur wildbuckwheat	Sagebrush-grassland to spruce-fir	Widespread in W North America
<b>PULVINATE/MAT-FORMING</b>			
<i>E. bicolor</i> M.E. Jones	pretty buckwheatbrush	Cold desert shrub, on Mancos Shale	Central Utah
<i>E. ovalifolium</i> Nutt.	cushion wild-buckwheat, roundleaf buckwheat	Wide range, from cold desert to alpine	Widespread, W North America
<i>E. shockleyi</i> S. Wats.	Shockley wild-buckwheat, mat buckwheat	Desert shrub to pinyon-juniper	Idaho & Colorado to SE California, Arizona, & New Mexico


Source: Meyer and Paulsen (2000).

Note: The genus *Eriogonum* is not that of the true, domesticated buckwheat, hence the common names of wild-buckwheat and buckwheatbrush.

**Figure 1**—*Eriogonum fasciculatum*, Mojave buckwheatbrush: achene in calyx (**left**) and achene without calyx (**right**).



**Figure 2**—*Eriogonum fasciculatum*, Mojave buckwheatbrush: longitudinal section through a seed excised from an achene.



pers or other containers. Combine harvesting has proven successful for sulfurflower wild-buckwheat in seed production fields (Stevens and others 1996). The harvested material will include achenes, perianths, involucre, and inflorescence branches. After the material is dried thoroughly, it may be threshed in a barley de-bearder and cleaned with a fanning mill. Species with tightly held achenes may require hand-rubbing through screens or on a rubbing board, which is also an alternative cleaning method for small seedlots of any species. The material should not be handled too rough-

ly, as the radicle end of the achene is often slender and easily damaged. Achene weights vary both among and within species but are usually in the range of 350 to 1,360/g (10,000 to 39,000/oz) (table 2). Seed quality is also variable (table 2).

There are few published reports of viability evaluation beyond germination percentages obtained without pretreatment, which may underestimate viability if there is a dormant fraction. Stevens and others (1996) report that viabilities of >90% may be expected from sulfurflower and Wyeth wild-buckwheats in an agronomic setting if seeds are harvested when fully mature; these values are comparable to those for wild-collected lots of many species (table 2). Insects may damage 10 to 35% of the fruits prior to harvest, but damaged seeds are normally eliminated in cleaning. Post-harvest damage from insect infestations is also possible (Stevens and others 1996). There is little information on maintenance of viability during storage for species of wild-buckwheat. Stevens and others (1996) report high viability for sulfurflower and Wyeth wild-buckwheats during 10 to 15 years in warehouse storage, which would indicate orthodox storage behavior. Other species are perhaps comparable.

**Seed germination and testing.** Germination is epigeal (figure 3). Seedlots of many species of wild-buckwheats contain at least a fraction that will germinate without any pretreatment (tables 2 and 3) (Young 1989). The size of this fraction depends on species and on the particular lot involved. Stevens and others (1996) report that seeds of sulfurflower and Wyeth wild-buckwheats lose dormancy during short periods of dry storage, and Mojave buckwheatbrush seeds are also reported to dry after-ripen (Kay and others 1977). Dormant seeds of most species we have examined lose dormancy during chilling at 1 °C for periods of 8 to 12 weeks (table 3).

To date there are no formal procedures for evaluating the seed quality of wild-buckwheat species, and tetrazolium (TZ) staining is probably the procedure most commonly employed. To evaluate using TZ, achenes are soaked overnight in water, clipped through both pericarp and seed coat at the cotyledon end (the wide end or hilum), and placed in 1% TZ solution for several hours at room temperature. Achenes are bisected longitudinally for evaluation (Belcher 1985).

**Field seeding and nursery practice.** Wild-buckwheats are generally readily established from direct seeding (Ratliff 1974; Stevens and others 1996; Tiedemann and Driver 1983; Zamora 1994). They establish best when seeded at a depth of 2 to 5 mm ( $1/16$  to  $3/16$  in), either by drilling or by broadcasting followed by covering (for example, raking). Seeding should take place before the season of maximum precipitation, which is generally fall or early winter in

**Table 2—*Eriogonum*, wild-buckwheat: achene weights and typical viability percentages**

Species	Achenes/weight		Viability	
	/g	/lb	%	Test
<b>SHRUBS</b>				
<i>E. corymbosum</i>	900	410,000	93	Post-chilling cut test
	2,000	900,000	—	—
<i>E. fasciculatum</i>	1,330	600,000	4–34	Germination %, no pretreatment
	520–1,085	236,000–490,000	20–46	Germination %, no pretreatment
<i>E. heermannii</i>	660	300,000	95	Post-chilling cut test
<b>SUBSHRUBS</b>				
<i>E. brevicaule</i>	700	320,000	84	Post-chilling cut test
<i>E. heracleoides</i>	350	160,000	95	Post-chilling cut test
	310	141,000	87	Post-chilling cut test
<i>E. jamesii</i>	350	160,000	—	—
<i>E. niveum</i>	1,290–1,360	585,000–620,000	52–72	Germination %; no pretreatment
<i>E. umbellatum</i>	470	213,000	86	Post-chilling cut test
	265	120,000	—	—
<b>PULVINATE/MAT-FORMING</b>				
<i>E. bicolor</i>	960	436,000	47	Post-chilling cut test
<i>E. ovalifolium</i>	990	450,000	95	Post-chilling cut test
<i>E. shockleyi</i>	750	340,000	86	Post-chilling cut test

**Sources:** Belcher (1985), Kay and others (1977), Meyer and Paulsen (2000), Stevens and others (1996), Tiedemann and Driver (1983).

\* Post-chilling cut tests (AOSA 1996) are considered accurate for recently harvested seedlots; however, tetrazolium staining (TZ) is required for seedlots stored for more than 2 years.

**Table 3—*Eriogonum*, wild-buckwheat: germination percentages**

Species	Samples	Germination* (% of total viable seeds)				
		No chill	4 weeks	8 weeks	12 weeks	16 weeks
<i>E. brevicaule</i>	2	3	28	65	86	96
<i>E. corymbosum</i>	3	28	79	100	100	100
<i>E. heracleoides</i>	3	4	11	30	55	77
<i>E. jamesii</i>	2	54	79	91	94	100
<i>E. ovalifolium</i>	2	22	74	98	98	100
<i>E. umbellatum</i>	4	7	30	74	99	100

**Source:** Meyer and Paulsen (2000).


\* Germination percentage determined after 0 to 16 weeks of chilling at 1 °C followed by 4 weeks of incubation at 10/20 °C

northern rainfall regions and midsummer in southern rainfall regions. Most wild-buckwheats are early seral and do not compete well with heavy stands of perennial grasses. Wild-buckwheats planted for field seed production are reported to reach 30 to 50% of maximum production, 200 to 400 kg/ha (180 to 360 lb/ac), the second year after planting (Stevens and others 1996).

Most species of wild-buckwheat are also easily propagated in a nursery setting. Shaw (1984) reported that Wyeth wild-buckwheat may be successfully produced as 1+0 bare-root stock. Because of the taprooted habit, plants must be

lifted carefully. Other woody wild-buckwheats could probably be produced as bareroot stock, but no published information is available. Wild-buckwheats may also be produced as container stock; book planters or tube containers such as those used for producing conifer seedlings are most appropriate. Nondormant lots may be direct-sown, whereas seedlots requiring chilling may be sown as chilled seed or as young germlings (Landis and Simonich 1984). Seedlings of many species grow rapidly and should not be held in small containers for more than a few months. Many species flower the first year and may even form flowering stalks while still in small tube containers.

**Figure 3**—*Eriogonum fasciculatum*, Mojave buckwheat-brush: very young seedling (left) and older seedling (right).



## References

- AOSA [Association of Official Seed Analysts]. 1996. Rules for testing seeds. *Journal of Seed Technology* 16(3): 1–113.
- Belcher E, ed. 1985. Handbook on seeds of browse-shrubs and forbs. Tech. Pub. R8-TP8. Atlanta: USDA Forest Service, Southern Region. 246 p.
- Kay BL, Ross CM, Graves WL. 1977. California buckwheat. Mojave Reveg. Notes 5. Davis: University of California, Department of Agronomy and Range Science. 4 p.
- Landis TD, Simonich EJ. 1984. Producing native plants as container seedlings. In: Murphy PM, comp. The challenge of producing native plants for the intermountain area. Proceedings, Intermountain Nurseryman's Association 1983 Conference; 1983 August 8–11; Las Vegas, NV. Gen. Tech. Rep. INT-168. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station: 16–25.
- Meyer SE, Paulsen A. 2000. Chilling requirements for seed germination of ten Utah species of wild buckwheat (*Eriogonum* Michx.: Polygonaceae). *Native Plants Journal* 1: 18–24.
- Ratliffe RD. 1974. *Eriogonum fasciculatum* Benth., California buckwheat. In: Schopmeyer CS, tech. coord. Seeds of woody plants in the United States. Agric. Handbk. 450. Washington, DC: USDA Forest Service: 382–383.
- Shaw NL. 1984. Producing bareroot seedlings of native shrubs. In: Murphy PM, comp. The challenge of producing native plants for the intermountain area. Proceedings, Intermountain Nurseryman's Association 1983 Conference; 1983 August 8–11; Las Vegas, NV. Gen. Tech. Rep. INT-168. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station: 6–15.
- Stevens R, Jorgensen KR, Young SA, Monsen SB. 1996. Forb and shrub seed production guide for Utah. Logan: Utah State University Extension Service. 51 p.
- Tiedemann AR, Driver CH. 1983. Snow eriogonum: a native halfshrub to revegetate winter game ranges. *Reclamation and Revegetation Research* 2: 31–39.
- Tiedemann AR, Lambert SM, Carlson JR, Perry CJ, Shaw NL, Welch BL, Driver CH. 1997. 'Umatilla' snow buckwheat for rangeland restoration in the interior Pacific Northwest. *Rangelands* 19(3): 22–25.
- Young JA. 1989. Germination of seeds of sulphur flower. *Journal of Seed Technology* 13: 31–38.
- Zamora BA. 1994. Use of *Eriogonum* for reclamation. *Hortus Northwest* 5(1): 9–11, 47.

Myrtaceae—Myrtle family

## *Eucalyptus* L'Her. eucalyptus

Stanley L. Krugman and Craig D. Whitesell

Dr. Krugman retired from the World Bank and from the USDA Forest Service's Washington Office;  
Dr. Whitesell retired from the USDA Forest Service's Pacific Southwest Research Station

**Growth habit, occurrence, and use.** The genus *Eucalyptus* comprises more than 523 species and 138 varieties, and new species and varieties are still being described (Blakely 1955; Johnston and Marryatt 1965; Penford and Willis 1961). Some are very tall trees, whereas others are woody shrubs (Jacobs 1979). Eucalypts are mainly native to Australia, but a few species are also native to the Philippines, New Guinea, and Timor (Hall and others 1963). Eucalypts are among the most widely cultivated forest trees in the world for ornamental use, shade, soil and site protection, wood production, and pulp making (Chippendale and others 1969). They are planted in southern Europe, the Middle East, Africa, India, Pakistan, China, North and South America—especially in Brazil, Uruguay, Chile, and Argentina (Jacobs 1979; Penford and Willis 1961).

This genus was first introduced into the United States with plantings in California and the Hawaiian Islands about 1853 (LeBarron 1962; Penford and Willis 1961). Eucalypts have also been planted, but to a limited extent, in Arizona, Florida, Georgia, Mississippi, and Texas (Geary and others 1983; Hunt and Zobel 1978; Metcalf 1961). About 250 species have been introduced into the United States and most of them are grown in California and Hawaii as ornamentals because of their decorative flowers and pleasing shapes (Jacobs 1979). Bluegum eucalyptus has been the most extensively planted eucalyptus in the United States, mainly in California, for the last 100 years (Metcalf 1961). It was initially grown for timber production but now is rarely used for this purpose. However, it is still widely used for fuel, shelterbelts, windbreaks, and has promise as a low-cost source of hardwood fiber (Krugman 1970). Among the other species that are promising for California conditions are river redgum, manna, mountain-gum, shining, and rosegum eucalyptuses (Ledig 1989). A number of other species are still being tested in California for their general landscape value (Hamilton 1979). In Florida, rosegum, robust, and river redgum eucalyptuses have been widely tested and are

most promising as a source of wood fiber (Geary and others 1983; Uhr 1976). In Hawaii, several species, including rosegum, robust, red-ironbark, saligna, and bluegum eucalyptuses, have been planted as windbreaks and for watershed protection, as well as for timber and biofuel production (LeBarron 1962; Whitesell and others 1992). There are numerous other species that hold promise for future fiber production, windbreaks, and for environmental forestry purposes (Fujii 1976; King and Krugman 1980) (table 1).

**Geographic races and hybrids.** Many eucalypts have an extensive natural distribution, and members of the same species often grow under very different environmental conditions (Boden 1964; Eldridge 1978; Hall and others 1963; Jacobs 1979). Although detailed scientific information as to the development of geographic races is lacking for most species, there is considerable genetic variation in those species with wide natural distribution and it can be assumed that numerous races do exist (Jacobs 1979; Miles 1990). For planting eucalypts in the United States, geographic origin must be considered in selecting a suitable seed source from species with extensive natural ranges, such as the widely grown river redgum eucalyptus (Eldridge 1975; Karschon 1967; Pryor and Byrne 1969). As a general rule, seed source selection should at least be based on a knowledge of the absolute minimal and maximal temperatures under which the species grows in its native range (Zon and Briscoe 1911). Differential low temperature tolerance has been demonstrated for different sources of broadleaf sallee (*E. camphora* R.T. Baker) and brown-barrel, lemon-gum, and manna eucalyptuses (Boden 1964; Hunt and Zobel 1978). Precipitation appears to be of less importance, but must also be considered in selecting the proper seed source.

Under natural conditions, hybridization between species of the same subgeneric group will take place. It is relatively common in some cases, for example, brown-barrel × narrow peppermint eucalyptus (*E. fastigata* × *radiata* Seibert ex DC.) and robusta × rosegum eucalyptuses (Boden 1964;

Table 1— <i>Eucalyptus</i> , eucalyptus: nomenclature and natural and extended ranges			
Scientific name & synonym(s)	Common name(s)	Natural range	Extension
<i>E. camaldulensis</i> Dehnhardt <i>E. rostrata</i> Schldl.	river redgum eucalyptus, red-gum, long-beak eucalyptus	Australia	California, Hawaii, & Arizona
<i>E. citriodora</i> Hook.	lemon-gum eucalyptus, lemon eucalyptus, lemon-gum	Central & N Queensland, Australia	California & Hawaii
<i>E. dalrympleana</i> Maiden	mountain-gum eucalyptus, white-gum, dalrymple eucalyptus	SE Australia	California & Hawaii
<i>E. delegatensis</i> R.T. Baker <i>E. gigantea</i> Hook. f.	alpine-ash eucalyptus, delegate eucalyptus	SE Australia	California
<i>E. fastigata</i> H. Deane & Maiden	brown-barrel eucalyptus, cuttail eucalyptus	SE Australia	California
<i>E. glaucescens</i> Maiden & Blakey	tingiringy-gum	SE Australia	California
<i>E. globulus</i> Labill.	bluegum eucalyptus, bluegum, Tasmania bluegum, Tasmanian blue eucalyptus	SE Australia	California, Hawaii, & Arizona
<i>E. grandis</i> W. Hill ex Maiden	rosegum eucalyptus, tooler eucalyptus.	E Australia	California, Florida, & Hawaii
<i>E. microcorys</i> F. Muell.	tallowwood eucalyptus	E Australia	California
<i>E. nitens</i> (H. Deane & Maiden) Maiden	shining eucalyptus, silver-top shining-gum	SE Australia	California & Hawaii
<i>E. obliqua</i> L'Her.	messmate stringybark eucalyptus	SE Australia	California
<i>E. paniculata</i> Sm.	gray ironbark eucalyptus, ironbark	E Australia	California & Hawaii
<i>E. pilularis</i> Sm.	blackbutt eucalyptus	E Australia	California & Hawaii
<i>E. regnans</i> F. Muell.	mountain-ash eucalyptus, swamp-gum giant eucalyptus	SE Australia	California
<i>E. robusta</i> Sm. <i>E. multiflora</i> Poir.	robusta eucalyptus, swamp-mahogany, beakpod eucalyptus	E Australia	California, Florida, Hawaii, & West Indies
<i>E. rudis</i> Sm.	desert eucalyptus, moitch eucalyptus, desert-gum	W Australia	California & Florida
<i>E. saligna</i> Sm.	saligna eucalyptus, Sidney bluegum eucalyptus, flooded-gum	E Australia	California & Hawaii
<i>E. sideoxylon</i> A. Cunningham	red ironbark eucalyptus, mulga ironbark eucalyptus, red-ironbark	SE Australia	California & Hawaii
<i>E. viminalis</i> Labill.	manna eucalyptus, ribbon eucalyptus, white-gum, ribbongum	SE Australia	California & Hawaii

Sources: Chippendale and others (1969), Johnston and Marryat (1965), Krugman (1974).

Jacobs 1979; Penford and Willis 1961; Pryor 1979). A number of hybrids have been described, but their value for planting in the United States must still be demonstrated. When grown under plantation conditions outside their natural habitat, species hybridization will occur more often, and seed collections from small plantations of closely related species should be discouraged if hybrid seeds are not desired (Boden 1964).

**Flowering and fruiting.** The flower clusters develop enclosed within an envelope formed by 2 bracteoles—small leafy structures. These bracteoles split and are shed during development, revealing the flower buds (Boland and others 1980; Penford and Willis 1961). The perfect flowers are white, yellow, or red, often in axillary umbels, corymbose,

or paniculate clusters (Blakely 1955). In a few cases, the flowers develop singly as with bluegum eucalyptus, but most often they are in 5- to 10-flowered axillary umbels as with river redgum and manna eucalyptuses (Blakely 1955). Sepals and petals are united to form a cap in the bud, which drops off at anthesis. The stigma is receptive within a few days after the cap drops (Barnard 1967) and pollination is mainly carried out by insects. The ovary has 3 to 6 locules with many ovules. There is a wide range in flowering times for the eucalypts (King and Krugman 1980). In California, some species such as manna eucalyptus may flower all year; other species, such as river redgum and gray ironbark eucalyptuses, flower in the spring; tingiringy-gum and mountain-gum eucalyptus in the summer; rosegum eucalyptus in the



fall; and tallowwood eucalyptus in winter (table 2) (King and Krugman 1980; Krugman 1970, 1974).

The fruit is a hemispherical, conical, oblong, or ovoid hard woody capsule 6 to 25 mm in diameter, that is loculicidally dehiscent at the apex by 3 to 6 valves (Blakely 1955; Boland and others 1980). The seeds are numerous and extremely small in most species (table 3; figure 1). The size of fertile seeds within a given seed collection varies widely. Usually only a few seeds are fertile in a single capsule, and capsule size may influence seed size (Blakely 1955). When

more than 1 seed ripens in a locule, the seeds are variously shaped and angular (figure 1). When solitary, the seed will be ovate or orbicular-compressed (Blakely 1955). The seedcoat is most often thin and smooth, but it can be ribbed, pitted, or sculptured in various ways (Blakely 1955; Penford and Willis 1961). Usually the seedcoat is black or dark brown in color as in manna eucalyptus or pale brown as with alpine-ash eucalyptus (table 3).

The embryo consists of bipartite or 2-lobed cotyledons that are folded or twisted over the straight radicle (Blakely

**Table 2—*Eucalyptus*, eucalyptus: height at maturity and phenology of flowering and fruiting of trees grown in California**

Species	Height at maturity (m)	Flowering	Fruit ripening	Seed dispersal
<i>E. camaldulensis</i>	18–36	Feb–Apr	July–Oct	Begins 8–9 months after flowering
<i>E. citriodora</i>	24–39	Nov–Jan	May–Aug	—
<i>E. dalrympleana</i>	18–36	June–Aug	Aug–Oct	Oct–Nov
<i>E. delegatensis</i>	30–83	Apr–June	Apr–July	May–July
<i>E. fastigata</i>	18–60	Apr–May	July–Aug	—
<i>E. glaucescens</i>	4–12	July–Aug	May–Sept	Nov–Feb
<i>E. globulus</i>	45–54	Nov–Apr	Oct–Mar	Oct–Mar
<i>E. grandis</i>	42–54	Sept–Nov	—	—
<i>E. microcorys</i>	30–45	Dec–Feb	—	—
<i>E. nitens</i>	30–90	Apr–July	May–June	May–June
<i>E. obliqua</i>	15–75	Apr–July	May–Aug	—
<i>E. paniculata</i>	24–42	Feb–May	—	—
<i>E. pilularis</i>	36–60	Dec–Mar	Jan–April	All year
<i>E. regnans</i>	52–105	Apr–July	June–Sept	—
<i>E. robusta</i>	24–27	Jan–Mar	—	—
<i>E. rudis</i>	9–15	Jan–Mar	—	—
<i>E. saligna</i>	15–45	Apr–June	Oct–Dec	—
<i>E. sideroxylon</i>	12–30	June–Sept	—	—
<i>E. viminalis</i>	15–45	All year	12 months after flowering	20–22 months after flowering

Source: Krugman (1970).


**Table 3—*Eucalyptus*, eucalyptus: description of viable seeds and chaff**

Species	Seed size (mm)		Seed color	Chaff color
	Length	Width		
<i>E. camaldulensis</i>	0.75–1.75	0.5–1.0	Yellow-brown	Yellow-brown to orange
<i>E. citriodora</i>	4.25	2.5	Black	Brownish red
<i>E. delegatensis</i>	1.25–3.75	1.0–1.75	Pale brown or brown	Pale brown or brown
<i>E. fastigata</i>	1.25–3.25	0.5–1.25	Pale brown or brown	Pale brown or brown
<i>E. glaucescens</i>	1.25–2.5	1.0–1.75	Black or dark brown	Pale red-brown
<i>E. globulus</i>	2.25	1.75	Dark brown	Brownish red
<i>E. nitens</i>	1.25–2.5	1.0–1.75	Black or dark brown	Pale red-brown
<i>E. obliqua</i>	1.0–2.0	0.75–1.25	Dark brown	Orange-brown or brown
<i>E. regnans</i>	1.25–2.5	0.5–1.25	Pale brown or brown	Pale brown or brown
<i>E. robusta</i>	1.5	0.75	Dark brown	Brownish red
<i>E. saligna</i>	1.25	1.0	Black	Brownish red
<i>E. sideroxylon</i>	1.0–2.0	0.75–1.5	Dark brown or black	Orange-brown
<i>E. viminalis</i>	1.25–2.5	1.5	Black or dark brown	Pale red-brown


1955; Krugman 1974). There is no endosperm (Blakely 1955; Krugman 1974) (figure 2). Fruits ripen at various times during the year, depending on the species (table 2). Dispersal is largely by wind within a month or two after ripening for most species, for example, bluegum and shining eucalyptuses. For other species, such as manna eucalyptus, dispersal may not take place until 10 months after ripening (table 2). Good seeds are produced by most species by 10 years of age (Grose 1969). For mature trees the interval between large seedcrops is from 2 to 5 years.

**Collection of fruits.** Collecting mature eucalyptus fruits should present no serious problems, other than reaching the fruit in very tall trees, because for most species there is a relatively long interval between seed ripening and opening of the capsule (table 2). However, it is important to take care to collect only well-developed, closed capsules, because capsules at different stages of maturity—as well as buds, flowers, and empty capsules—will be found on a single branch (Boland and others 1980; Krugman 1974). The capsules should be spread in a thin layer to permit rapid drying and to prevent mold formation (Boland and others 1980;

**Figure 1**—*Eucalyptus*, eucalyptus: seeds (from left to right) of *E. camaldulensis*, river redgum eucalyptus; *E. delegatensis*, alpine-ash eucalyptus; and *E. fastigata*, brown-barrel eucalyptus (top). *E. grandis*, rosegum eucalyptus; *E. microcorys*, tallowwood eucalyptus; *E. nitens*, shining eucalyptus (second row). *E. obliqua*, messmate stringybark eucalyptus; *E. paniculata*, gray ironbark eucalyptus; and *E. pilularis*, blackbutt eucalyptus (third row). *E. regnans*, mountain-ash eucalyptus, *E. robusta*, robusta eucalyptus; and *E. rudis*, desert eucalyptus (fourth row). *E. saligna*, saligna eucalyptus; *E. sideroxyton*, red ironbark eucalyptus; and *E. viminalis*, manna eucalyptus (bottom row).



**Figure 2**—*Eucalyptus rudis*, desert eucalyptus: longitudinal section through a seed.



Grose and Zimmer 1958b). The most common method is to air-dry the capsules for a few hours to a few days, depending on the maturity of the capsules (Boland and others 1980; Grose and Zimmer 1958b). Drying temperatures should not exceed 37.7 °C for prolonged periods, because high temperature may strengthen the dormancy of species such as alpine-ash, brown-barrel, shining, and mountain-ash eucalyptuses and tingiringy-gum (Grose 1969; Grose and Zimmer 1958b). Capsules can also be kiln-dried for relatively short periods (Boland and others 1980; Grose 1969). Fruit drying schedules for some of the common species are listed in table 4.

**Extraction and cleaning.** Once the capsules are open, they should be vigorously shaken to remove the seeds. Shaking is especially important if the capsules are somewhat immature, because viable seeds may not have separated completely from the capsule's placenta. Thus, unless the capsules are shaken, only infertile seeds will be extracted (Boland and others 1980; LeBarron 1962). When examined, immature capsules may appear empty after the aborted seeds are removed, because viable seeds are normally attached at the base of the capsule (LeBarron 1962). Viable seeds are extracted along with the unfertilized or aborted ovules, which are known collectively as "chaff" (Boland and others 1980; Grose and Zimmer 1958b). Large impurities such as the remains of twigs, capsules, and leaves can be removed by screening. Smaller impurities can be removed by specific-gravity separators such as the one used in the air-column method (Boland and others 1980; Grose 1969). For a few species, viable seeds can be separated from the remaining chaff by employing sieves of the appropriate mesh size (Boland and others 1980). Because viable seeds and chaff of

**Table 4**—*Eucalyptus*, eucalyptus: fruit drying schedules

Species	Air-drying		Kiln-drying	
	Temp (°C)	Time (days)	Temp (°C)	Time (hrs)
<i>E. camaldulensis</i>	32	1	59	3
<i>E. delegatensis</i>	32	3	59	6
<i>E. globulus</i>	21	5	—	—
<i>E. obliqua</i>	32	3	59	5
<i>E. regnans</i>	32	3	59	6
<i>E. sideroxylon</i>	21	4	—	—
<i>E. viminalis</i>	21	6	—	—

most species cannot be separated by the usual methods, commercial seed suppliers sell chaff along with the fertile seeds. The proportion by weight of chaff to viable seeds is in the range of 5:1 to 30:1 (Grose and Zimmer 1958b). For some species, such as mountain-ash eucalyptus, the seeds and chaff are identical in size and color; for others, such as river red-gum eucalyptus, the seeds and chaff are similar in color but different in size (table 3). For most species, there are some differences in color and size, so that viable seeds can be separated from chaff to some extent if necessary. Because of their very small size, seeds of eucalyptus species are normally sold by weight with the chaff. The average number of viable seeds plus chaff per weight ranges from 770/g (21,900/oz) for river redgum eucalyptus to 35/g for blackbutt eucalyptus (1,000/oz) (table 5). There may be as many as 2,100 seeds/g (59,500/oz) for river redgum eucalyptus or as few as 7 seeds/g (200/oz) for blackbutt eucalyptus.

**Storage.** Eucalyptus seeds are orthodox in storage behavior. They should be stored in air-tight containers that are as completely filled as possible to reduce the amount of air (Boland and others 1980). Prior to storage, the seeds should be treated to kill insect pests, by either fumigation or placing paradichlorobenzene crystals in the container (Boland and others 1980).

Eucalypts seeds have germinated after 30 years of storage at room temperature, but the germination was very low (Penford and Willis 1961). Most seeds can be successfully stored for periods up to 10 years in air-tight containers at moisture contents of 4 to 6% and temperatures of 0 to 5 °C (Boland and others 1980; Grose 1969; Grose and Zimmer 1958b). It should be possible to store these seeds successfully for even longer at temperatures below 0 °C (Krugman 1970).

**Pregermination treatments.** Most eucalyptus seeds need no pretreatment to ensure adequate germination if fresh seeds are used (Boland and others 1980; Grose and Zimmer

1958b). A few species—such as tingiringy-gum and alpine-ash, brown-barrel, shining, and mountain-ash eucalyptuses—are normally dormant at the time of collection and will require pretreatment. Stratification of moist seeds stored in a plastic bag at temperatures of 3 to 5 °C for a period of 3 weeks will break the dormancy of these 5 species, except for alpine-ash eucalyptus, which should be stratified for 4 weeks (Boland and others 1980; Grose 1969). Longer stratification periods of 6 to 8 weeks are often recommended.

Dormancy between different seedlots of the same species can vary considerably. In addition, different methods of extraction and storage can induce dormancy in nondormant seed or strengthen primary dormancy in normally dormant seeds (Krugman 1970). If the seeds fail to germinate after the recommended shorter stratification periods, then a longer period should be tried before the seeds are considered nonviable. Because most seeds are stored before they are used, stratification for 3 to 4 weeks at a temperature of 3 to 5 °C is recommended for all eucalyptus seeds to ensure faster and more uniform germination (Hinkle 1968; Krugman 1970).

In a few cases, chemicals have been employed to overcome seed dormancy. The germination of unstratified and dormant seeds of alpine-ash, brown-barrel, and mountain-ash eucalyptuses was improved by germinating the seeds in a solution of gibberellic acid (Bachelord 1967). However, not all seedlots of the same species responded to gibberellic acid (Krugman 1970).

**Germination tests.** Standard methods for testing germination in other seeds are not used for eucalyptus seeds because of their small size and the presence of so much chaff, which can exceed the weight of viable seeds. Instead, samples for germination are of equal weight, not number (Boland and others 1980; Grose and Zimmer 1958b; ISTA 1993; Turnbull and Doran 1987). Such methods as the excised-embryo and tetrazolium tests are impractical (Boland and others 1980; Grose and Zimmer 1958b). The

**Table 5**—*Eucalyptus*, eucalyptus: seed yield data

Species	Viable seeds/(weight of seeds + chaff)				Samples #
	Range		Average		
	lg	loz	lg	loz	
<i>E. camaldulensis</i>	65–2,100	1,800–59,500	770	21,900	41
<i>E. citriodora</i>	80–220	2,200–6,200	140	4,000	15
<i>E. dalrympleana</i>	65–285	1,800–8,100	195	5,500	7
<i>E. delegatensis</i>	40–125	1,100–3,500	75	2,100	13
<i>E. fastigata</i>	90–210	2,500–5,900	150	4,300	6
<i>E. glaucescens</i>	40–120	1,000–3,000	35	2,000	2
<i>E. globulus</i>	20–70	500–9,100	150	2,500	10
<i>E. grandis</i>	200–1,200	5,600–34,000	700	20,000	13
<i>E. microcorys</i>	530–900	1,500–25,600	85	6,800	22
<i>E. nitens</i>	230–550	6,600–15,700	385	10,900	7
<i>E. obliqua</i>	20–160	500–4,500	85	2,400	18
<i>E. paniculata</i>	65–340	1,800–9,600	75	5,000	8
<i>E. pilularis</i>	7–85	200–2,400	35	1,000	28
<i>E. regnans</i>	20–530	600–15,000	315	8,900	11
<i>E. robusta</i>	220–700	6,200–20,000	390	11,000	12
<i>E. rudis</i>	270–1,100	7,600–31,000	600	17,000	9
<i>E. saligna</i>	85–915	2,400–26,000	460	13,000	9
<i>E. sideroxylon</i>	65–440	1,800–12,500	240	6,800	16
<i>E. viminalis</i>	265–445	7,500–12,600	350	10,000	6

Sources: Grose and Zimmer (1958b), Larsen (1965).

International Seed Testing Association (1993) recommends a sample unit of 0.10 to 1.0 g of seeds, depending on the species. Seeds are placed on 1 or more layers of moist paper and germinated at a constant temperature of 15 to 35 °C, depending on the species (Grose 1969; Scott 1972). Some species may require alternating temperatures of 20 °C for 16 hours and 30 °C for 8 hours (ISTA 1993). The tests are normally conducted under lights, although lights are not necessary for all species. Recommendations for individual species are listed in table 6. Official rules (ISTA 1993) provide recommendations for many more species. Immature seeds of mountain-ash eucalyptus should be tested under lights (Penford and Willis 1961).

If an approximate estimate of viability is desired, a known weight of dry seeds can be soaked in water and then squashed systematically. All seeds that show a firm white embryo can be recorded as viable (Grose and Zimmer 1958b).

Soundness of eucalyptus seeds is highly variable. Seeds collected from individual trees of bluegum eucalyptus in California showed from 2 to 80% germination after 30 days (Krugman 1970). Germination of from 11 to 98% has been reported for other species (table 7).

**Nursery practice.** On the United States mainland, eucalyptus seeds are rarely sown directly in the nursery, a practice once common in Hawaii. The most common practice for growing eucalyptus seedlings is to germinate the

**Table 6**—*Eucalyptus*, eucalyptus: germination test conditions

Species	Daily light exposure (hrs)	Temp (°C)	Days
<i>E. camaldulensis</i>	24	35	14
<i>E. citriodora</i>	0	25	14
<i>E. dalrympleana</i>	24	25	14
<i>E. delegatensis</i> †	0	20	14
<i>E. fastigata</i> ‡	0	25	14 or 21
<i>E. glaucescens</i> ‡	24	20	14 or 21
<i>E. globulus</i>	24	25	14
<i>E. grandis</i>	0	25	14
<i>E. microcorys</i>	24	20§	28
<i>E. nitens</i> ‡	24	20	14 or 21
<i>E. obliqua</i>	0	20	14
<i>E. paniculata</i>	24	20§	28
<i>E. pilularis</i>	0	20	14
<i>E. regnans</i> ‡	24	25	21
<i>E. robusta</i>	24	20	28
<i>E. rudis</i>	24	35	14
<i>E. saligna</i>	24	25	28
<i>E. sideroxylon</i>	24	20	14
<i>E. viminalis</i>	24	25	14

Sources: Grose (1969), ISTA (1966, 1993).

\* Seeds germinated on 2 layers of moist, filter paper in a petri dish (Grose 1969).

† Prechilled for 28 days at 3.3 to 5 °C (Grose 1969).

‡ Prechilled for 21 days at 3.3 to 5 °C (Grose 1969).

§ Treated at 20 °C for 16 hours, then 30 °C for 8 hours.

**Table 7**—*Eucalyptus*, eucalyptus: germination test results

Species	% Germination	Duration (days)
<i>E. citriodora</i>	51	15
<i>E. grandis</i>	98	29
<i>E. microcorys</i>	76	24
<i>E. pilularis</i>	11	29
<i>E. robusta</i>	84	18
	100	21
<i>E. sideroxylon</i>	69	49

Sources: Floyd (1964), Ganguli (1966), Krugman (1974), Scott (1972).

seeds in small pots, boxes, or wooden flats. Commonly used containers are wooden flats or plastic containers 45 to 50 cm (18 to 20 in) long and 40 to 50 cm (16 to 20 in) wide, and 10 to 12.5 cm (4 to 5 in) deep with good bottom drainage (Hinkle 1968; Jacobs 1955). The planting medium should be porous, friable, and light textured, such as a light, sandy loam (Hinkle 1968; Holmes and Floyd 1969; Willan 1985). The medium must permit good drainage and should not cake or become hard on the surface after watering. Because of possible weed and disease problems, the soil should be sterilized. Various mixtures are also used, with the most common consisting of equal parts (by volume) of sand, soil, and organic matter. The flats are filled to a depth of 7.5 to 10 cm (3 to 4 in) and the soil surface is leveled.

Because of their small size, the seeds are mixed with a little sand, and the mixture then is spread evenly over the soil surface (Hinkle 1968; Jacobs 1955; Penford and Willis 1961). The seeds are covered with 3 mm of fine sand, peat, or sphagnum moss to prevent surface drying.


A newer method to ensure more precision in sowing eucalyptus seeds is to coat them and form a mini-pellet (Geary and Millier 1982; Willan 1985). The seed dressing is usually an inert material serving as a sticker on which other materials can be added. One successful method involves a coating of fine silica sand filler and a polyvinyl alcohol binder (Geary and Millier 1982). In addition, fungicides and insecticides can also be added.

Enough seeds should be sown to raise between 500 and 2,000 plants/flat (table 8). Depending on seed size, this should represent about 7 g ( $1/4$  oz) of seed/flat (Krugman 1970). The flats should be well watered and drained just before planting and should be protected from wind, heavy rains, and excessive heat. The emerging seedlings may require protection from birds and rodents in some locations.

**Table 8**—*Eucalyptus*, eucalyptus: seedlings produced per weight of seeds

Species	Seedlings produced/seed weight	
	/g	/oz
<i>E. camaldulensis</i>	175	5,000
<i>E. citriodora</i>	60	1,700
<i>E. glaucescens</i>	99	2,830
<i>E. globulus</i>	28	800
<i>E. microcorys</i>	7	180
<i>E. paniculata</i>	6	160
<i>E. robusta</i>	14	400
<i>E. viminalis</i>	111	3,160

Sources: Zon and Briscoe (1911).

**Figure 3**—*Eucalyptus*, eucalyptus: seedling development for manna eucalyptus at 1 day (left) and 8 days (center) and desert eucalyptus at 42 days (right).

**Seedling care.** Germination is epigeal (figure 3), begins in 7 to 10 days, and is completed in 3 to 4 weeks (Boland and others 1980; Jacobs 1955; Krugman 1974). Because the seeds are small and the seedlings very delicate, overhead watering should be with a fine spray and care must be taken to maintain adequate soil moisture.

When the seedlings are about 6 to 8 weeks old and have developed 2 pairs of leaves and a third pair is just visible above the cotyledons, they can be transplanted into suitable containers for further growth (Hinkle 1968; Jacobs 1955).

In transplanting, the seedlings should be lifted by the tip of a sturdy leaf, and not by the soft delicate stem. A dibble should be used to protect the fibrous root system. Prior to lifting, the seedlings should be hardened off by exposure to the open air away from full sunlight and strong winds for a few days to a week. A variety of different containers, from Jiffy™ pots to tin cans, have been used with success (Holmes and Floyd 1969; Jacobs 1955). The containers should be large enough to permit the development of strong plants, but small enough to permit ease of transportation. Tubes should be at least 4 cm (1.5 in) in diameter and 15 to 30 cm (6 to 12 in) long (Jacobs 1955). When metal tubes are employed, they are made so that they can be readily opened in the field and then later cleaned and reused. After transplants are placed in containers, care must be taken to prevent damage to them. Seedlings should be well watered and shaded from full sunlight. Fine gravel can be placed on the surface of the container to restrict slime molds. After several weeks the transplants can be placed in the open so that they can become hardy. They should be ready for outplanting in 4 to 5 months, depending on the species and growing conditions (Jacobs 1955).

Because of the rapid growth of eucalyptus seedlings, care must be taken lest they become pot-bound. Seedlings should not be permitted to grow in small containers for extended periods before outplanting.

Seeds can be sown directly in a standard 1.2-m-wide (4-ft-wide) nurserybed. The soil should be first fumigated to kill weed seeds and pathogens, then watered well and drained before sowing. Because seeds are small, even distribution is difficult when they are broadcast sown. Seeds should be sown in narrow strips or rows, covered with a thin layer of sand or peat, and watered thoroughly (Penford and Willis 1961). Under very hot conditions the nursery beds should be shaded. Young eucalyptus seedlings need a great amount of light, so only moderate shade is recommended. If bareroot stock is desired, the seedlings are left in the beds for about 6 to 12 months. More commonly, the seedlings are lifted after 5 to 10 weeks and planted in individual containers. Because of differences in seed size and purity among species seedlots, the variation in number of seedlings produced, by a given weight of seeds, will vary widely (Zon and Briscoe 1911).

**Vegetative propagation.** Vegetative propagation by rooting and grafting has been successful in some of the eucalypts. The following species have been rooted successfully: lemon-gum, red-ironbark, river redgum, mountain-ash, robust, desert, mountain-gum, blue-gum, rosegum, and manna eucalyptuses (Blomstedt and others 1991; Jacobs 1979; Linnard 1969; Penford and Willis 1961). But in the main, only shoots with juvenile leaves have been rooted, and these most often from trees younger than 5 years (Jacobs 1979; Penford and Willis 1961). Eucalyptus can also be propagated by grafting. At the present time, vegetative propagation appears to be a practical method for producing eucalypts in large numbers only in countries where labor is inexpensive, for example, Brazil (Jacobs 1979). The production of possibly useful cultivars is both difficult and expensive, for cuttings from mature eucalyptus trees do not readily root (Chippendale and others 1969).

## References

- Bachelord EP. 1967. Effects of gibberellic acid, kinetin, and light on the germination of dormant seeds of some eucalypt species. *Australian Journal of Botany* 15: 393–401.
- Barnard RC. 1967. Some garden eucalypts. *Australian Plants* 4(30): 69–70, 71–72.
- Blakely WF. 1955. *A key to the eucalypts*. 2nd ed. Canberra: Forest and Timber Bureau. 359 p.
- Blomstedt C, Cameron J, Whiteman P, Chandler SF. 1991. Micropropagation of juvenile *Eucalyptus regnans* (mountain ash). *Australian Journal of Botany* 39(2): 179–186.
- Boden RW. 1964. Hybridization in eucalyptus. *Indian Forester* 90(9): 581–586.
- Boland DJ, Brooker MIH, Turnbull JW. 1980. *Eucalyptus seed*. Sydney, Australia: Hogbin Poole Pty. Ltd. 191 p.
- Chippendale GM, Johnston, RD, Kelly S. 1969. *Eucalypts*. Melbourne, Australia: Thomas Nelson (Australia), Ltd. 82 p.
- Eldridge KG. 1975. An annotated bibliography of genetic variation in *Eucalyptus camaldulensis*. Trop. For. Pap. 8. Oxford, UK: Oxford University, Department of Forestry. 59 p.
- Eldridge KG. 1978. Genetic improvement of eucalypts. *Silvae Genetica* 27(5): 205–209.
- Floyd AG. 1964. Germination test methods for tree seeds. Tech. Pap. 7. Sydney, Australia: Commonwealth Forestry. 15 p.
- Fujii DM. 1976. The Nuuanu eucalyptus planting: growth, survival, stand development after 64 years. Res. Note PSW-318. Berkeley, CA: USDA Forest Service, Pacific Southwest Forest and Range Experiment Station. 5 p.
- Ganguli BN. 1966. Application of Czabator's approach to germination studies of *Eucalyptus*. *Science and Culture* 32(9): 466–468.
- Geary TF, Millier WF. 1982. Pelleting small volumes of eucalyptus seeds for precision sowing. *South African Forestry Journal* 121: 79–83.

- Geary TF, Meskimen GF, Franklin EC. 1983. Growing eucalypts in Florida for industrial wood production. Gen. Tech. Rep. SE-23. USDA Forest Service, Southeastern Forest Experiment Station. 43 p.
- Grose RJ. 1969. Personal communication. Melbourne, Australia: Commonwealth Forestry.
- Grose RJ, Zimmer WJ. 1958a. A description of the seeds of 70 Victorian eucalypts. Bull. 8. Melbourne, Australia: Commonwealth Forestry. 24 p.
- Grose RJ, Zimmer WJ. 1958b. The collection and testing of seed from some Victorian eucalypts with results of viability tests. Bull. 10. Melbourne, Australia: Commonwealth Forestry. 14 p.
- Hall N, Johnston RD, Marryatt R. 1963. The natural occurrence of the eucalypts. Leaflet. 65. 2nd ed. Canberra, Australia: Forestry and Timber Bureau. 122 p.
- Hamilton WD. 1979. The landscape eucalyptus tree: an evaluation study 1971–1978. *Journal of Arboriculture* 5(9): 210–216.
- Hinkle EH. 1968. Eucalyptus in Taiwan. *Quarterly Journal of Chinese Forestry* 1(2): 87–98.
- Holmes DA, Floyd AG. 1969. Nursery techniques for raising eucalypts in jiffy pots on the New South Wales north coast. Res. Note 22. Sydney, Australia: Commonwealth Forestry. 15 p.
- Hunt R, Zobel B. 1978. Frost-hardy eucalyptus grow well in the Southeast. *Southern Journal of Applied Forestry* 2(1): 6–10.
- ISTA [International Seed Testing Association]. 1966. International rules for seed testing. *Proceedings of the International Seed Testing Association* 31(1): 52–106.
- ISTA [International Seed Testing Association]. 1993. International rules for seed testing: rules 1993. *Seed Science and Technology* 21 (Suppl.): 1–259.
- Jacobs MR. 1955. Growth habits of the eucalypts. Canberra, Australia: Forestry and Timber Bureau. 262 p.
- Jacobs MR. 1979. Eucalypts for planting. For. Ser. 11. Rome: FAO. 677 p.
- Johnston RD, Marryatt R. 1965. Taxonomy and nomenclature of eucalypts. Leaflet. 92. Canberra, Australia: Forestry and Timber Bureau. 24 p.
- Karschon R. 1967. Ecotypic variation in *Eucalyptus camaldulensis* Dehn. In: Contributions on eucalypts in Israel. [National University Institute of Agricultural, Ilanot, and Land Development Authority]: Kiriath Hayim III: 35–53.
- King JP, Krugman SL. 1980. Tests of 36 *Eucalyptus* species in northern California. Res. Pap. PSW-152. Berkeley, CA: USDA Forest Service, Pacific Southwest Forest and Range Experiment Station. 6 p.
- Krugman SL. 1970. Observations recorded 1965–1970. Berkeley, CA: USDA Forest Service, Pacific Southwest Forest and Range Experiment Station. 11 p.
- Krugman SL. 1974. *Eucalyptus, eucalyptus*. In: Schopmeyer CS, tech. coord. Seeds of woody plants in the United States. Agric. Handbk. 450. Washington, DC: USDA Forest Service: 384–392.
- Larsen E. 1965. Germination of *Eucalyptus* seed. Leaflet. 94. Canberra, Australia: Forestry and Timber Bureau. 16 p.
- LeBarron RK. 1962. Eucalypts in Hawaii: a survey of practices and research programs. Misc. Pap. 64. Berkeley, CA: USDA Forest Service, Pacific Southwest Forest and Range Experiment Station. 24 p.
- Ledig FT. 1989. Improvement of eucalypts for fuel and fiber in California. In: Pereira JS, Landsberg JJ, eds. Biomass production by fast growing trees. Proceedings, NATO Advanced Research Workshop, Series E. Dordrecht, The Netherlands: Kluwer Academic Publishers: 231–245.
- Linnard W. 1969. Cultivation of eucalypts in the USSR. *Forestry Abstracts* 30(2): 199–209.
- Metcalfe W. 1961. Progress with eucalypts in North America [unpublished office report]. Section A, United States mainland. National Report, 2nd World Eucalyptus Conference, Brazil: 1–18.
- Miles M. 1990. Principles of species and clonal selection for farm forestry. *California Eucalyptus Grower* 5(1): 8, 9, 11.
- Penford AR, Willis JL. 1961. The eucalypts: botany, cultivation, chemistry, and utilization. New York: Interscience Publishers. 551 p.
- Pryor ID. 1979. Reproductive habits of the eucalypt. *Unasylva* 30(119/120): 42–46.
- Pryor LD, Byrne OR. 1969. Variation and taxonomy in *Eucalyptus camaldulensis*. *Silvae Genetica* 18(3): 64–71.
- Scott L. 1972. Viability testing of eucalypt seeds. Leaflet. 116. Canberra, Australia: Forestry and Timber Bureau. 24 p.
- Toyama S, Moromizato S. 1957. Breeding of eucalyptus trees (report 1): characteristics of eucalyptus seed [in Japanese]. Government Forest Experiment Station (Tokyo) Bulletin 103: 1–24.
- Turnbull JW, Doran JC. 1987. Germination of Australian native plant seed. In: Langkamp PJ, ed. Melbourne, Australia: CSIRO, Division of Forest Research: 46–57, 186–198.
- Uhr SC. 1976. Eucalypt: the wonder tree. *American Forests* 82(10): 42–43, 59–63.
- Whitesell CD, DeBell DS, Schubert TS, Strand RF, Crabb TB. 1992. Short-rotation management of eucalyptus: guidelines for plantations in Hawaii. Gen. Tech. Rep. PSW-137. Albany, CA: USDA Forest Service Pacific Southwest Research Station. 30 p.
- Willan, RL. 1985. A guide to forest seed handling. For. Pap. 20/2. Rome: FAO. 379 p.
- Zon R, Briscoe JM. 1911. Eucalypts in Florida. Bull. 87. Washington, DC: USDA Forest Service. 47 p.



Celastraceae—Bittersweet family

## *Euonymus* L.

### euonymus, spindletree

John C. Zasada and Paul O. Rudolf

Dr. Zasada retired from the USDA Forest Service's North Central Forest Research Station;  
Dr. Rudolf (deceased) retired from the USDA Forest Service's North Central Forest Experiment Station

**Growth habit, occurrence, and use.** The genus *Euonymus* includes about 170 species of deciduous or evergreen shrubs and small trees, sometimes creeping or climbing, native to North and Central America, Europe, Asia, Madagascar, and Australia (Krüssmann 1960; Rehder 1940). The majority of species are native to east Asia from 52°N latitude to the Tropics (Nikolaeva 1967). Because of their attractive fruits and foliage, the euonymus species are planted widely for ornamental purposes. Winged spindletree, described by Dirr (1990) as “one of the finest landscape plants for American gardens,” has brilliant red foliage in the fall (it is commonly known as burning-bush) and prominent corky wings on the stem that add variety to the winter landscape. *Euonymus* species show a large amount of variability: for example, Dirr (1990) listed about 70 cultivars recognized

by horticulturists. At least one of these introduced species—European spindletree—has become naturalized and is considered invasive in the Northeast; other species do not appear to be as aggressive (Dirr 1990; Fernald 1950; Gleason and Cronquist 1963; Voss 1985). The deciduous and evergreen euonymus used as ornamentals in Britain have been described by Lancaster (1981). They also have value for wildlife food, shelterbelts, and minor wood products; at least 1 species is a source of gutta (Nikolaeva 1967). Eight species that have been used for conservation plantings are described in tables 1 and 2.

The 3 native species (table 1) described by Fernald (1950) occur in sites generally described as “rich” and with a mesic to wet soil water regime. In Wisconsin, eastern wahoo is most common in southern wet forests that are

**Table 1**—*Euonymus*, euonymus: nomenclature and occurrence

Scientific name & synonym(s)	Common name(s)	Occurrence
<i>E. alata</i> (Thunb.) Sieb. <i>Celastris alatus</i> Thunb.	winged spindletree, winged euonymus, corkbush, burning-bush	Central China, Manchuria, E Siberia, Korea, Japan, & Sakhalin
<i>E. americana</i> L.	American strawberry-bush, bursting-heart, hearts-a-bustin, brook euonymus	New York to Illinois to Texas to Florida
<i>E. atropurpurea</i> Jacq.	eastern wahoo, burning-bush, wahoo	W New York to S Ontario, central Michigan & Minnesota, SE North Dakota, S to NW Nebraska, central Kansas, & E Texas, E to Arkansas, Tennessee, & N Alabama
<i>E. bungeana</i> Maxim.	winterberry euonymus	N China, Manchuria, & Korea
<i>E. europaea</i> L.	European spindletree, European euonymus	Europe to W Asia (to 900 m in mtns)
<i>E. hamiltoniana</i> spp. <i>maackii</i> (Rupr.) Komarov	Maack euonymus	N China & Korea
<i>E. obovata</i> Nutt.	running strawberry-bush, running euonymus	W New York & S Ontario to central Michigan, Illinois, S to West Virginia, Kentucky, & Missouri
<i>E. verrucosa</i> Scop.	warty-bark euonymus, warty spindletree	S Europe & W Asia

Sources: Dirr (1990), Rudolf (1974).

**Table 2**—*Euonymus*, euonymus: height and year first cultivated

Species	Height at maturity (m)	Year first cultivated
<i>E. alata</i>	0.9–3.1	1860
<i>E. americana</i>	0.9–1.8	1697
<i>E. atropurpurea</i>	1.8–6.2	1756
<i>E. bungeana</i>	4.0–6.2	1883
<i>E. europaea</i>	3.1–7.1	Long ago
<i>E. hamiltoniana</i> ssp. <i>maackii</i>	1.5–5.2	1880
<i>E. obovata</i>	0.3–0.6	1820
<i>E. verrucosa</i>	0.9–2.2	1763


Sources: Dirr (1990), Fernald (1950), Lancaster (1981), Rehder (1940).

dominated by silver maple (*Acer saccharinum* L.), black willow (*Salix nigra* Marsh.), and other trees characteristically found on wet sites (Curtis 1959); in Michigan, it is described as a floodplain species (Voss 1985). European spindletree, the naturalized exotic, is generally found on moist to wet sites, and floodplains in central Europe where it occurs naturally (Lee and others 1991). Historically, the roots, bark, and seeds were used for medicinal purposes with the warning that products of each may be poisonous to some individuals (Foster and Duke 1990; Snow and Snow 1988).

**Flowering and fruiting.** The usually perfect flowers, borne in clusters, bloom in the spring. The fruit, which ripens in late summer or fall, is a 4- to 5-celled (occasionally 2- to 3-celled) capsule that is usually lobed and sometimes winged (figure 1). Ostrobuka and Bencat (1987) found that winged spindletree pollen germinated in sucrose concentrations of 15, 20, and 25%, with 20% giving best results. Each fruit cell contains 1 or 2 seeds enclosed in a fleshy, usually orange aril (figure 2). Natural seed dispersal usually occurs soon after the fruits are fully ripe. Seed dispersal of European spindletree is primarily by birds, with robins (*Erithacus rubecula*) being a principal disperser in Britain. Some species ingest the entire aril, whereas others carry it to a perch and remove the pulp and drop the seeds (Snow and Snow 1988). Fruits of European spindletree have some of the highest lipid and protein contents reported for plants (Snow and Snow 1988). The flowering and fruiting habits of 8 species are summarized in tables 3 and 4.

Fruits are generally available annually. Flower bud differentiation occurs from early June to mid-August and weather conditions during this period will affect fruit production potential (Tomita and Uematsu 1978).

The fruit of European spindletree contains 1 to 5 seeds. Dry weight of the fruit is 0.17 g, with the seed accounting for about 45% of the dry weight. At maturity the water con-

**Figure 1**—*Euonymus*, euonymus: top views of open capsules of *E. americana*, American strawberry-bush (left) and *E. atropurpurea*, eastern wahoo (right).**Figure 2**—*Euonymus americana*, American strawberry-bush: seeds enclosed in their fleshy arils.

tent is about 50% (fresh weight basis) (Lee and others 1991; Nielsen and Iroquoian 1988). Lee and others (1991) described the seeds of European spindletree as poisonous and little used by birds; however, Snow and Snow (1988) document a substantial use of fruits by birds, stating that the seeds are not consumed.

**Collection of fruits.** Seeds may be collected in late summer or fall by picking the ripe fruits from the bushes or trees by hand or by shaking them onto an outspread canvas. They should then be spread out to dry for several days in a warm room but need not be completely dry to be cleaned (Myatt and others 1991).

**Extraction and storage of seeds.** Seeds can be processed with a macerator (Stein and others 1974). The plate on the separator should be set slightly larger than the seeds and adjusted as necessary to prevent too many seeds from being lost with the pulp (Myatt and others 1991).

Smaller seedlots can be extracted by beating the fruits in a canvas bag and then rubbing them through a coarse, round-hole grain screen. The fruits may be macerated in water and the seeds extracted by flotation (Rudolf 1974). Following dry extraction, the chaff can be removed by

**Table 3**—*Euonymus, euonymus*: phenology of flowering and fruiting

Species	Location	Flowering	Fruit ripening
<i>E. alata</i>	New England & Japan	May–June	Sept–Oct
<i>E. americana</i>	Carolinas	May–June	Sept–Oct
<i>E. atropurpurea</i>	—	May–June	Sept–Oct
<i>E. bungeana</i>	NE US	June	Sept–Oct
<i>E. europaea</i>	NE US & Europe	May–June	Aug–Nov
<i>E. hamiltoniana</i> ssp. <i>maackii</i>	NE US	June	Oct
<i>E. obovata</i>	—	April–June	Aug–Oct
<i>E. verrucosa</i>	NE US & Germany	May–June	Aug–Oct

**Sources:** Fernald (1950), Lancaster (1981), Radford and others (1964), Rehder (1940), Sus (1925), Snow and Snow (1988), Voss (1985), Wappes (1932), Wyman (1947).

**Table 4**—*Euonymus, euonymus*: fruit form and color of flowers, fruits, and seeds

Species	Fruit form	Color			
		Flower	Ripe fruit	Seed	Aril
<i>E. alata</i>	Divided nearly to base in 4 separate pods (sometimes 1–3)	Yellowish	Reddish-purplish	Brown*	Orange-red
<i>E. americana</i>	3- to 5-lobed	Reddish green–greenish purple	Pink-rose	Yellowish white	Scarlet
<i>E. atropurea</i>	Smooth, deeply 3- to 4-lobed, 4-celled	Purple	Pink–purple	Light brown	Scarlet
<i>E. bungeana</i>	Deeply 4-lobed & 4-angled	Yellowish	Yellowish–pinkish white	Whitish or pinkish	Orange
<i>E. europaea</i>	Smooth, 4-lobed, 3 to 5-celled	Yellowish green	Rose red–pink†	White	Orange
<i>E. hamiltoniana</i> ssp. <i>maackii</i>	4-lobed	Yellowish	Pink	Red	Orange
<i>E. obovata</i>	Usually 3-lobed	Greenish purple	Crimson	Tan	Orange-scarlet
<i>E. verrucosa</i>	Deeply 4-lobed	Brownish	Yellowish red	Black	Orange-red‡

**Sources:** Bailey (1939), Dirr (1990), Fernald (1950), Rehder (1940), Snow and Snow (1988).  
\* Black, in one variety.  
† Whitish, in one variety.  
‡ Seed not wholly covered by aril.


winning. The pulpy arils can be removed (figure 3) by rubbing the seeds through coarse-mesh wire cloth after they have dried several weeks, but this is difficult to do without breaking the relatively thin seedcoats (Lee and others 1991) and injuring the seeds. If the arils (which are sometimes oily) are not removed, the seeds may not store as well. As a result, commercial seeds are treated rather gently in extraction and seedlots usually contain seeds with parts of the arils still attached, along with completely clean seeds (figure 4).

The numbers of seeds per weight cleaned to this variable degree are shown in table 5. Forty-five kilograms (100 lb) of ripe fruits will yield about 4.5 to 9.1 kg (10 to 20 lb) and an average of 7.1 kg (16 lb) of cleaned seeds, based upon data for American strawberry-bush, eastern wahoo, European spindletree, and warty-bark euonymus (Gorshenin 1941; Rudolf 1974; Swingle 1939).

Seed weights can vary significantly within a population. Nielsen and Iroquoian (1988) reported that the variation in the dry weight of 1,000 seeds ranged from about 28 to 40 g among 8 individual European spindletree plants. Mature seeds from different positions in the plant varied significantly in seed weight; seeds from the top and shaded parts of the crown were 5% greater and 7% less than mean seed weight, respectively (Nielsen and Iroquoian 1988).

Seeds of European spindletree and warty-bark euonymus can be kept satisfactorily for 2 years in ordinary dry storage (Gorshenin 1941; Sus 1925), or in dry cold storage in sealed containers at 1 to 2 °C (Heit 1967). However, more recent Russian reports have shown high viability maintained for at least 7 years under moist conditions at constant temperatures, either warm (15 to 20 °C) or cold (Nikolaeva 1967). Any drying in storage reduced viability (Nikolaeva

**Figure 3**—*Euonymus*, euonymus: seeds with arils removed of *E. americana*, american strawberry-bush (**top**); *E. atropurpurea*, eastern wahoo (**middle**); and *E. obovata*, running euonymus (**bottom**).



1967). Moist cold storage may be the most practical and effective way of retaining high viability of euonymus seeds for extended periods (table 6).

**Pregermination treatments.** Seeds of most euonymus species have dormant embryos. Cold stratification is adequate to break dormancy for some species, but warm stratification followed by a cold period is needed for maximum germination for other species (table 7) (Dirr 1990; Dirr and Heuser 1987; Nikolaeva 1967; Singh 1985; Yu and others 1976). The length of the warm period should be adjusted, depending on the temperature used for cold stratification. For example, Nikolaeva (1967) suggests a 2- to 3-month period of warm stratification if cold stratification is at 0 to 3 °C. Table 7 provides the range of temperatures for warm and cold stratification that have been effective for breaking dormancy. Nikolaeva (1967) provides a thorough discussion of the effects of temperature, water availability, seed maturation, and storage alone and in combination on germination. There may also be some variation in germination among European spindle tree seeds formed under different climatic conditions (Dawidowicz-Grezgorzewska and Beranger-Novat 1989).

Variation in dormancy can be significantly different among plants. Nielsen (1988), for example, reported that germination of European spindle tree seeds collected from 10 different plants varied from 0 to 30% following stratification

**Table 5**—*Euonymus*, euonymus: seed yield data

Species	Place collected	Cleaned seeds/weight				Samples
		Range		Average		
		/kg	/lb	/kg	/lb	
<i>E. alata</i>	NE US	41,110–69,620	18,600–31,500	55,250	25,000	2+
<i>E. americana</i>	Durham Co., North Carolina	63,300–100,113	30,000–45,300	77,571	35,100	3+
<i>E. atropurpurea</i>	Carver Co., Minnesota; Cole Co., Missouri; & rangewide	19,227–88,400	8,700–40,000	37,349	16,900	8
<i>E. bungeana</i>	US	—	—	29,835	13,500	1+
<i>E. europaea</i>	Russia, Netherlands, & NE US	18,785–35,360	8,500–16,000	29,393	13,300	32+
<i>E. obovata</i>	Clinton Co., Michigan	—	—	49,725	25,500	1
<i>E. verrucosa</i>	Russia	35,950–58,870	16,300–26,700	45,084	20,400	10+

Sources: Barnes (1969), Gorshenin (1941), Heit (1968a), NBV (1946), Nielsen and Eriksen (1988), Rudolf (1974), Sus (1925), Swingle (1939).

**Table 6**—*Euonymus*, euonymus: seed storage conditions

Species	Seed storage conditions		Viable period (yrs)
	Seed moisture	Temp (°C)	
<i>E. atropurpurea</i> *	Air-dry	–1.1–3.3	—
<i>E. europaea</i>	Dry	—	1–2
	Moist	15–20 or 2.8	7
<i>E. obovatas</i> *	Air-dry	1.1–3.3	—
<i>E. verrucosa</i>	Air-dry	20	2
	Moist	15–20 or 2.8	7


Sources: Gorshenin (1941), Heit (1967), NBV (1946), Nikolaeva (1967), Rudolf (1974), Sus (1925), Swingle (1939).

\* Seed stored in closed containers.

**Table 7**—*Euonymus*, euonymus: stratification treatments

Species	Moisture-holding medium	Warm period		Cold period	
		Temp (°C)	Days	Temp (°C)	Days
<i>E. alata</i>	Sand or peat	—	0	0–10	90–100
<i>E. americana</i>	Perlite–peat mix	—	0	5	139
<i>E. atropurpurea</i>	Sand	20–30	60	5	60
	Sand	—	0	2.8–5	60–180
<i>E. bungeana</i>	Sand, peat, or filter paper	—	0	2.8–10	61–120
<i>E. europaea</i>	Sand, peat, or filter paper	20–25	60–90	32–50	60–120
	Sand or filter paper	—	—	2.8–5	60–120
<i>E. hamiltoniana</i> ssp. <i>maackii</i>	Sand or filter paper	—	0	0–10	60–90
<i>E. obovata</i>	Sand	—	0	2.8–5	60–150
<i>E. verrucosa</i>	Sand or filter paper	15–20	60–90	0–10	120–150

Sources: Heit (1968a), Nikolaeva (1967), Rudolf (1974), Shumailina (1949), Swingle (1939).

**Figure 4**—*Euonymus europaea*, European spindletree: longitudinal section through a seed.

at 4 to 6 °C. This variation may have been significantly different if seeds had been subjected to warm stratification before cold stratification (table 7) (Nikolaeva 1967).

The morphological changes that occur during pretreatment are important indicators of the adequacy of the pregermination treatment. An increase in seed volume, cracking of the seedcoat, and protrusion of the tip of the hypocotyl occur during warm stratification or warm stratification hastens these changes when seeds are moved to cold stratification. Completeness of germination depends on these changes in seed morphology (Nikolaeva 1967).

There seems to be little information on the natural germination pattern of euonymus seeds. Untreated seeds of

European spindletree germinated mainly in the second year after sowing (Lee and others 1991), suggesting that the alternation of warm and cold stratification also regulates germination under field conditions. Although birds are important in the dispersal of European spindletree seeds, the seeds may (which can cause some degree of scarification) or may not pass through the digestive tract of birds, depending on the species taking the seeds (Snow and Snow 1988).

**Germination tests.** Germination is epigeal (figure 5). Germination tests on stratified seeds can be run in sand flats, germinators, or petri dishes. ISTA (1993) recommends 45 days of stratification at 3 to 5°C, then a 28-day test at 20/30 °C for European spindletree. Viability can also be estimated by the embryo excision method (Heit 1966) or tetrazolium staining (ISTA 1993). Germination test conditions are summarized in table 8.


X-radiography has been used to assess viability in fresh European spindletree seeds. However, stored seeds should be stained with tetrazolium to determine viability (Smirnova and Tikhomirova (1980).

**Nursery practice.** Seedlings can be grown in containers or nurserybeds. For best results, cleaned euonymus seeds should be sown in the fall soon after collection, before the seeds have dried out (Heit 1968a&b; NBV 1946). If sowing after seeds have been collected is not feasible, stratified seeds can be planted (table 6) early the next spring or the next fall (NBV 1946). Details for most species are lacking, but for European spindletree, recommendations are to sow the seeds 6 mm ( $\frac{1}{4}$  in) deep at a density to produce 422 seedlings/m<sup>2</sup> (40/ft<sup>2</sup>) of nurserybed (NBV 1946). The beds should be mulched with pine straw (NBV 1946). Tree percentages range from about 10% for winterberry euonymus to 20% for eastern wahoo and 25% for European spindletree (Swingle 1939).

**Table 8—*Euonymus*, euonymus:** germination test conditions and results on stratified seeds

Species	Germination test conditions				Days	Avg germ capacity (%)	Purity (%)	Soundness (%)	Samples
	Medium	Temp (°C)							
		Day	Night						
<i>E. americana</i>	Filter paper	21.1	21.1	14	15	—	—	1	
<i>E. atropurpurea</i>	Sand flats	30	20	61	40	75	88	2	
<i>E. bungeana</i>	Germinator	10	0	60	20	—	—	1	
<i>E. europaea</i>	Sand flats, germinators	25	20	60	71	75	96	22+	
<i>E. hamiltoniana</i> ssp. <i>maackii</i>	Germinators	20	15	60	75	—	—	3+	
<i>E. verrucosa</i>	Sand flats, filter paper	20	12	60	70	75	96	7+	

Sources: NBV (1946), Nikolaeva (1967), Rudolf (1974), Swingle (1939).

**Figure 5—*Euonymus europaea*, European spindle tree:** seedlings 1 and 12 days after germination.

### References

- Bailey LH. 1939. The standard cyclopedia of horticulture. New York: Macmillan 3639 p.
- Curtis JT. 1959. The vegetation of Wisconsin: an ordination of plant communities. Madison: University of Wisconsin Press. 657 p.
- Dawidowicz-Grzegorzewska A, Beranger-Novat N. 1989. Ultrastructural studies on development in mature seeds: embryos of *Euonymus europaea* L. dormant or cultured at +25 °C. Journal of Experimental Botany 40: 913–918.
- Dirr MA. 1990. Manual of woody landscape plants: their identification, ornamental characteristics, culture, propagation, and uses. Champaign, IL: Stipes Publishing Co. 1007 p.
- Dirr MA, Heuser CW Jr. 1987. The reference manual of woody plant propagation: from seeds to tissue culture. Athens, GA: Varsity Press. 239 p.
- Fernald ML. 1950. Gray's manual of botany. New York: American Book Co. 1632 p.
- Foster S, Duke JA. 1990. A field guide to medicinal plants: eastern and central North America. New York: Houghton Mifflin. 366 p.
- Gleason HA, Cronquist A. 1963. Manual of vascular plants of northeastern United States and adjacent Canada. Boston: Willard Grant Press. 810 p.
- Gorshenin NM. 1941. Agrolesomelioratsiya [Agro-forest melioration]. Moscow. 392 p.
- Heit CE. 1966. Propagation from seed: 2. Testing extremely dormant seeds. American Nurseryman 124(10): 10, 11, 38, 40.
- Heit CE. 1967. Propagation from seed: 11. Storage of deciduous tree and shrub seeds. American Nurseryman 126(10): 12–13, 86–94.
- Heit CE. 1968a. Propagation from seed: 15. Fall planting of shrub seeds for successful seedling production. American Nurseryman 128(4): 8–10, 70–80.
- Heit CE. 1968b. Thirty-five years' testing of tree and shrub seeds. Journal of Forestry 66(8): 632–634.
- ISTA [International Seed Testing Association]. 1993. International rules for seed testing: rules 1993. Seed Science and Technology (suppl.): 1–259.
- Krüssman G. 1960. Handbuch der Laubgehölze. Volume 1 and Volume 2. 495 & 608 p.
- Lancaster R. 1981. An account of *Euonymus* in cultivation and its availability in cultivation. Plantsman 3: 133–166.
- Lee WG, Grubb PJ, Wilson JB. 1991. Patterns of resource allocation in fleshy fruits of nine European tall shrubs. Oikos 61: 307–315.
- Myatt A, Huffman G, Odell J. 1991. Seed processing manual. Stillwater: Oklahoma Department of Agriculture, Forestry Division, Forest Restoration Center: 1 p.
- NBV [Nederlandsche Boschbouw Vereeniging]. 1946. Boomzaden: handleiding inzake het oegsten, behandelen, bewaren en uitzaaien van boomzaden. Wageningen, the Netherlands: Ponsen and Looijen. 171 p.
- Nielsen KK. 1988. Dormancy in seeds from different positions on individual plants. Acta Horticulturae 226(I): 255–261.
- Nielsen KK, Ericksen EN. 1988. Dry weight and moisture content in seeds from different positions in individual plants. Acta Horticulturae 226(II): 525–528.

- Nikolaeva MG. 1967. Fiziologiya glubokogo pokoya semyan. Akad. Nauk SSSR, Bot. Inst. V. L. Komarova. Izdatel'stvo "Nauka," Leningrad. [Physiology of deep dormancy in seeds. Transl. TT 68-50463, 1969. Springfield, VA: USDC CFSTI. 220 p.]
- Ostrolucka MG, Bencat F. 1987. Pollen viability of some introduced woody species in the in-vitro conditions. Lesnictvi 33: 789–810.
- Radford AE, Ahles HE, Bell CR. 1964. Guide to the vascular flora of the Carolinas. Chapel Hill: University of North Carolina Book Exchange. 383 p.
- Rehder A. 1940. Manual of cultivated trees and shrubs hardy in North America. New York: Macmillan. 996 p.
- Rudolf PO. 1974. *Euonymus*, euonymus. In: Schopmeyer CS, tech. coord. Seeds of woody plants in the United States. Agric. Handbk. 450. Washington, DC: USDA Forest Service: 393–397.
- Shumilina ZK. 1949. Podgotovka posevu semyan drevesnykh I kustarnikovykh porod. Vsesoyuznyi Nauchno-issledovatel'skii Inst. Agrolesomelior., Goslesbumizdat, Moskva-Leningrad [Preparation of tree and shrub seed for sowing. Transl. TT 67-51300, 36 p., 1967. CFSTI, U.S. Dept. Commerce, Springfield, VA].
- Singh CP. 1985. A comparison between low temperature and gibberellic acid removed dormancy of *Euonymus europaeus* L. embryos with respect to seedling growth and development [abstract]. Phytos 45: 143–148.
- Smirnova NG, Tikhomirova NI. 1980. Combined use of X-ray and the tetrazolium method for assessing seed viability [in Russian, English abstract]. Byulleten Glavnogo Botanicheskogo Sada 117: 81–85.
- Snow B, Snow D. 1988. Birds and berries: a study of an ecological interaction. Calton, Scotland: T & AD Poyser. 268 p.
- Stein WI, Slabaugh PE, Plummer AP. 1974. Harvesting, processing, and storage of fruits and seeds. In: Schopmeyer CS, tech. coord. Seeds of woody plants in the United States. Agric. Handbk. 450. Washington, DC: USDA Forest Service: 98–125.
- Sus NI. 1925. Pitomnik [in Russian: The forest nursery.] Moscow. 227 p.
- Swingle CF, comp. 1939. Seed propagation of trees, shrubs, and forbs for conservation planting. SCS-TP-27. Washington, DC: USDA Soil Conservation Service. 198 p.
- Tomita H, Uematsu J. 1978. Studies on the flowering in some potted ornamental trees and shrubs: 2. On the time of flower bud differentiation and flower bud development in 6 species [in Japanese, English abstract]. Bulletin of the Saitama Horticultural Experiment Station 7: 55–63.
- Voss EG. 1985. Michigan flora: 2. Dicots. Cranbrook Institute of Science Bulletin 59: 1–724.
- Wappes L. 1932. Wald und Holz ein Nachschlagebuch für die Praxis der Forstwirte, Holzhändler und Holzindustriellen. Volume 1. Berlin: J. Neumann. 872 p.
- Wyman D. 1947. Seed collecting dates of woody plants. Arnoldia 7(9): 53–56.
- Yu TY, Yeam DY, Kim YJ. [1976]. Studies on seed propagation in ornamental plants: I. The chilling requirement of *Euonymus alatus*, *Euonymus japonicus*, *Pyracantha coccinea*, and *Thuja occidentalis* seeds for germination. Journal of the Korean Society of Horticultural Science 17: 164–168.