



# The Value of Forests to Pollinating Insects Varies with Forest Structure, Composition, and Age

Michael D. Ulyshen<sup>1</sup> · Kimberly M. Ballare<sup>2</sup> · Christopher J. Fettig<sup>3</sup> · James W. Rivers<sup>4</sup> · Justin B. Runyon<sup>2</sup>

Accepted: 5 June 2024

This is a U.S. Government work and not under copyright protection in the US; foreign copyright protection may apply 2024

## Abstract

**Purpose of Review** Forests play an essential role in conserving pollinating insects and supporting pollination services in mixed-use landscapes and are particularly important to species that require resources restricted to forests. However, some forests provide higher quality habitat for these organisms than others. The primary objectives of this article are to 1) review how pollinator communities are influenced by changes in forest structure, composition, and age, 2) explore how these patterns differ between conifer and broadleaf forests, and 3) provide recommendations for managers interested in optimizing forest conditions for pollinating insects.

**Recent Findings** Although biodiversity generally increases as forests mature and become more structurally and compositionally complex, patterns exhibited by pollinating insects vary depending on forest type and prevailing disturbance regimes. For example, conifer forests can either sustain pollinator diversity comparable to open habitats or experience reduced pollinator diversity depending on the openness of the canopy. In broadleaf forests, pollinator diversity often increases with age (following the stem exclusion stage) and increasing tree diversity, and diversity in these areas can exceed what is observed in open habitats even under closed-canopy conditions. Such patterns likely reflect the importance of flowering broadleaf trees to pollinators, including many forest-dependent species, and suggest that optimal management practices for conserving pollinators differ between conifer and broadleaf forests.

**Summary** We conclude that: 1) the quality of forests to pollinating insects is a function of forest structure and composition as mediated by forest age and disturbance history and 2) best management practices need to be developed separately for conserving pollinators in conifer and broadleaf forests. Research aimed at better understanding the value of different broadleaf tree taxa to pollinators, especially forest-dependent species, is needed.

**Keywords** Bees · Early seral forests · Forest canopy · Forest succession · Mature forests · Nesting resources

## Introduction

Recognition that forests are important to pollinating insects (hereafter ‘pollinators’) has been building for decades [1–5], and interest in this topic has surged over the past several

years [6••, 7, 8•, 9••, 10]. Far from providing suboptimal conditions for these organisms as suggested by some authors, it is now clear that forests support a large fraction of pollinator diversity endemic to many regions [11••]. Forests not only play a key role in maintaining regional pollinator diversity but they also promote pollination in adjacent land uses including many crops [11••], sometimes with substantial economic benefits [12, 13•].

To fully understand the value of forests to pollinators, it is first necessary to distinguish forest-dependent species from those that use forests opportunistically [14]. Forest-dependent species require resources restricted to forests at some point during their life cycle. Although such resources often include the flowers of trees or other forest plants, they can also include other plant structures, such as leaf or stem material used for food (e.g., by lepidopteran larvae), and substrates or microhabitats used for nesting or breeding (e.g.,

✉ Michael D. Ulyshen  
michael.d.ulyshen@usda.gov

<sup>1</sup> USDA Forest Service, Southern Research Station, Athens, GA, USA

<sup>2</sup> USDA Forest Service, Rocky Mountain Research Station, Bozeman, MT, USA

<sup>3</sup> USDA Forest Service, Pacific Southwest Research Station, Davis, CA, USA

<sup>4</sup> Department of Forest Engineering, Resources, and Management, Oregon State University, Corvallis, OR, USA

dead wood, hollow trunks, or tree cavities). By contrast, pollinators that are habitat generalists can persist in landscapes devoid of forests yet may still make use of forest resources to meet their life history needs.

Recent work suggests that approximately a third of bee species are associated with forests in the northeastern U.S. [9••]. Work from other regions indicates that extensive deforestation results in the replacement of forest-dependent pollinators by habitat generalists or species favored by anthropogenic disturbances [15–19]. Moreover, beyond a certain level of forest loss, such ‘regime shifts’ are expected to be difficult to reverse even with major restoration efforts [11••]. Therefore, it is urgent to determine how much forest cover is needed to conserve forest-dependent pollinators [7, 11••, 20, 21]. There is an equally important need to understand how the structure, composition, and age of forests affect pollinators.

As forest succession proceeds, forest structure and composition change and the canopy can, depending on tree composition, become an increasingly important source of flowers and other resources for pollinators. Changes in structure with time depend largely on prevailing disturbance regimes which vary with region and forest type, topography, climate, human activity, among other factors [22, 23]. Whereas some mature forests are characterized by mostly closed canopies, others are not. More open structures occur in mature forests when disturbances kill sufficient numbers of dominant and co-dominant trees [24, 25]. Changes in forest composition include the predictable succession of tree species (e.g., shade-tolerant taxa replacing shade-intolerant taxa) as well as pests, diseases, natural disturbances, and management activities that select for a subset of tree species. While it is clear that pollinators respond to structural and compositional changes in forests, as reviewed here, inconsistent findings among studies suggest that such effects vary with forest type. Most notably, open stand conditions appear to be particularly important to pollinators when forests are dominated by trees that provide few floral resources [26, 27••, 28••].

The objectives of this article are to 1) provide a brief overview of global changes in forest cover, structure, composition, and age, 2) review how pollinator communities are affected by forest structure, composition, and age, 3) explore how these patterns may vary among forest types, particularly gymnosperm- vs. angiosperm-dominated forests (hereafter conifer vs. broadleaf, respectively), and 4) provide recommendations for managers interested in optimizing forest conditions for pollinator conservation. Efforts to define the term ‘forest’ are challenged by the variability in forest structure and composition observed globally. Therefore, to be as inclusive as possible we use the term ‘forest’ to refer to any ecosystem consisting of arborescent woody plants, regardless of their size or spacing. We adopt a similarly broad definition of ‘pollinator’ to refer to any flower-visiting insect that

has the potential to contribute to pollination. Although most of the literature on insect pollinators has focused on bees, non-bee insects can enhance pollination beyond what can be achieved by bees alone and are the primary pollinators of many plant species [5, 29]. Finally, this review aims to be global in coverage but we note that most recent research on these topics comes from work conducted in North America and Europe.

## Global Changes in Forest Structure, Composition, and Age

Forests throughout the world continue to experience changes in structure, composition, and age, with important implications for biodiversity. Globally, forests encompass ~31% (~4.1 billion ha) of land area [30] with tropical forests representing the largest proportion (45%), followed by boreal (27%), temperate (16%), and subtropical (11%) forests. The amount of forest has been declining globally although the rate of decline (net forest loss) has slowed to ~4.7 million ha yr<sup>-1</sup> [30]. This trend is influenced by many factors, including declines in deforestation rates in some countries and natural expansions of forests in others. Most recently, there have been several regional, national, and global initiatives to plant more trees, such as the The Trillion Tree Campaign which aims to plant one trillion trees worldwide as a nature-based solution to combat climate change [31].

While the loss of forests, such as through agricultural expansion [32], poses a major threat to global biodiversity, forest degradation, defined as the loss of key forest attributes that reduce the capacity of a forest to provide goods, services, and biodiversity [33], is also of concern, particularly in tropical regions [34]. Forest degradation often occurs in addition to forest loss, but this is not always the case. In the U.S., for example, forest cover has remained relatively stable over the past century [35] but a variety of forces are acting to reshape the structure and composition of these forests. These include non-native species, climate change, urbanization, and expansion of the wildland-urban interface (i.e., where houses and other developments meet or mix with undeveloped natural areas) [36–38].

Globally, old forests tend to be concentrated in regions that are cold and dry or warm and wet, whereas those of intermediate ages span large climatic gradients [39]. In North America, forests in the southeastern U.S. and in parts of central Canada are relatively young, reflecting the effects of industrial forest practices in the southeastern U.S. and the frequency of natural disturbances in central Canada [40]. A recent inventory in the U.S. found 64% of the ~58.4 million ha of forest managed by the USDA Forest Service were mature or old-growth [41]. Young forests (≤30 yrs old) comprise ~25% of forests in the U.S. whereas very old

forests ( $\geq 200$  years old) comprise  $\sim 4\%$  (calculations based on USDA Forest Service Forest Inventory and Analysis data, for all ownerships).

Climate changes driven by anthropogenic forces—including warming and altered precipitation patterns—can have direct and indirect effects on forests. Warming alters the phenology of vegetation (e.g., the timing of flowering and leaf out) and competitive relationships among species [42] and influences functional diversity, which has been shown to affect the vigor, growth, and survival of trees in some forest types [43, 44]. Many indirect effects of climate change are mediated through changes in the frequency and severity of disturbances that shape forest ecosystems by driving changes in vegetation growth, mortality, regeneration, and recruitment. Globally, approximately 98 million ha of forest were impacted by wildfire in 2015 [30], with more than two-thirds occurring in Africa and South America. Insects, diseases, and severe weather events damaged  $\sim 40$  million ha of forests in 2015 [30], mainly in temperate and boreal regions. In western North America, tens of millions of ha were impacted by bark beetle (Curculionidae: Scolytinae) outbreaks during 2000–2020 due, in part, to warming and drought with billions of trees killed [45]. Exotic, invasive insects and diseases have greatly altered the structure and composition of some forests and are a growing conservation concern [46]. Invasive tree diseases, including chestnut blight and Dutch elm disease in North America (caused by *Cryphonectria parasitica* and *Ophiostoma ulmi*, respectively), have threatened the existence of entire tree genera. With climate change, novel forest ecosystems are becoming increasingly prevalent and characterized by forest conditions, environmental drivers, and disturbance regimes with no or few past analogs [47]. Continued warming and changes in precipitation patterns are expected to amplify interactions among forest disturbances. However, vulnerability of the world's forests varies due to differences in biophysical conditions, disturbances, and local and regional variations in climate [48].

Humans play a significant role in determining the distribution, structure, composition, and age of forests. Numerous social and economic drivers influence how, when, and where forests are managed for goods and services, with each of these decisions having the potential to affect pollinator conservation. Seventy-three percent of the world's forests are publicly owned, with the share of publicly owned forests declining globally since 1990 [20]. In the U.S., more than half of forest land is privately owned; most private ownership occurs in the eastern U.S. with most western U.S. forests publicly owned. Increases in landowner turnover rates, parcellation of forest properties, and forest fragmentation are important conservation concerns (due to habitat loss, among other factors) in the eastern U.S. and likely to intensify in the future [49]. Globally, tropical forests experienced the

most fragmentation between 2000 and 2020. However, 75% of the world's forests experienced a decrease in fragmentation between 2000 and 2020, including highly fragmented temperate and subtropical regions in northern Eurasia and southern China [50]. In the future, forest landowners and natural resource managers will be increasingly challenged to maintain forests and associated goods and services, including pollination services, in the face of environmental and socioeconomic changes.

## Forest Structure

Forest structure—the horizontal and vertical arrangement of both living and dead vegetation—plays an important role in determining the amount and distribution of floral resources and breeding substrates used by pollinators. Here, we address the following three topics: 1) benefits and drivers of canopy openness, 2) effects of the shrub layer, and 3) the availability and suitability of breeding substrates. Although we discuss the effects of forest structure and composition on pollinators separately here, it should be noted that neither can be fully understood in isolation and the relative importance of each can be hard to disentangle [51].

Open canopies have increased floral herb abundance and diversity in the understory [52] and can have microclimates (e.g., temperature and soil moisture) that improve nesting habitat and enhance foraging conditions for pollinators [53–55]. The amount and diversity of floral resources in forests are highly linked to pollinator abundance and species richness [56, 57], including bees and wasps [58–62], flies [63], beetles [64], and butterflies [65]. Both natural disturbances and forest management activities can help maintain open canopies and enhance floral resource availability. The most common forest management treatments that increase canopy openness and floral resource availability in the understory include mechanized tree harvest and prescribed burning. Pollinator communities tend to respond positively to tree removal, especially in conifer forests where herbaceous floral diversity usually increases the first few years following thinning [27••, 66, 67]. For example, Davies et al. [68•] found that bee diversity and numbers of plant-bee interactions were greater in thinned versus unthinned ponderosa pine (*Pinus ponderosa*) stands in Colorado, U.S. Forest thinning in this system resulted in enhanced canopy openness and floral abundance in the understory, which were correlated with native bee abundance and diversity [68•]. In Europe, maintaining open habitats within forests (e.g., glades and clearings) and thinning forest stands is recommended to reverse recent declines in butterfly diversity [69]. Although thinning tends to benefit herbaceous floral abundance in the understory, thinning in some systems, such as tropical or temperate broadleaf forests, may reduce floral

resources for pollinators if it includes the removal of flowering trees or shrubs [70, 71].

Fire also creates more open canopies and increases floral abundance and diversity [72–75]. Some studies have shown that floral resources increase in response to the more open conditions following low-severity burns in both conifer and broadleaf forests, and that this results in higher pollinator diversity [76, 77•]. However, benefits of prescribed fire to floral and pollinator diversity tend to decline over time, sometimes within just a few years [78•]. For example, Gelles et al. [76] found that the highest floral diversity occurred one year after prescribed fire in ponderosa pine forests in Colorado whereas no difference in bee diversity was observed between burned and unburned stands after three years. In the U.S., a combination of mechanical thinning and prescribed fire has been shown in several studies to result in higher diversity of herbaceous plants and pollinators in temperate broadleaf and conifer forests [72, 77•, 79•, 80].

Both native and non-native species can form dense shrub layers that reduce the availability of flowers near the forest floor. Invasive non-native flowering plants can outcompete native plant species [81, 82] and may provide resources primarily for generalist and/or exotic pollinator species [83, 84]. Managing the shrub layer may help restore native plant and pollinator communities and their interactions [83, 85•, 86] (Fig. 1 D, E). However, this may not be the best course of action for flowering native shrubs [78•] and there is a shortage of research on the short and long-term consequences of removing invasive flowering species from forests [87–90].


In addition to floral resources, pollinators require nesting and overwintering sites which vary by taxa. Most bee species are ground-nesters and generally prefer exposed, malleable, dry soils at warm microsites with high light availability [91, 92]. While some forest bees are known to nest beneath leaf litter [93], the amount of bare soil is often positively correlated with abundance of ground-nesting bees [92]. This relationship has rarely been studied in forests, but in the southeastern U.S. prescribed fire has been shown to favor ground-nesting bees in pine (*Pinus* spp.) forests, likely via increases in bare soil [94, 95•], and soil-nesting bees declined with increasing depth of duff in broadleaf and conifer forests [96]. This trend seems to extend to other forest types [97, 98]. It is likely that most disturbances that create patches of bare soil could benefit bees that nest below ground. For example, thinning of the forest canopy can also promote bare soil, immediately via ground disturbance during harvesting and later through reduced litter deposition over time, and increase the amount of light reaching the forest floor with benefits to ground-nesting bees [55]. However, fire can compact soils which could limit the ability of bees to excavate nests [77•] and the extensive use of heavy (logging) equipment may have a similar effect [99], although this is not always the case [98].

Other pollinators, including many beetles (e.g., Cerambycidae, Scarabaeidae, Mordellidae) and hover flies (Syrphidae) require decaying wood for larval development, including that provided by tree hollows [100, 101]. While more structurally complex forests are hypothesized to increase nesting opportunities and larval feeding sites for pollinators [6••, 77•, 102, 103], relatively few studies have explicitly examined the effects of forest structure on nesting resources for pollinators. Abundance of dead wood (i.e., snags, stumps, and coarse woody debris) has been repeatedly linked to increases in cavity-nesting bees across a wide range of global forest types [6••, 79•, 96, 97, 104–106]. This may also be true for hoverflies and saproxylic beetles which also benefit from structural mosaics, increased canopy cover, and the presence of old and decadent trees [105, 107, 108]. Although some disturbances (e.g., high-severity wildfire) might be expected to reduce the availability of dead wood, several studies have found that woody debris and nesting opportunities for bees increase after severe wildfires [66, 79•, 109]. The effects of more exposed conditions on the suitability of breeding substrates for saproxylic pollinators probably vary among taxa. Although many beetle species that nest in dead wood benefit from sun exposure [110], more open conditions may negatively affect species dependent on wetter microhabitats, including many fly taxa [111–113]. More research is needed to clarify these relationships.

## Forest Composition

In addition to defining which floral and nesting resources are available, the composition of forest vegetation also determines how such resources are distributed across space and time. For example, forest canopies can either represent zones of floral abundance or scarcity depending on the overstory tree community. Moreover, climatic adaptations can result in forest plant assemblages characterized by highly seasonal periods of blooming [17, 114]. Here, we review four topics of particular relevance to managers: 1) the relative value of forests dominated by conifers vs. broadleaf trees, 2) the relationship between pollinator diversity and tree diversity, 3) the conservation value of plantation forests, and 4) the effects of non-native plants, insects, and pathogens on forest pollinators.

The first trees were gymnosperms and their descendants still dominate about 32% of global forest cover [115]. They remain particularly widespread at high latitudes and high elevations, as well as in regions experiencing frequent fire. Conifer forests are characterized by relatively low plant diversity, especially in the canopy, although some open fire-maintained forests (e.g., longleaf pine, *Pinus palustris*) have some of the highest understory plant diversity outside of the tropics, including many endemic species [116].



**Fig. 1** Pollinator diversity changes with forest age, often declining as young stands (A) develop into dense stands (B) during the stem exclusion phase [26]. Pollinator diversity can then increase as forests mature, especially if fire or other disturbances maintain open understories (C) [26] or if floral resources are provided by the overstory trees [120]. Midstory thickets formed by invasive shrubs such as Chinese privet (*Ligustrum sinense*) (D) can greatly reduce floral resource

availability near the forest floor [85•]. Efforts to eradicate such plants benefit flowering plants in the understory (E) as well as pollinators [85•]. Plantation forests such as this Monterey pine (*Pinus radiata*) plantation in New Zealand (F) are less structurally and compositionally complex than native forests (G), with important implications for biodiversity (photos taken by Michael Ulyshen)

By contrast, the canopies of broadleaf forests commonly consist of dozens to hundreds of tree species per ha [117], including both insect- and wind-pollinated taxa. The few comparisons of pollinator diversity between conifer- and broadleaf-dominated forests suggest diversity tends to be higher in broadleaf forests. For example, this was shown for butterflies in India [118], and, more recently, work from the southeastern U.S. (where both forest types occur naturally) found that bee diversity is negatively correlated with the amount of pine forest in the surrounding landscape [119] as well as the proportion of pine locally [120]. Such findings may reflect differences in pollen quality between coniferous and broadleaf trees. Although a number of studies have reported conifer pollen on bees or in their nest provisions [121, 122••], this is typically in very small quantities [123] and may represent contamination rather than active collection [124]. Furthermore, other work suggests conifer pollen may be a less nutritious and less preferred resource than that of many angiosperm taxa [125]. Despite evidence that conifers provide less favorable floral resources to pollinators than broadleaf trees, it is important to stress that the value of conifer forests is mediated by canopy openness. As discussed in the previous section, open conifer forests are characterized by abundant floral resources in the understory and support diverse pollinator assemblages [126]. Moreover, in regions where both open conifer forests and broadleaf forests occur together, such as the southeastern U.S., the two forest types support distinct pollinator communities [119], indicating that both are important to regional diversity.

The relationship between overstory tree diversity and pollinator diversity remains understudied. Several studies reported no strong relationship between tree diversity and the richness of cavity-nesting bees [62, 127]. However, few bee species were collected in those studies and more comprehensive sampling efforts suggest tree diversity may indeed be important to bee diversity. In England, for example, the presence of flowering *Acer pseudoplatanus* greatly increased bee activity in the forest canopy [128•]. In forests of Ohio and Indiana, U.S., Cunningham-Minnick and Crist [83] reported a positive correlation between spring bee diversity and the diversity of woody plants. Moreover, in the southeastern U.S., Traylor et al. [120] found bee diversity to be positively correlated with the number of insect-pollinated broadleaf tree species nearby. Similar patterns were reported for hoverflies in a beech (*Fagus*) forest in France [107]. Although this question has not, to our knowledge, been specifically tested for other groups of pollinating insects, the diversity of beetles generally increases with increasing tree species diversity [129] and this is also likely to be the case for Lepidoptera given their host specificity as larvae. For pollen-limited tree species, pollinator abundance and diversity may be critical for maintaining fruit and seed set [130].

Planted forests cover ~294 million ha of which 131 million ha (~3% of global forest cover) consist of plantations focused on the production of goods (wood or fiber) or providing ecosystem services such as protection from desertification, wind, soil degradation, or other concerns [30]. Because the total area of plantation forests continues to increase, often replacing native forests or naturally treeless habitats, there is an urgent need to better understand how they affect biodiversity [30, 131, 132]. Globally, the most widely planted tree genera are *Pinus*, *Cunninghamia*, *Eucalyptus*, *Populus*, and *Acacia* [133]. In some parts of the world, such as North and Central America, plantations consist mostly of native species (e.g., *Pinus*) whereas in other areas, such as South America and New Zealand, most plantations consist of introduced species (e.g., *Eucalyptus*, *Pinus radiata*) [30] (Fig. 1F). Despite supporting pollinators in some landscapes, there is agreement among studies that intensively managed non-native plantations harvested on short rotations are typically of limited benefit to pollinators [134]. This is true even when the planted trees produce flowers likely to attract pollinators. For example, poor seed production observed in both *Eucalyptus* and *Acacia* plantations suggests inadequate flower visitation, possibly resulting from pesticides or cultivation practices that reduce foraging or nesting resources within and around plantations [135]. Moreover, as with any monoculture, the benefits of flowering plantation trees are limited to the flowering period of that individual species, which may be short-lived. Thus, it is critical to provide other sources of flowers nearby. This can include preserving semi-natural habitats and native trees [136–139], planting a mix of tree species [134], maintaining unplanted open spaces [140], harvesting in ways that result in uneven-aged stands [141], and employing thinning practices that increase the availability of flowers in the understory of forest plantations [142, 143].


The intentional or accidental introduction of non-native plant species has led to marked alterations of forest composition worldwide. In addition to the 44% of all plantation forests consisting of introduced species [30], many other non-native species of trees, shrubs, grasses, and herbs are naturalized or invasive. Furthermore, introduced insects and pathogens have caused strong changes in overstory tree composition, including the extirpation of mass-flowering trees (e.g., American chestnut, *Castanea dentata*) that were historically of great importance to pollinators [144]. While the effects of some introduced plants may be neutral or even positive [145, 146], others, such as certain woody shrubs, negatively affect native plants in the understory [147] and can pose a threat to entire forest ecosystems by preventing tree regeneration [85•]. Controlling such species is critical to restoring conditions for pollinators in invaded forests.

## Forest Age

Forests undergo pronounced changes in both structure and composition as they age with a general trend toward greater complexity. As traditionally understood, there are four main stages of forest development [148]. These are the stand initiation stage, the stem exclusion stage when trees compete for space, the understory reinitiation stage when forest herbs and shrubs return to the understory, and the old growth stage when overstory trees begin to die. For the purposes of this review, it is helpful to add a fifth stage, marking the maturity and flowering of overstory trees, which occurs between the stem exclusion stage and the old growth stage. In addition to flowering forest trees and other plants, there is a proliferation of nesting and breeding microhabitats, such as tree hollows and dead wood, as trees age, senesce, and die. In most forests, natural disturbances result in an uneven age structure including patches of early successional habitat at landscape scales.

Comparisons of pollinator diversity among forests of different ages suggest conifer- and broadleaf-dominated forests exhibit different patterns due to variation in both disturbance regimes and availability of resources within the forest canopy. In conifer systems managed with regular prescribed fire (e.g., every few years), pollinator diversity appears to be unrelated to forest age [28••, 149, but see 150]. For example, Dixon et al. [28••] found no significant difference in bee richness among restored pine savannas across a gradient of stand ages (5–15, 15–30, 20–50, 50–75, and 75–100 yrs) on the southeastern U.S. Coastal Plain that were burned at two-year intervals. However, the same study [28••] and work by Hanula et al. [26] found bee diversity to be significantly lower in denser unburned pine stands compared to fire-maintained pine savannas. In Japan, Taki et al. [67] reported steep declines in bee richness with age in conifer plantations and Zitomer et al. [27••] reported similar results from northwestern North America where bee diversity declined by almost 50% for every five years since anthropogenic disturbance (i.e., clearcut harvest) in early successional, intensively managed Douglas-fir (*Pseudotsuga menziesii*) stands. Together, these studies suggest that conifer forests are of limited value to pollinators unless overstory tree density is reduced by fire or other disturbances (Figs. 1 A-C, Fig. 2). They also lend support to the idea that conifer trees provide insufficient floral rewards within the canopy to compensate for shaded understories. It is possible that old conifer forests characterized by uneven-aged structure and more canopy openings provide value to pollinators [150], consistent with the U-shaped curve described by Hilmers et al. [151], but more research is needed to evaluate this idea.

Unlike conifer forests, the canopies of broadleaf forests become increasingly valuable sources of floral resources as



**Fig. 2** Hypothesized relationships between pollinator diversity and forest age predict strong differences between conifer- and broadleaf-dominated forests. In the absence of disturbances, pollinator diversity in conifer forests declines rapidly with canopy closure and as the amount of light reaching the forest floor decreases [27••]. Reductions in conifer density—such as achieved with prescribed fire, thinning, or natural senescence—can restore pollinator diversity to levels comparable to that observed in more open habitats [26, 28••, 150, 151]. Although pollinator diversity declines similarly with time in young broadleaf forests [152•], it can increase as broadleaf trees mature and begin to flower, sometimes exceeding the diversity observed in open habitats [26]. This subsequent increase likely results from both the production of flowers in the canopy as well as nesting or breeding microhabitats essential to some forest-dependent pollinators. How pollinators may respond to thinning in broadleaf forests remains unknown. However, it is possible that reduced tree diversity and more xeric conditions may have a negative impact on some forest specialists. Given these differences, it is also hypothesized that a smaller proportion of pollinators in conifer forests are forest-dependent compared to broadleaf forests

trees age and begin to flower [122••]. Therefore, closed-canopy broadleaf forests can support high pollinator abundance and diversity, including many forest specialists. This may help explain why Taki et al. [67] reported higher pollinator diversity in mature broadleaf forests than in conifer forests of the same age in Japan. As in conifer forests, pollinator diversity declines from the open stand initiation stage to the stem exclusion stage when few flowers are available in the understory or canopy [67, 152•]. However, there is some support for the hypothesis that pollinator diversity increases as broadleaf trees mature and begin to produce flowers, and can reach levels exceeding what is observed in open habitats [26] (Fig. 2). In the southeastern U.S., Traylor et al. [120] reported that nearly half of the bee species examined were favored by older forests with a lower proportion of pine

(primarily *Pinus taeda*) in the canopy. Similar patterns are observed in tropical rainforests where diverse assemblages of stingless bees depend on old trees with cavities for nesting [153]. Thus, the relationship between pollinator diversity and forest age is hypothesized to differ between broadleaf and conifer forests (Fig. 2).

Another important consideration concerns changes in the composition of pollinator communities as forests age. Although little studied, pollinator composition can be expected to change strongly as forests transition from the youngest and most open stages to mature forests with flowering overstory trees and microhabitats required by forest-dependent species for breeding. While a considerable fraction of bee species in conifer forests depend on tree-derived resources (e.g., dead wood or resin) for nesting [126], the fact that conifer pollen is of comparatively limited value [124, 125] suggests that fewer pollinator species are dependent on conifer forests than on broadleaf forests. There is a strong need for research on changes in pollinator community composition as forests age, particularly work aimed at distinguishing between forest-dependent species and habitat generalists.

The differences described above regarding effects of forest age in broadleaf versus conifer forests have important implications for managers. Based on the information currently available, a stronger case can be made for prescribed burning or mechanically thinning to improve pollinator habitat in conifer forests as they age than in broadleaf forests. While thinning or burning broadleaf forests can be expected to increase floral resources near the ground, and this will likely favor many pollinator species, it is unclear how any associated changes in overstory tree diversity and composition may affect forest-dependent pollinators in these systems. Although a growing number of studies suggest the canopies of broadleaf forests are important to maintaining diverse pollinator assemblages, including forest-dependent species [10, 122••, 154•], much more research in this area is needed to anticipate how changes to the canopy are likely to impact this fauna at different stages of succession.

## Discussion

As we have shown, forest structure, composition, and age are highly interrelated and strongly influence the diversity and composition of pollinator assemblages. In many parts of the world, human activities are resulting in forests that bear little resemblance to historical conditions. Such changes can have negative outcomes for pollinators, particularly for forest specialists. For example, many countries in South America and Asia have implemented subsidies and other incentive programs for establishing forest plantations [155]. The displacement of older and more diverse native forests by plantation

monocultures of non-native tree species represent a loss of resources critical to native pollinator diversity. Other negative effects of human activities in forests include the alteration of historic disturbance regimes, such as fire exclusion, which has negative consequences for many plant species and pollinators that are adapted to disturbance. Fortunately, steps can be taken to help remedy the situation. Below we highlight some general recommendations for managers interested in improving conditions for pollinators in forested landscapes and provide suggestions for future research.

It is clear that the complexity of both forest structure and composition promotes insect diversity, including pollinators, with forest-dependent species benefiting the most. Most forests naturally become more structurally and compositionally complex as they age, although plant diversity can peak at mid-successional stages [156], with some critical resources being largely restricted to older forests [157]. While some of these resources, like large hollow-bearing trees and dead wood, can be found in younger forests and occasionally in other land cover types, older forests provide the critical benefit of ensuring temporal continuity in the availability of these resources. Thus, the conservation of old forests, where possible, will help ensure adequate resource availability for many sensitive species over time and space [158].

One characteristic typical of older forests is an uneven-aged stand structure maintained by regular natural disturbances such as wind events, wildfires, and outbreaks of forest insects and diseases [157]. Such disturbances are critical for maintaining forest biodiversity [156], so management that permits or emulates such disturbances at historic levels should improve conditions for pollinators. For example, the value of mature pine forests in the southeastern U.S. to pollinators is largely a function of fire frequency. Forests that burn infrequently tend to have higher tree densities as well as thicker understories that have fewer flowering species and low pollinator abundance and diversity [26]. By contrast, frequently burned forests have greater floral resource availability and more pollinators near the forest floor [26, 28••]. Timber harvest can also benefit pollinators by creating areas of good forage. Even clearcutting can benefit forest-dependent species provided there is suitable forest cover nearby [20]. Thus, it is important to ensure adequate availability of key resources throughout the landscape at appropriate temporal and spatial scales. For example, because most stingless bee species in the tropics nest in large cavity-bearing trees [159, 160], which are more common in older forests, it is important to incorporate old forests in management plans for this group.

Forests at stand initiation, understory reinitiation, and old growth stages of forest dynamics [148] often provide the greatest concentration of flowers within the herbaceous layer. Floral resources and pollinator diversity often decline from stand initiation to stem exclusion and will either remain low or increase

during understory reinitiation depending on the resources provided by the maturing trees in the canopy and on stand openness [161]. Maintaining open conditions may be more critical for pollinators in conifer forests than in broadleaf forests which often include mass-flowering canopy trees. While it may be possible that managing for more open canopies will benefit pollinators in all forest types, it remains poorly understood how practices that favor certain tree species over others, such as selective harvests or prescribed fire, may affect pollinator diversity within the canopies of mixed broadleaf-dominated forests. Future research aimed at clarifying the importance of different overstory tree species to forest-dependent taxa, as well as their response to alternative management actions, would be beneficial. There is a particularly strong need for more pollinator research in broadleaf forests.

A guiding principle in forest management is to mimic the disturbance regime to which an ecosystem is adapted. In some cases, conifer forests commonly experience more frequent or more widespread disturbances than broadleaf forests, such as wildfire or insect outbreaks. Disturbances are also common in broadleaf forests but often result in localized canopy gaps that arise from small-scale tree mortality events. Small clearings within regions extensively covered by broadleaf forests can be especially important to pollinators [78•], especially during times of the year when floral resources within closed-canopy forests are lacking. It is possible that more widespread thinning operations in broadleaf forests could continue to provide adequate resources for forest-dependent species while also improving conditions for habitat generalists. However, this should not be based on patterns observed in conifer forests given the many dissimilarities between these forest types (Fig. 2).

In plantations, incorporating native tree species and, when possible, planting a combination of tree species, can be expected to be of greater benefit to pollinators than monocultures of non-native tree species. As with any other crop, the negative effects of plantations on pollinators can also be ameliorated by protecting patches of native forests or other semi-natural habitats at regular intervals across the landscape [139].

## Conclusions

As forest loss and degradation continue throughout much of the world, there is an urgent need to understand and satisfy the resource requirements of species endemic to each region. Because pollinators are essential to global food security, it is in our best interest to protect them and the ecosystems upon which they depend. Although the abundance and diversity of pollinators are declining in many parts of the world [162], including in some forests [163], steps can be taken to improve conditions for these organisms. It is particularly

important to recognize that there is no substitute for semi-natural areas, and it is imperative that an adequate amount of such areas is conserved if we are to minimize local losses of endemic species. While targets remain poorly defined [21], a number of studies suggest strong declines in forest-dependent pollinators can be expected at <20% native forest cover [15, 18, 19, 164] and that considerably more than that is needed to avoid declines in some taxa [20]. Such targets need to be developed separately for different regions and forest types and depending on conservation priorities.

It should be stressed that the amount of forest cover necessary to support endemic pollinator diversity will depend on the quality of forests for pollinator foraging and nesting. Native forests can generally be expected to be of greater value to pollinators than plantations of non-native tree species. Furthermore, the value of native forests to pollinators can be reduced through alterations of historic disturbance regimes, addition of invasive species, removal of hollow, moribund, or otherwise unmarketable trees, elimination of dead wood, and other interventions that result in less structurally and compositionally complex systems. Thus, in addition to protecting adequate forest cover, managers should be mindful of forest quality when working to conserve native pollinators in forests.

**Acknowledgements** We thank three anonymous reviewers for comments that greatly improved this manuscript. This research was supported in part by the USDA Forest Service. The findings and conclusions in this publication are those of the authors and should not be construed to represent any official USDA or U.S. Government determination or policy.

**Author Contributions** M.D.U. conceived of and coordinated the manuscript. All authors contributed sections of text and provided final edits. M.D.U. prepared the figures.

## Compliance with Ethical Standards

**Competing Interests** The authors declare that they have no competing interests.

**Human and Animal Rights and Informed Consent** This article does not contain any studies with human or animal subjects performed by any of the authors.

## References

Papers of particular interest, published recently, have been highlighted as:

- Of importance
- Of major importance

1. Irvine A, Armstrong J. Beetle pollination in tropical forests of Australia. In: Bawa KS, Hadley M, editors. *Reproductive Ecology of Tropical Forest Plants*. Park Ridge, NJ: The Parthenon Publishing Group Inc.; 1990. p. 135–49.

2. Roubik DW. Ecology and natural history of tropical bees. Cambridge University Press; 1992.
3. Bawa KS. Plant-pollinator interactions in tropical rain forests. *Annu Rev Ecol Syst.* 1990;21:399–422. <https://doi.org/10.1146/annurev.es.21.110190.002151>.
4. Birtele D, Hardersen S. Analysis of vertical stratification of Syrphidae (Diptera) in an oak-hornbeam forest in northern Italy. *Ecol Res.* 2012;27:755–63. <https://doi.org/10.1007/s11284-012-0948-2>.
5. Haber WA, Frankie GW. A tropical hawkmoth community: Costa Rican dry forest Sphingidae. *Biotropica.* 1989;21:155–72. <https://doi.org/10.2307/2388706>.
6. ●● Urban-Mead KR, Muñiz P, Gillung J, Espinoza A, Fordyce R, van Dyke M, et al. Bees in the trees: Diverse spring fauna in temperate forest edge canopies. *Forest Ecol Manag.* 2021;482:118903. <https://doi.org/10.1016/j.foreco.2020.118903>. **Provides evidence that trees provide important floral resources to bees in temperate deciduous forests.**
7. Rahimi E, Barghjeh S, Dong P. Amount, distance-dependent and structural effects of forest patches on bees in agricultural landscapes. *Agric Food Secur.* 2022;11:10. <https://doi.org/10.1186/s40066-022-00360-x>.
8. ● Mola JM, Hemberger J, Kochanski J, Richardson LL, Pearse IS. The importance of forests in bumble bee biology and conservation. *BioScience.* 2021;1(12):1234–48. <https://doi.org/10.1093/biosci/biab121>. **Reviews the importance of forests to bumble bee conservation.**
9. ●● Smith C, Harrison T, Gardner J, Winfree R. Forest-associated bee species persist amid forest loss and regrowth in eastern North America. *Biol Conserv.* 2021;260:109202. <https://doi.org/10.1016/j.biocon.2021.109202>. **Estimates the proportion of bees in the northeastern U.S. associated with forests.**
10. Dorey JB, Davies OK, Magnacca KN, Schwarz MP, Gilpin A-M, Ramage T, et al. Canopy specialist *Hylaeus* bees highlight sampling biases and resolve Michener’s mystery. *Front Ecol Evol.* 2024;12:1339446. <https://doi.org/10.3389/fevo.2024.1339446>.
11. ●● Ulyshen MD, Urban-Mead KR, Dorey J, Rivers JW. Forests are critically important to global pollinator diversity and enhance pollination in adjacent crops. *Biol Rev.* 2023;98:1118–41. **Establishes the importance of forests to global pollinator diversity.**
12. Ricketts TH, Daily GC, Ehrlich PR, Michener CD. Economic value of tropical forest to coffee production. *PNAS.* 2004;101(34):12579–82. <https://doi.org/10.1073/pnas.0405147101>.
13. ● Severiano-Galeana F, Rosas-Guerrero V, Alemán-Figueroa L, Lucas-García R, Almazán-Núñez RC, Kuk-Dzul JG. Orchards closer to forest patches produced fewer malformed fruits and more commercial fruits: The importance of legitimate floral visitors. *Agr Ecosyst Environ.* 2024;363:108872. <https://doi.org/10.1016/j.agee.2023.108872>. **Provides evidence that forest bees promote pollination in adjacent crops.**
14. Smith C, Weinman L, Gibbs J, Winfree R. Specialist foragers in forest bee communities are small, social or emerge early. *J Anim Ecol.* 2019;88:1158–67. <https://doi.org/10.1111/1365-2656.13003>.
15. Kerr WE, Carvalho GA, Nascimento VA. The probable consequences of the destruction of Brazilian stingless bees. *Adv Econ Bot.* 1999;13:395–403.
16. Collado MÁ, Sol D, Bartomeus I. Bees use anthropogenic habitats despite strong natural habitat preferences. *Divers Distrib.* 2019;25:924–35. <https://doi.org/10.1111/ddi.12899>.
17. Harrison T, Gibbs J, Winfree R. Forest bees are replaced in agricultural and urban landscapes by native species with different phenologies and life-history traits. *Glob Change Biol.* 2018;24:287–96. <https://doi.org/10.1111/gcb.13921>.
18. Hall MA, Nimmo DG, Cunningham SA, Walker K, Bennett AF. The response of wild bees to tree cover and rural land use is mediated by species’ traits. *Biol Conserv.* 2019;231:1–12. <https://doi.org/10.1016/j.biocon.2018.12.032>.
19. Liow LH, Sodhi NS, Elmqvist T. Bee diversity along a disturbance gradient in tropical lowland forests of south-east Asia. *J Appl Ecol.* 2001;38:180–92. <https://doi.org/10.1046/j.1365-2664.2001.00582.x>.
20. Viljur M-L, Tiitsaar A, Gimbutas M, Kaasik A, Valdma D, Õunap E, et al. Conserving woodland butterflies in managed forests: Both local and landscape factors matter. *Forest Ecol Manag.* 2020;462:118002. <https://doi.org/10.1016/j.foreco.2020.118002>.
21. Carroll C, Noss RF. How percentage-protected targets can support positive biodiversity outcomes. *Conserv Biol.* 2022;36:e13869. <https://doi.org/10.1111/cobi.13869>.
22. Guyette RP, Stambaugh MC, Dey DC, Muzika R-M. Predicting fire frequency with chemistry and climate. *Ecosystems.* 2012;15:322–35. <https://doi.org/10.1007/s10021-011-9512-0>.
23. Stanturf JA, Goodrick SL, Outcalt KW. Disturbance and coastal forests: A strategic approach to forest management in hurricane impact zones. *For Ecol Manag.* 2007;250:119–35. <https://doi.org/10.1016/j.foreco.2007.03.015>.
24. Scherer SS, D’Amato AW, Kern CC, Palik BJ, Russell MB. Long-term impacts of prescribed fire on stand structure, growth, mortality, and individual tree vigor in *Pinus resinosa* forests. *For Ecol Manag.* 2016;368:7–16. <https://doi.org/10.1016/j.foreco.2016.02.038>.
25. Quine CP, Gardiner BA, Moore J. Chapter Four - Wind disturbance in forests: The process of wind created gaps, tree overturning, and stem breakage. In: Johnson EA, Miyanishi K, editors. *Plant Disturbance Ecology*. 2nd ed. San Diego: Academic Press; 2021. p. 117–84.
26. Hanula JL, Horn S, O’Brien JJ. Have changing forests conditions contributed to pollinator decline in the southeastern United States? *Forest Ecol Manag.* 2015;348:142–52. <https://doi.org/10.1016/j.foreco.2015.03.044>.
27. ●● Zitomer RA, Galbraith SM, Betts MG, Moldenke AR, Progar RA, Rivers JW. Bee diversity decreases rapidly with time since harvest in intensively managed conifer forests. *Ecol Appl.* 2023;33(5):e2855. <https://doi.org/10.1002/eap.2855>. **Provides evidence that early successional forests provide important resources for bees in Pacific Northwestern forests.**
28. ●● Dixon CM, Robertson KM, Ulyshen MD, Sikes BA. Pine savanna restoration on agricultural landscapes: The path back to native savanna ecosystem services. *Sci Total Environ.* 2022;818:151715. <https://doi.org/10.1016/j.scitotenv.2021.151715>. **Shows the importance of open canopy conditions for bees in southeastern U.S. pine ecosystems.**
29. Rader R, Bartomeus I, Garibaldi LA, Garratt MPD, Howlett BG, Winfree R, et al. Non-bee insects are important contributors to global crop pollination. *PNAS.* 2016;113(1):146–51. <https://doi.org/10.1073/pnas.1517092112>.
30. FAO. *Global Forest Resources Assessment 2020: Main report*. Rome. 2020.
31. Lippke B, Puettmann M, Oneil E, Dearing OC. The plant a trillion trees campaign to reduce global warming – Fleshing out the concept. *J Sustain Forest.* 2021;40(1):1–31. <https://doi.org/10.1080/10549811.2021.1894951>.
32. Zabel F, Delzeit R, Schneider JM, Seppelt R, Mauser W, Václavík T. Global impacts of future cropland expansion and intensification on agricultural markets and biodiversity. *Nat Commun.* 2019;10(1):2844. <https://doi.org/10.1038/s41467-019-10775-z>.
33. Ghazoul J, Burivalova Z, Garcia-Ulloa J, King LA. Conceptualizing forest degradation. *Trends Ecol Evol.* 2015;30(10):622–32. <https://doi.org/10.1016/j.tree.2015.08.001>.

34. Lapola DM, Pinho P, Barlow J, Aragão LEOC, Berenguer E, Carmenta R, et al. The drivers and impacts of Amazon forest degradation. *Science*. 2023;379(6630):eabp8622. <https://doi.org/10.1126/science.abp8622>.
35. Oswalt SN, Smith WB, Miles PD, Pugh SA. Forest resources of the United States, 2017: A technical document supporting the Forest Service 2020 RPA Assessment. WO-GTR-97. Washington, DC: U.S. Department of Agriculture, Forest Service, Washington Office; 2019.
36. Radeloff VC, Helmers DP, Kramer HA, Mockrin MH, Alexandre PM, Bar-Massada A, et al. Rapid growth of the US wildland-urban interface raises wildfire risk. *PNAS*. 2018;115(13):3314–9. <https://doi.org/10.1073/pnas.1718850115>.
37. Mockrin MH, Helmers D, Martinuzzi S, Hawbaker TJ, Radeloff VC. Growth of the wildland-urban interface within and around U.S. National Forests and Grasslands, 1990–2010. *Landscape Urban Plan*. 2022;218:104283. <https://doi.org/10.1016/j.landurbplan.2021.104283>.
38. Stewart SI, Radeloff VC, Hammer RB, Hawbaker TJ. Defining the wildland–urban interface. *J Forest*. 2007;105:201–7. <https://doi.org/10.1093/jof/105.4.201>.
39. Besnard S, Koirala S, Santoro M, Weber U, Nelson J, Gütter J, et al. Mapping global forest age from forest inventories, biomass and climate data. *Earth Syst Sci Data*. 2021;13(10):4881–96. <https://doi.org/10.5194/essd-13-4881-2021>.
40. Pan Y, Birdsey RA, Phillips OL, Jackson RB. The structure, distribution, and biomass of the world's forests. *Annu Rev Ecol Evol S*. 2013;44(1):593–622. <https://doi.org/10.1146/annurev-ecolsys-110512-135914>.
41. USDA Forest Service. Mature and old-growth forests: Definition, identification, and initial inventory on lands managed by the Forest Service and Bureau of Land Management. FS-1215a. 2023.
42. Vitasse Y, François C, Delpierre N, Dufrêne E, Kremer A, Chuine I, et al. Assessing the effects of climate change on the phenology of European temperate trees. *Agr Forest Meteorol*. 2011;151(7):969–80. <https://doi.org/10.1016/j.agrformet.2011.03.003>.
43. Forrester DI, Ammer C, Annighöfer PJ, Barbeito I, Bielak K, Bravo-Oviedo A, et al. Effects of crown architecture and stand structure on light absorption in mixed and monospecific *Fagus sylvatica* and *Pinus sylvestris* forests along a productivity and climate gradient through Europe. *J Ecol*. 2018;106(2):746–60. <https://doi.org/10.1111/1365-2745.12803>.
44. Looney CE, Long JW, Fettig CJ, Fried JS, Wood KEA, Audley JP. Functional diversity affects tree vigor, growth, and mortality in mixed-conifer/hardwood forests in California, U.S.A., in the absence of fire. *Forest Ecol Manag*. 2023;544:121135. <https://doi.org/10.1016/j.foreco.2023.121135>.
45. Fettig CJ, Asaro C, Nowak JT, Dodds KJ, Gandhi K, Moan JE, et al. Trends in bark beetle impacts in North America during a period (2000–2020) of rapid environmental change. *J Forest*. 2022;120:693–713.
46. Poland TM, Patel-Weynand T, Finch DM, Miniati CF, Hayes DC, Lopez VM. Invasive species in forests and rangelands of the United States: A comprehensive science synthesis for the United States forest sector. Springer Nature; 2021.
47. Golladay SW, Martin KL, Vose JM, Wear DN, Covich AP, Hobbs RJ, et al. Achievable future conditions as a framework for guiding forest conservation and management. *Forest Ecol Manag*. 2016;360:80–96. <https://doi.org/10.1016/j.foreco.2015.10.009>.
48. Lecina-Diaz J, Martínez-Vilalta J, Alvarez A, Banqué M, Birkmann J, Feldmeyer D, et al. Characterizing forest vulnerability and risk to climate-change hazards. *Front Ecol Environ*. 2021;19(2):126–33. <https://doi.org/10.1002/fee.2278>.
49. Fettig CJ, Vose JM, et al. America's changing biophysical and socioeconomic context. In: Schuler TM, Patel-Weynand T, editors. Active management for resilient and productive forests WO-GTR-103. Washington, DC: US Department of Agriculture, Forest Service, Washington Office; 2024. <https://doi.org/10.2737/WO-GTR-103-Chap2>.
50. Ma J, Li J, Wu W, Liu J. Global forest fragmentation change from 2000 to 2020. *Nat Commun*. 2023;14:3752. <https://doi.org/10.1038/s41467-023-39221-x>.
51. Traylor CR, Ulyshen MD, Wallace D, Loudermilk EL, Ross CW, Hawley C, et al. Compositional attributes of invaded forests drive the diversity of insect functional groups. *Glob Ecol Conserv*. 2022;35:e02092. <https://doi.org/10.1016/j.gecco.2022.e02092>.
52. Radhamoni HVN, Queenborough SA, Arietta AZA, Suresh HS, Dattaraja HS, Kumar SS, et al. Local- and landscape-scale drivers of terrestrial herbaceous plant diversity along a tropical rainfall gradient in Western Ghats. *India J Ecol*. 2023;111:1021–36. <https://doi.org/10.1111/1365-2745.14075>.
53. Davis TS, Comai N. Canopy cover and seasonality are associated with variation in native bee assemblages across a mixed pine-juniper woodland. *Agr Forest Entomol*. 2022;24(4):497–505. <https://doi.org/10.1111/afe.12511>.
54. Hanula JL, Ulyshen MD, Horn S. Conserving pollinators in North American forests: A review. *Nat Area J*. 2016;36(4):427–39. <https://doi.org/10.3375/043.036.0409>.
55. Polatto LP, Chaud-Netto J, Alves-Junior VV. Influence of abiotic factors and floral resource availability on daily foraging activity of Bees. *J Insect Behav*. 2014;27(5):593–612. <https://doi.org/10.1007/s10905-014-9452-6>.
56. Ebeling A, Klein A-M, Schumacher J, Weisser WW, Tschardt T. How does plant richness affect pollinator richness and temporal stability of flower visits? *Oikos*. 2008;117(12):1808–15. <https://doi.org/10.1111/j.1600-0706.2008.16819.x>.
57. Hyjazie BF, Sargent RD. Floral resources predict the local bee community: Implications for conservation. *Biol Conserv*. 2022;273:109679. <https://doi.org/10.1016/j.biocon.2022.109679>.
58. Brooks DR, Nocera JJ. Bumble bee (*Bombus* spp.) diversity differs between forested wetlands and clearcuts in the Acadian forest. *Can J Forest Res*. 2020;50:1399–404. <https://doi.org/10.1139/cjfr-2020-0094>.
59. Klein A-M, Steffan-Dewenter I, Tschardt T. Rain forest promotes trophic interactions and diversity of trap-nesting Hymenoptera in adjacent agroforestry. *J Anim Ecol*. 2006;75:315–23. <https://doi.org/10.1111/j.1365-2656.2006.01042.x>.
60. Roulston TH, Goodell K. The role of resources and risks in regulating wild bee populations. *Annu Rev Entomol*. 2011;56:293–312. <https://doi.org/10.1146/annurev-ento-120709-144802>.
61. Rubene D, Schroeder M, Ranius T. Diversity patterns of wild bees and wasps in managed boreal forests: Effects of spatial structure, local habitat and surrounding landscape. *Biol Conserv*. 2015;184:201–8. <https://doi.org/10.1016/j.biocon.2015.01.029>.
62. Sobek S, Tschardt T, Scherber C, Schiele S, Steffan-Dewenter I. Canopy vs. understory: Does tree diversity affect bee and wasp communities and their natural enemies across forest strata? *Forest Ecol Manag*. 2009;258(5):609–15. <https://doi.org/10.1016/j.foreco.2009.04.026>.
63. Souza-Silva M, Fontenelle JCR, Martins RP. Seasonal abundance and species composition of flower-visiting flies. *Neotrop Entomol*. 2001;30:351–9.
64. Kirmse S, Chaboo CS. Flowers are essential to maintain high beetle diversity (Coleoptera) in a Neotropical rainforest canopy. *J Nat Hist*. 2020;54(25–26):1661–96. <https://doi.org/10.1080/00222933.2020.1811414>.

65. Gordon SCC, Kerr JT. Floral diversity increases butterfly diversity in a multitrophic metacommunity. *Ecology*. 2022;103(8):e3735. <https://doi.org/10.1002/ecy.3735>.
66. Galbraith SM, Cane JH, Moldenke AR, Rivers JW. Salvage logging reduces wild bee diversity, but not abundance, in severely burned mixed-conifer forest. *Forest Ecol Manag*. 2019;453: 117622. <https://doi.org/10.1016/j.foreco.2019.117622>.
67. Taki H, Okochi I, Okabe K, Inoue T, Goto H, Matsumura T, et al. Succession influences wild bees in a temperate forest landscape: The value of early successional stages in naturally regenerated and planted forests. *PLoS ONE*. 2013;8:e56678. <https://doi.org/10.1371/journal.pone.0056678>.
68. • Davies CB, Davis TS, Griswold T. Forest restoration treatments indirectly diversify pollination networks via floral- and temperature-mediated effects. *Ecol Appl*. 2023;3(8):e2927. <https://doi.org/10.1002/eap.2927>. **Presents evidence that thinning benefits bees in western U.S. conifer forests.**
69. Bubová T, Vrabec V, Kulma M, Nowicki P. Land management impacts on European butterflies of conservation concern: a review. *J Insect Conserv*. 2015;19(5):805–21. <https://doi.org/10.1007/s10841-015-9819-9>.
70. Coulin C, Aizen MA, Garibaldi LA. Contrasting responses of plants and pollinators to woodland disturbance. *Austral Ecol*. 2019;44(6):1040–51. <https://doi.org/10.1111/aec.12771>.
71. Ghazoul J. Alien Abduction: Disruption of native plant-pollinator interactions by invasive species. *Biotropica*. 2004;36(2):156–64. <https://doi.org/10.1111/j.1744-7429.2004.tb00308.x>.
72. Campbell JW, Hanula JL, Waldrop TA. Effects of prescribed fire and fire surrogates on floral visiting insects of the blue ridge province in North Carolina. *Biol Conserv*. 2007;134(3):393–404. <https://doi.org/10.1016/j.biocon.2006.08.029>.
73. Carbone LM, Tavella J, Pausas JG, Aguilar R. A global synthesis of fire effects on pollinators. *Global Ecol Biogeogr*. 2019;28(10):1487–98. <https://doi.org/10.1111/geb.12939>.
74. He T, Lamont BB, Pausas JG. Fire as a key driver of Earth's biodiversity. *Biol Rev*. 2019;94(6):1983–2010. <https://doi.org/10.1111/brev.12544>.
75. Potts SG, Vulliamy B, Dafni A, Ne'eman G, O'Toole C, Roberts S, et al. Response of plant-pollinator communities to fire: Changes in diversity, abundance and floral reward structure. *Oikos*. 2003;101:103–12. <https://doi.org/10.1034/j.1600-0706.2003.12186.x>.
76. Gelles RV, Davis TS, Barrett KJ. Prescribed fire is associated with increased floral richness and promotes short-term increases in bee biodiversity in the ponderosa pine forest of the Southern Rocky Mountains. *Agri Forest Entomol*. 2023;25(3):435–48. <https://doi.org/10.1111/afe.12565>.
77. • Chase MH, Fraterrigo JM, Charles B, Harmon-Threatt A. Wild bee response to forest management varies seasonally and is mediated by resource availability. *Forest Ecol Manag*. 2023;548:121426. <https://doi.org/10.1016/j.foreco.2023.121426>. **Reports that bees respond most strongly to a combination of thinning and burning.**
78. • Ulyshen M, Elliott K, Scott J, Horn S, Clinton P, Liu N, et al. Effects of Rhododendron removal and prescribed fire on bees and plants in the southern Appalachians. *Ecol Evol*. 2022;12(3):e8677. <https://doi.org/10.1002/ece3.8677>. **Documents the concentration of bees in small clearings compared to neighboring closed canopy forests in the Southern Appalachians.**
79. • Gelles RV, Davis TS, Stevens-Rumann CS. Wildfire and forest thinning shift floral resources and nesting substrates to impact native bee biodiversity in ponderosa pine forests of the Colorado Front Range. *Forest Ecol Manag*. 2022;510:120087. <https://doi.org/10.1016/j.foreco.2022.120087>.
80. Lettow MC, Brudvig LA, Bahlai CA, Gibbs J, Jean RP, Landis DA. Bee community responses to a gradient of oak savanna restoration practices. *Restor Ecol*. 2018;26(5):882–90. <https://doi.org/10.1111/rec.12655>.
81. Daehler CC. Performance comparisons of co-occurring native and alien invasive plants: Implications for conservation and restoration. *Annu Rev Ecol Evol S*. 2003;34(1):183–211. <https://doi.org/10.1146/annurev.ecolsys.34.011802.132403>.
82. Dietzsch AC, Stanley DA, Stout JC. Relative abundance of an invasive alien plant affects native pollination processes. *Oecologia*. 2011;167:469–79. <https://doi.org/10.1007/s00442-011-1987-z>.
83. Cunningham-Minnick MJ, Crist TO. Floral resources of an invasive shrub alter native bee communities at different vertical strata in forest-edge habitat. *Biol Inv*. 2020;22(7):2283–98. <https://doi.org/10.1007/s10530-020-02248-y>.
84. Morales CL, Aizen MA. Does Invasion of exotic plants promote invasion of exotic flower visitors? A case study from the temperate forests of the Southern Andes. *Biol Inv*. 2002;4(1):87–100. <https://doi.org/10.1023/A:1020513012689>.
85. • Ulyshen MD, Horn S, Hanula JL. Decadal patterns of forest and pollinator recovery following the eradication of an invasive shrub. *Front Ecol Evol*. 2022;10:832268. <https://doi.org/10.3389/fevo.2022.832268>. **Demonstrates the lasting value of efforts to remove invasive shrubs on forest pollinator communities.**
86. McKinney AM, Goodell K. Plant–pollinator interactions between an invasive and native plant vary between sites with different flowering phenology. *Plant Ecol*. 2011;212(6):1025–35. <https://doi.org/10.1007/s11258-010-9882-y>.
87. Drossart M, Michez D, Vanderplanck M. Invasive plants as potential food resource for native pollinators: A case study with two invasive species and a generalist bumble bee. *Sci Rep*. 2017;7(1):16242. <https://doi.org/10.1038/s41598-017-16054-5>.
88. Kovács-Hostyánszki A, Szigeti V, Miholcsa Z, Sándor D, Soltész Z, Török E, et al. Threats and benefits of invasive alien plant species on pollinators. *Basic Appl Ecol*. 2022;64:89–102. <https://doi.org/10.1016/j.baee.2022.07.003>.
89. Stout JC, Tiedeken EJ. Direct interactions between invasive plants and native pollinators: Evidence, impacts and approaches. *Funct Ecol*. 2017;31(1):38–46. <https://doi.org/10.1111/1365-2435.12751>.
90. Vilà M, Bartomeus I, Dietzsch AC, Petanidou T, Steffan-Dewenter I, Stout JC, et al. Invasive plant integration into native plant-pollinator networks across Europe. *Proc R Soc Lond B Biol Sci*. 2009;276(1674):3887–93. <https://doi.org/10.1098/rspb.2009.1076>.
91. Antoine CM, Forrest JRK. Nesting habitat of ground-nesting bees: A review. *Ecol Entomol*. 2021;46(2):143–59. <https://doi.org/10.1111/een.12986>.
92. Harmon-Threatt A. Influence of nesting characteristics on health of wild bee communities. *Annu Rev Entomol*. 2020;65(1):39–56. <https://doi.org/10.1146/annurev-ento-011019-024955>.
93. Ulyshen M, Traylor CR, Danforth BN. Patterns of nest-site selection by *Colletes thoracicus* within a forested watershed. *Apidologie*. 2023;54:56. <https://doi.org/10.1007/s13592-023-01035-7>.
94. Moylett H, Youngsteadt E, Sorenson C. The impact of prescribed burning on native bee communities (Hymenoptera: Apoidea: Anthophila) in longleaf pine savannas in the North Carolina sandhills. *Environ Entomol*. 2019;49(1):211–9. <https://doi.org/10.1093/ee/nvz156>.
95. • Ulyshen MD, Wilson AC, Ohlson GC, Pokswinski SM, Hiers JK. Frequent prescribed fires favour ground-nesting bees in southeastern U.S. forests. *Insect Conserv Diver*. 2021;14(4):527–34. <https://doi.org/10.1111/icad.12484>. **Shows that frequent burns improve habitat for many ground-nesting bees in southeastern U.S. conifer forests.**
96. Fortuin CC, Gandhi KJK. Functional traits and nesting habitats distinguish the structure of bee communities in clearcut and managed hardwood & pine forests in Southeastern USA. *Forest Ecol Manag*. 2021;496:119351. <https://doi.org/10.1016/j.foreco.2021.119351>.

97. Potts SG, Vulliamy B, Roberts S, O'Toole C, Dafni A, Ne'eman G, et al. Role of nesting resources in organising diverse bee communities in a Mediterranean landscape. *Ecol Entomol.* 2005;30(1):78–85. <https://doi.org/10.1111/j.0307-6946.2005.00662.x>.
98. Rivers JW, Mathis CL, Moldenke AR, Betts MG. Wild bee diversity is enhanced by experimental removal of timber harvest residue within intensively managed conifer forest. *GCB Bioenergy.* 2018;10(10):766–81. <https://doi.org/10.1111/gcbb.12531>.
99. Christmann S. Regard and protect ground-nesting pollinators as part of soil biodiversity. *Ecol Appl.* 2022;32(3):e2564. <https://doi.org/10.1002/eap.2564>.
100. Ulyshen MD, editor. Saproxylic insects: Diversity, Ecology and Conservation. Cham, Switzerland: Springer; 2018.
101. Traylor CR, Ulyshen MD, McHugh JV. New host records and a review of deadwood associations for Mordellidae (Coleoptera) in North America. *Coleopta Bull.* 2023;77:375–81. <https://doi.org/10.1649/0010-065X-77.3.375>.
102. Flores LMA, Zanette LRS, Araujo FS. Effects of habitat simplification on assemblages of cavity nesting bees and wasps in a semiarid neotropical conservation area. *Biodivers Conserv.* 2018;27(2):311–28. <https://doi.org/10.1007/s10531-017-1436-3>.
103. Rodríguez A, Kouki J. Disturbance-mediated heterogeneity drives pollinator diversity in boreal managed forest ecosystems. *Ecol Appl.* 2017;27(2):589–602. <https://doi.org/10.1002/eap.1468>.
104. Glennly W, Runyon JB, Burkle LA. Habitat characteristics structuring bee communities in a forest-shrubland ecotone. *Forest Ecol Manag.* 2023;534:120883. <https://doi.org/10.1016/j.foreco.2023.120883>.
105. Proesmans W, Bonte D, Smagghe G, Meeus I, Decocq G, Spicher F, et al. Small forest patches as pollinator habitat: Oases in an agricultural desert? *Landscape Ecol.* 2019;34(3):487–501. <https://doi.org/10.1007/s10980-019-00782-2>.
106. Vázquez DP, Alvarez JA, Debandi G, Aranibar JN, Villagra PE. Ecological consequences of dead wood extraction in an arid ecosystem. *Basic Appl Ecol.* 2011;12(8):722–32. <https://doi.org/10.1016/j.baae.2011.08.009>.
107. Larriue L, Cabanettes A, Sarthou JP. Hoverfly (Diptera: Syrphidae) richness and abundance vary with forest stand heterogeneity: Preliminary evidence from a montane beech fir forest. *Eur J Entomol.* 2015;112:755–69. <https://doi.org/10.14411/eje.2015.083>.
108. Ricarte A, Jover T, Marcos-García MA, Micó E, Brustel H. Saproxylic beetles (Coleoptera) and hoverflies (Diptera: Syrphidae) from a Mediterranean forest: Towards a better understanding of their biology for species conservation. *J Nat Hist.* 2009;43(9–10):583–607. <https://doi.org/10.1080/00222930802610527>.
109. Simanonok MP, Burkle LA. Nesting success of wood-cavity-nesting bees declines with increasing time since wildfire. *Ecol Evol.* 2019;9(22):12436–45. <https://doi.org/10.1002/ece3.5657>.
110. Seibold S, Bässler C, Brandl R, Büche B, Szallies A, Thorn S, et al. Microclimate and habitat heterogeneity as the major drivers of beetle diversity in dead wood. *J Appl Ecol.* 2016;53:934–43. <https://doi.org/10.1111/1365-2664.12607>.
111. Ulyshen MD. Saproxylic Diptera. In: Ulyshen MD, editor. Saproxylic Insects: Diversity, Ecology and Conservation. Cham: Springer International Publishing; 2018. p. 167–92.
112. Jonsell M, Nordlander G, Ehnström B. Substrate associations of insects breeding in fruiting bodies of wood-decaying fungi. *Ecol Bull.* 2001;49:173–94.
113. Rotheray GE, Hancock G, Hewitt S, Horsfield D, MacGowan I, Robertson D, et al. The Biodiversity and Conservation of Saproxylic Diptera In Scotland. *J Insect Conserv.* 2001;5:77–85. <https://doi.org/10.1023/A:1011329722100>.
114. Heinrich B. Flowering phenologies: bog, woodland, and disturbed habitats. *Ecology.* 1976;57:890–9.
115. FAO. The State of the World's Forests 2020. Forests, biodiversity and people. Rome; 2020.
116. Peet RK. Ecological classification of longleaf pine woodlands. In: Jose S, Jokela EJ, Miller DL, editors. The longleaf pine ecosystem: Ecology, silviculture, and restoration. New York, NY: Springer New York; 2006. p. 51–93.
117. Phillips OL, Hall P, Gentry AH, Sawyer SA, Vasquez R. Dynamics and species richness of tropical rain forests. *PNAS.* 1994;91:2805–9. <https://doi.org/10.1073/pnas.91.7.2805>.
118. Kunte K, Joglekar A, Utkarsh G, Padmanabhan P. Patterns of butterfly, bird and tree diversity in the Western Ghats. *Curr Sci.* 1999;77(4):577–86.
119. Ulyshen MD, Adams C, Adams J, Adams SB, Bland M, Bragg DC, et al. Spatiotemporal patterns of forest pollinator diversity across the southeastern United States. *Divers Distrib.* 2024;e13869. <https://doi.org/10.1111/ddi.13869>.
120. Traylor CR, Ulyshen MD, Bragg DC, McHugh JV. Forest bees benefit from compositionally diverse broadleaf canopies. *For Ecol Manag.* 2024;566:122051. <https://doi.org/10.1016/j.foreco.2024.122051>.
121. Saunders ME. Insect pollinators collect pollen from wind-pollinated plants: implications for pollination ecology and sustainable agriculture. *Insect Conserv Diver.* 2018;11:13–31. <https://doi.org/10.1111/icaad.12243>.
- 122.●● Urban-Mead KR, van Dyke M, Muniz P, Young A, Danforth BN, McArt SH. Early spring orchard pollinators spill over from resource-rich adjacent forest patches. *J Appl Ecol.* 2023;60:553–64. **Provides evidence that bees collect pollen from broadleaf trees.**
123. Tucker SK, Ginsberg HS, Alm SR. Eastern carpenter bee (Hymenoptera: Apidae): Nest structure, nest cell provisions, and trap nest acceptance in Rhode Island. *Environ Entomol.* 2019;48:702–10. <https://doi.org/10.1093/ee/nvz032>.
124. Pound MJ, Vinkenoog R, Hornby S, Benn J, Goldberg S, Keating B, et al. Determining if honey bees (*Apis mellifera*) collect pollen from anemophilous plants in the UK. *Palynology.* 2023;47:2154867. <https://doi.org/10.1080/01916122.2022.2154867>.
125. Pernal SF, Currie RW. Pollen quality of fresh and 1-year-old single pollen diets for worker honey bees (*Apis mellifera* L.). *Apidologie.* 2000;31:387–409. <https://doi.org/10.1051/apido:2000130>.
126. Ulyshen MD, Robertson KM, Horn S, Dixon CM. Characterization of the bee community and pollination network in a southeastern U.S. pine savanna. *Front Ecol Evol.* 2024;12:1403602. <https://doi.org/10.3389/fevo.2024.1403602>.
127. Guo P-F, Wang M-Q, Orr M, Li Y, Chen J-T, Zhou Q-S, et al. Tree diversity promotes predatory wasps and parasitoids but not pollinator bees in a subtropical experimental forest. *Basic Appl Ecol.* 2021;53:134–42. <https://doi.org/10.1016/j.baae.2021.03.007>.
- 128.● Allen G, Davies RG. Canopy sampling reveals hidden potential value of woodland trees for wild bee assemblages. *Insect Conserv Diver.* 2023;16(1):33–46. <https://doi.org/10.1111/icaad.12606>. **Provides evidence that broadleaf forest canopies are important for bees.**
129. Sobek S, Steffan-Dewenter I, Scherber C, Tschardt T. Spatiotemporal changes of beetle communities across a tree diversity gradient. *Divers Distrib.* 2009;15(4):660–70. <https://doi.org/10.1111/j.1472-4642.2009.00570.x>.
130. McLaughlin R, Keller J, Wagner E, Biddinger D, Grozinger C, Hoover K. Insect visitors of black cherry (*Prunus serotina*) (Rosales: Rosaceae) and factors affecting viable seed production. *Environ Entomol.* 2022;51:471–81. <https://doi.org/10.1093/ee/nvab141>.
131. Van Holt T, Binford MW, Portier KM, Vergara R. A stand of trees does not a forest make: Tree plantations and forest transitions. *Land Use Policy.* 2016;56:147–57. <https://doi.org/10.1016/j.landusepol.2016.04.015>.
132. Pérez-Gómez A, Godoy O, Ojeda F. Beware of trees: Pine afforestation of a naturally treeless habitat reduces flower and pollinator diversity. *Glob Ecol Conserv.* 2024;50:e02808. <https://doi.org/10.1016/j.gecco.2024.e02808>.

133. Kanninen M. Plantation forests: Global perspectives. In: Bauhus J, van der Meer P, Kanninen M, editors. Ecosystem goods and services from plantation forests. London, UK: Earthscan; 2010. p. 1–15.
134. Wang C, Zhang W, Li X, Wu J. A global meta-analysis of the impacts of tree plantations on biodiversity. *Glob Ecol Biogeogr*. 2022;31(3):576–87. <https://doi.org/10.1111/geb.13440>.
135. Moncur MW, Mitchell A, Frupp Y, Kleinschmidt GJ. The role of honey bees (*Apis mellifera*) in eucalypt and acacia seed production areas. *Commonw Forest Rev*. 1995;74(4):350–4.
136. Wu P, Axmacher JC, Li X, Song X, Yu Z, Xu H, et al. Contrasting effects of natural shrubland and plantation forests on bee assemblages at neighboring apple orchards in Beijing China. *Biol Conserv*. 2019;237:456–62. <https://doi.org/10.1016/j.biocon.2019.07.029>.
137. Vasconcelos RN, Cambui ECB, Mariano-Neto E, da Rocha PLB, Cardoso MZ. The role of Eucalyptus planted forests for fruit-feeding butterflies' conservation in fragmented areas of the Brazilian Atlantic forest. *Forest Ecol Manag*. 2019;432:115–20. <https://doi.org/10.1016/j.foreco.2018.09.017>.
138. Aguiar WMD, Sofia SH, Melo GAR, Gaglianone MC. Changes in orchid bee communities across forest-agroecosystem boundaries in Brazilian Atlantic forest landscapes. *Environ Entomol*. 2015;44(6):1465–71. <https://doi.org/10.1093/ee/nvv130>.
139. van Halder I, Barbaro L, Corcket E, Jactel H. Importance of semi-natural habitats for the conservation of butterfly communities in landscapes dominated by pine plantations. In: Brockerhoff EG, Jactel H, Parrotta JA, Quine CP, Sayer J, Hawksworth DL, editors. *Plantation Forests and Biodiversity: Oxymoron or Opportunity?* Dordrecht: Springer, Netherlands; 2009. p. 225–45.
140. Gittings T, O'Halloran J, Kelly T, Giller PS. The contribution of open spaces to the maintenance of hoverfly (Diptera, Syrphidae) biodiversity in Irish plantation forests. *Forest Ecol Manag*. 2006;237(1):290–300. <https://doi.org/10.1016/j.foreco.2006.09.052>.
141. Straw NA, Williams DT, Fielding NJ, Jukes M, Connolly T, Forster J. The influence of forest management on the abundance and diversity of hoverflies in commercial plantations of Sitka spruce: The importance of sampling in the canopy. *Forest Ecol Manag*. 2017;406:95–111. <https://doi.org/10.1016/j.foreco.2017.06.010>.
142. Taki H, Inoue T, Tanaka H, Makiyama H, Sueyoshi M, Isono M, et al. Responses of community structure, diversity, and abundance of understory plants and insect assemblages to thinning in plantations. *Forest Ecol Manag*. 2010;259(3):607–13. <https://doi.org/10.1016/j.foreco.2009.11.019>.
143. Harrington TB, Edwards MB. Understory vegetation, resource availability, and litterfall responses to pine thinning and woody vegetation control in longleaf pine plantations. *Can J Forest Res*. 1999;29(7):1055–64. <https://doi.org/10.1139/x99-118>.
144. Larue C, Austruy E, Basset G, Petit RJ. Revisiting pollination mode in chestnut (*Castanea* spp): An integrated approach. *Bot Lett*. 2021;168(3):348–72. <https://doi.org/10.1080/23818107.2021.1872041>.
145. Totland Ø, Nielsen A, Bjercknes A-L, Ohlson M. Effects of an exotic plant and habitat disturbance on pollinator visitation and reproduction in a boreal forest herb. *Am J Bot*. 2006;93(6):868–73. <https://doi.org/10.3732/ajb.93.6.868>.
146. Tepedino VJ, Brosi AB, Griswold TL. Might flowers of invasive plants increase native bee carrying capacity? Intimations from Capitol Reef National Park. *Utah Nat Area J*. 2008;28(1):44–50. [https://doi.org/10.3375/0885-8608\(2008\)28\[44:MFOIPI\]2.0.CO;2](https://doi.org/10.3375/0885-8608(2008)28[44:MFOIPI]2.0.CO;2).
147. McKinney AM, Goodell K. Shading by invasive shrub reduces seed production and pollinator services in a native herb. *Biol Inv*. 2010;12(8):2751–63. <https://doi.org/10.1007/s10530-009-9680-4>.
148. Oliver CD, Larson BC. *Forest stand dynamics*. New York: John Wiley & Sons, Inc.; 1996.
149. Ulyshen MD, Pokswinski S, Hiers JK. A comparison of bee communities between primary and mature secondary forests in the longleaf pine ecosystem. *Sci Rep*. 2020;10(1):2916. <https://doi.org/10.1038/s41598-020-59878-4>.
150. Ealy N, Pawelek J, Hazlehurst J. Effects of forest management on native bee biodiversity under the tallest trees in the world. *Ecol Evol*. 2023;13(7):e10286. <https://doi.org/10.1002/ece3.10286>.
151. Hilmers T, Friess N, Bässler C, Heurich M, Brandl R, Pretzsch H, et al. Biodiversity along temperate forest succession. *J Appl Ecol*. 2018;55:2756–66. <https://doi.org/10.1111/1365-2664.13238>.
152. Mathis CL, McNeil DJ, Lee MR, Grozinger CM, King DI, Otto CRV, et al. Pollinator communities vary with vegetation structure and time since management within regenerating timber harvests of the Central Appalachian Mountains. *Forest Ecol Manag*. 2021;496:119373. <https://doi.org/10.1016/j.foreco.2021.119373>. **Reports changes in pollinator diversity in early successional broadleaf forests.**
153. Roubik DW. Stingless bee nesting biology. *Apidologie*. 2006;37(2):124–43. <https://doi.org/10.1051/apido:2006026>.
154. Cunningham-Minnick MJ, Roberts HP, Milam J, King DI. Sampling the understory, midstory, and canopy is necessary to fully characterize native bee communities of temperate forests and their dynamic environmental relationships. *Front Ecol Evol*. 2024;12:1352266. <https://doi.org/10.3389/fevo.2024.1352266>. **Shows changes in bee composition with height in broadleaf forests.**
155. Bopp C, Engler A, Jara-Rojas R, Arriagada R. Are forest plantation subsidies affecting land use change and off-farm income? A farm-level analysis of Chilean small forest landowners. *Land Use Policy*. 2020;91:104308. <https://doi.org/10.1016/j.landusepol.2019.104308>.
156. Connell JH. Diversity in tropical rain forests and coral reefs. *Science*. 1978;199(4335):1302–10. <https://doi.org/10.1126/science.199.4335.1302>.
157. Messier C, Posada J, Aubin I, Beaudet M. Functional relationships between old-growth forest canopies, understory light and vegetation dynamics. In: Wirth C, Gleixner G, Heimann M, editors. *Old-growth forests: Function, fate and value*. Berlin, Heidelberg: Springer Berlin Heidelberg; 2009. p. 115–39.
158. DellaSala DA, Mackey B, Norman P, Campbell C, Comer PJ, Kormos CF, et al. Mature and old-growth forests contribute to large-scale conservation targets in the conterminous United States. *Front For Glob Change*. 2022;5:979528. <https://doi.org/10.3389/ffgc.2022.979528>.
159. Roubik DW. Nest and colony characteristics of stingless bees from Panamá (Hymenoptera: Apidae). *J Kansas Entomol Soc*. 1983;56(3):327–55.
160. Salmah S, Inoue T, Sakagami SF. An analysis of apid bee richness (Apidae) in central Sumatra. In: Ohgushi R, Sakagami SF, Roubik DW, editors. *Natural history of social wasps and bees in equatorial Sumatra*. Sapporo: Hokkaido University Press; 1990. p. 139–74.
161. De Schuyter W, De Lombaerde E, Depauw L, De Smedt P, Stachurska-Swakoń A, Orczewska A, et al. Declining potential nectar production of the herb layer in temperate forests under global change. *J Ecol*. 2024;112(4):832–47. <https://doi.org/10.1111/1365-2745.14274>.
162. Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. Global pollinator declines: Trends, impacts and drivers. *Trends Ecol Evol*. 2010;25(6):345–53. <https://doi.org/10.1016/j.tree.2010.01.007>.

163. Ulyshen MD, Horn S. Declines of bees and butterflies over 15 years in a forested landscape. *Curr Biol.* 2023;33:1346–50.
164. Jha S, Vandermeer JH. Impacts of coffee agroforestry management on tropical bee communities. *Biol Conserv.* 2010;143(6):1423–31. <https://doi.org/10.1016/j.biocon.2010.03.017>.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.