

Table 2. Taxa of *Chamaecyparis*, *Cupressus*, and *Platycladus* reported as hosts of *Arceuthobium oxycedri*.

Preferred name	Name cited	Natural range	Country of report	Reference ^a
1. <i>Chamaecyparis funebris</i> (Chinese weeping cypress)	<i>Cupressus funebris</i>	China	Ukraine (Crimea)	36
2. <i>Chamaecyparis thyoides</i> (Atlantic white-cedar)	<i>Chamaecyparis sphaeroides</i> var. <i>pendula</i> <i>Chamaecyparis thyoides</i>	Atlantic coast, USA	Central Europe Croatia	10 ^b , 33 ^c 63
1. <i>Cupressus arizonica</i> (Arizona cypress)	<i>Cupressus arizonica</i> <i>Cupressus arizonica</i>	Southwest USA	Spain Ukraine (Crimea)	58 36
2. <i>Cupressus lusitanica</i>	<i>Cupressus lusitanica</i>	Portugal ^d	Ukraine (Crimea)	36
3. <i>Cupressus benthamii</i>	<i>Cupressus lusitanica</i> var. <i>benthamii</i>	Mexico	Ukraine (Crimea)	36
4. <i>Cupressus macnabiana</i> (MacNab cypress)	<i>Cupressus macnabiana</i>	California, USA	Ukraine (Crimea)	36
5. <i>Cupressus macrocarpa</i> (Monterey cypress)	<i>Cupressus macrocarpa</i>	California, USA	Ukraine (Crimea)	36, 75
1. <i>Platycladus orientalis</i> (Oriental arbor-vitae)	<i>Biota orientalis</i> <i>Platycladus orientalis</i>	North and west China and Korea	Central Europe Ukraine (Crimea)	10 ^b 36

^aSee [References](#) and [Methods](#).

^bRecord based on transmission by graft from *Juniperus communis*.

^cRecord based on artificial inoculation.

^dRecent analysis of leaf essential oils and DNA fingerprinting of planted *C. lusitanica* near Bussaco, Portugal indicate that it is different from populations of *Cupressus* in Mexico and Central America, previously believed to be the origin of this plantation (Adams et al 1997).