

Long Distance Movement by a Coyote within the Rocky Mountains

Abstract

We documented a long distance movement of a juvenile male coyote (*Canis latrans*) between February 2002 and February 2003. The radiocollared coyote, last located in west central Montana, U.S.A. traveled ≥ 310 km to southwestern Alberta, Canada, where it was trapped. This is the longest documented movement by a coyote in western North America.

A juvenile male coyote (*Canis latrans*) was trapped and radiocollared near Seeley Lake, Montana on September 16, 2001 as one of 25 animals marked during a study of coyote-winter ecology. The coyote left the study area November 20, 2001 and the radio signal was lost on February 21, 2002. This coyote was killed by a fur trapper near Crowsnest Pass, Alberta February 2, 2003 after traveling a straight line distance of ≥ 310 km from its last recorded location. To our knowledge, this is the longest documented movement by a coyote in western North America.

The coyote was captured in a modified Victor #3 Softcatch[®] (Oneida Victor Inc. Ltd., Euclid,

OH) coil spring trap by a front foot, anesthetized, examined, and fitted with a radio collar. It was uninjured and apparently healthy when released. The animal remained within 20 km of the capture location until late November 2001 when it left the study area. In February 2003, this male was trapped in good condition along with a female coyote; both animals were killed by the trapper (J. Clarke, Alberta Environment, personal communication).

Although this movement is exceptional for western North America, movements of equal or greater lengths have been documented within the central plains and eastern forest regions (Table 1).

TABLE 1. Long distance movements by coyotes documented in North America.

Region	Movement Distance (km)	Location ^a	Source
West ^b	310	Montana, USA./ Alberta, Can.	Kolbe and Squires (This article)
	58	Alberta, Can.	Roy and Dorance (1985)
	152	Montana, U.S.A.	Pyrah (1984)
	57	Idaho, U.S.A.	Woodruff and Keller (1982)
	85	Montana, U.S.A.	R. Crabtree (pers. comm.)
	155	Alberta, Can.	Nellis and Keith (1976)
	140	California, USA	Hawthorn (1971)
	185	Montana, USA	Robinson and Cummings (1951)
	161	Wyoming, USA.	Garlough (1940)
Central ^c	544	Manitoba/ Saskatchewan, Can.	Carbyn and Paquet (1986)
	323	Iowa/Missouri, USA	Andrews and Boggess (1978)
East ^d	320	Ontario, Can.	Rosatte (2002)
	342	Maine/Vermont, USA	Harrison (1992)

^a Locations listed as "moved from/moved to" where applicable

^b "West" is defined as having an initial capture location west of 110°W Longitude

^c "Central" is defined as having an initial capture location between 90°W and 110°W Longitude

^d "East" is defined as having an initial capture location east of 90°W Longitude

¹ Author to whom correspondence should be addressed.

E-mail: jaykolbe@hotmail.com

This coyote was initially captured at an elevation of 1470m and trapped at an elevation of roughly 1700m. Although the movement path on this male was not documented, the area between these two points is mostly mountainous terrain with elevations varying between 1100m to over 3100m.

Long-distance movements of coyotes are apparently rare. However, biologists should consider the maximum regional dispersal distance when

managing rabies, sarcoptic mange and other disease epizootics (Rosatte 2002).

Acknowledgements

We thank R. Hendrickson for reporting his capture of the marked coyote in Alberta. We also thank J. Clarke of Alberta Environment for his assistance in documenting the capture location and returning the collar to us for analysis.

Literature Cited

- Andrews, R. D., E. K. Boggess. 2001. Ecology of Coyotes in Iowa. Pages 249-265 *In* M. Bekoff (editor). *Coyotes: Biology, Behavior, and Management*. The Blackburn Press, Caldwell, New Jersey.
- Carbyn, L. N., and P. C. Paquet. 1986. Long distance movement of a coyote from Riding Mountain National Park. *Journal of Wildlife Management* 50: 89.
- Garlough, F. E. 1940. Study of the Migratory Habits of Coyotes. Denver Wildlife Research Lab. Denver, Colorado. (Mimeo.)
- Harrison, D. J., 1992. Dispersal characteristics of juvenile coyotes in Maine. *Journal of Wildlife Management* 56: 128-138.
- Hawthorn, V. M. 1971. *California Fish and Game* 57: 154-161.
- Nellis, C. H., and L. B. Keith. 1976. Population dynamics of coyotes in central Alberta, 1964-1968. *Journal of Wildlife Management* 40: 389-399.
- Pyrah, D., 1984. Social distribution and population estimates of coyotes in north-central Montana. *Journal of Wildlife Management* 48: 679-690.
- Robinson, W. B., and M. W. Cummings. 1951. U. S. Fish and Wildlife Service Special Scientific Report: Wildlife Number 11.
- Rosatte, R. C., 2002. Long distance movement by a coyote, *Canis latrans*, and red fox, *Vulpes vulpes*, in Ontario: implications for disease spread. *The Canadian Field Naturalist* 116: 129-131.
- Roy, L. D., and M. J. Dorrance. 1985. Coyote movements, habitat use, and vulnerability in central Alberta. *Journal of Wildlife Management* 49: 307-313.
- Woodruff, R. A., and B. L. Keller. 1982. Dispersal, daily activity, and home range of coyotes, *Canis latrans*, in southeastern Idaho, U.S.A. *Northwest Science* 56: 199-207.

Received 11 November 2004

Accepted for Publication 4 December 2004