

Environmental DNA particle size distribution from Brook Trout (*Salvelinus fontinalis*)

Taylor M. Wilcox^{1,2} · Kevin S. McKelvey² · Michael K. Young² · Winsor H. Lowe¹ · Michael K. Schwartz³

Received: 10 December 2014 / Accepted: 8 April 2015
© Springer Science+Business Media Dordrecht (outside the USA) 2015

Abstract Environmental DNA (eDNA) sampling has become a widespread approach for detecting aquatic animals with high potential for improving conservation biology. However, little research has been done to determine the size of particles targeted by eDNA surveys. In this study, we conduct particle distribution analysis of eDNA from a captive Brook Trout (*Salvelinus fontinalis*) in a flow-through mesocosm. Our data suggest that 1.2–10- μ m particles are the most common size of eDNA from Brook Trout, which is consistent with our hypotheses that eDNA in the environment is comprised of loosely aggregated smaller particles, resulting in high inter-sample heterogeneity. These findings are similar to those of a study on Common Carp (*Cyprinus carpio*) in lentic systems.

Keywords Detection · eDNA · Mitochondria · Fish

Sampling water for environmental DNA (eDNA) allows for detection of aquatic organisms without ever seeing a live individual (Lodge et al. 2012). These approaches may be more sensitive and cost-effective than traditional sampling methods, making eDNA a potentially powerful tool for conservation biology. eDNA has recently been applied

to detect rare invasive species (e.g., Jerde et al. 2011) and endangered native species (e.g., Biggs et al. 2015) including fishes (Takahara et al. 2013) amphibians (Pilliod et al. 2013), reptiles (Piaggio et al. 2013), mollusks (Goldberg et al. 2013), arthropods (Mächler et al. 2014), and mammals (Foote et al. 2012). However, to fulfill its potential as a conservation tool, we need to learn more about those factors that affect both the production and persistence of eDNA in complex aquatic environments.

A key example is the particle size of eDNA, which has not been determined empirically for most species, but has been attributed in the literature to everything from extracellular DNA to sloughed tissues (e.g., Dejean et al. 2011). DNA of different particle sizes may degrade or be deposited at different rates in the environment, and eDNA sampling approaches (e.g., precipitation vs. filtration) vary in the size of particles that they target. As a result, determining the size of eDNA particles sampled for species detection is a critical step in understanding eDNA dynamics and capture.

Turner et al. (2014) sought to determine the size of eDNA particles from Common Carp (*Cyprinus carpio*) in lentic environments using sequential filtration of environmental samples. Particles 1–10 μ m across, corresponding to the size of individual mitochondria or small cells, constituted the most common size class of Common Carp eDNA (Turner et al. 2014). Here, we present the results of a comparable experiment to determine the particle size distribution of eDNA from Brook Trout (*Salvelinus fontinalis*) under different environmental conditions (cold, lotic mesocosms). Brook Trout are a species of special conservation concern within their native range (Hudy et al. 2008) and an important invasive species globally (e.g., Dunham et al. 2002; Bosch et al. 2006). Further, studies of Brook Trout eDNA may be directly transferable to eDNA studies

✉ Taylor M. Wilcox
taylor.wilcox@umontana.edu

¹ Division of Biological Sciences, University of Montana, 32 Campus Drive, Missoula, MT, USA

² U.S. Forest Service, Rocky Mountain Research Station, 800 East Beckwith Avenue, Missoula, MT, USA

³ United States Department of Agriculture, Forest Service, National Genomics Center for Wildlife and Fish Conservation, Rocky Mountain Research Station, 800 East Beckwith Avenue, Missoula, MT, USA

of other salmonids of conservation concern (e.g., Laramie et al. 2014).

In August 2014 a single Brook Trout was captured by angling in West Fork Lolo Creek, MT, USA (N 46.687168°, W 114.557371°) and placed in a flow-through mesocosm (0.35 m × 0.5 m × 0.22 m) fed by a fishless, ephemeral tributary. This tributary was confirmed free of Brook Trout DNA by sampling 1 month and again several days prior to the experiment (5-L samples; see protocol below). Twenty-four hours after fish addition, we collected five sequential 0.5-L samples in disposable polyethylene bottles from the tank outflow. The fish was then returned to its capture site.

For each 0.5-L sample, 100 mL was sequentially filtered through four filter pore sizes from largest to smallest. This filter series included three 47-mm-diameter polycarbonate track-etched (PCTE) filters (0.4, 1.2, and 10- μ m pores) and one 47-mm-diameter nylon net filter (60- μ m pores; no PCTE filter was available in this size; Millipore). We did not assess eDNA particles smaller than 0.4 μ m because, in a pilot study, long filtration times made smaller pore sizes impractical for many field applications (≥ 1 h per filter for 1–5 L sample volumes). As a result, we may not have detected particles of eDNA of smaller sizes, such as extracellular DNA. Filters were held in disposable filter holder funnels (Fisher Scientific) attached to silicone housing (Masterflex) and were pumped using a peristaltic pump (GeoTech Inc.,) so that the outflow was captured by the next filter in the sequence. New filter holders and hoses were used between each sample. We also filtered 100 mL of distilled water through a sequence of filters as an equipment control. Using new forceps for each sample, filters were folded in half and then rolled into sterile 1.5-mL tubes and stored on ice until arrival at the lab (within 6 h), where they were stored in a -20 °C freezer until extraction. All samples were filtered in the field within 2 h of collection from the mesocosm tank.

We extracted DNA from these samples and equipment controls using the DNeasy Blood and Tissue Kit and QIAshredder columns (Qiagen; protocol adapted from Goldberg et al. 2013; final elution into 100 μ L sterile TE; Integrated DNA Technologies; IDT). All extractions were performed in a room reserved for extracting non-invasive genetic samples where no PCR products or other sources of high concentration DNA are handled. Extracted samples were stored in a -20 or -80 °C freezer until qPCR analysis.

We used a species-specific TaqMan-MGB assay (*BRK2*; Wilcox et al. 2013) to estimate Brook Trout mitochondrial DNA (mtDNA) concentrations in all samples and equipment controls. Experiments were performed in 15- μ L volumes with 4 μ L of template DNA and final concentrations of 1 × TaqMan Environmental Mastermix 2.0 (Life

Technologies), a 1 × assay mix (primers each at 900 nM, probe at 250 nM), and a VIC-labeled exogenous internal amplification control (Life Technologies TaqMan Exogenous Control Kit). We used cycling conditions of 95 °C/10 min (95 °C/15 s, 60 °C/60 s) × 45 cycles on a StepOne Plus Real-time PCR Instrument (Life Technologies). The internal amplification control acted as a test for PCR inhibitors in the samples, as evidence by a >1 cycle threshold (C_t) shift in the amplification curve relative to that in a no template control. There was no evidence of inhibition in this study.

The qPCR plate also included a negative PCR setup control and a series of five known-quantity target standards diluted to 6250, 1250, 250, 50 and 10 copies/reaction. This standard was composed of a linearized, synthetic gene from IDT containing the 139-bp sequence of interest (Wilcox et al. 2013). Samples were quantified by comparison with the standard curve ($r^2 = 0.93$, efficiency = 99.6 %) using the C_{y0} method, which is less sensitive to low levels of PCR inhibitors than the C_t method (Guescini et al. 2008). C_{y0} values were calculated from raw amplification data using the package *qpcR* (Ritz and Spiess 2008) in R v. 3.0.1 (R Core Team 2013). All standards, controls, and samples were run in triplicate and copy number is reported as the mean estimate for all triplicate wells.

There was no amplification in any of the equipment or PCR setup controls; there was amplification for all 60 sample reactions (20 samples run in triplicate). For each sequential filtration replicate, mtDNA copy number for all four samples was summed (median = 24×10^6 , range = $14\text{--}41 \times 10^6$ copies/L), and the copy number estimates from each filter were divided by this total to report the proportion of eDNA in each size class (Fig. 1).




Fig. 1 Bars show the mean proportion of the brook trout eDNA that comes from each size class (0.4–1.2, 1.2–10, 10–60, and ≥ 60 μ m) for each of five water samples (individual samples shown with white points). As in Turner et al. (2014), particles approximately 1–10 μ m across are the most abundant source of the fish eDNA

Our results were generally consistent with the findings of Turner et al. (2014) in that particles 1.2–10 μm were the largest source of Brook Trout eDNA. This suggests that this particle size is likely to be the most useful to target for capture in the field (e.g., via in situ filtration). Also, as in Turner et al. (2014), we observed high inter-sample variability (the maximum DNA concentration was almost 3 \times the minimum in our five samples), which is surprising because the particle sizes captured by filtration suggest that the eDNA is dominated by individual cells or mitochondria. In Turner et al. (2014) and in this study, a well-mixed solution containing many independent mitochondria or cells should exhibit low intra-sample variability. A parsimonious explanation for this apparent discrepancy is that larger particles were captured, but were broken apart with filtration. This hypothesis—that fish eDNA is transported primarily in weak aggregations that are subject to breakdown—is also consistent with measured eDNA losses in lotic systems (Jane et al. 2015) and high inter-sample heterogeneity observed in other field studies (Pilliod et al. 2013). Jane et al. (2015) observed relatively rapid downstream Brook Trout eDNA losses in streams during low flows, which is consistent with rapid settling expected by larger particle sizes (e.g., Maggi 2013), but also found that small quantities of eDNA were transported distantly downstream, which is consistent with small particle sizes. Previous studies on the physical transport of other organic particles in streams could help us better understand the behavior of fish DNA in the environment, further informing sampling design.

Acknowledgments TW is supported by a NSF Graduate Research Fellowship (Grant No. DGE-1313190). This work was done in accordance with Montana Fish, Wildlife and Parks Scientific Collectors Permit #SCP-33-14. This manuscript was improved by comments from Kellie Carim, Katie Zarn, and two anonymous reviewers.

References

- Biggs J, Ewald N, Valentini A et al (2015) Using eDNA to develop a national citizen science-based monitoring programme for the great crested newt (*Triturus cristatus*). *Biol Conserv* 183:19–28. doi:10.1016/j.biocon.2014.11.029
- Bosch J, On PAR, Boyero L, Martinez-Solano I (2006) Effects of introduced salmonids on a montane population of Iberian frogs. *Conserv Biol* 20:180–189. doi:10.1111/j.1523-1739.2006.00296.x
- Dejean T, Valentini A, Duparc A et al (2011) Persistence of environmental DNA in freshwater ecosystems. *PLoS One* 6:e23398. doi:10.1371/journal.pone.0023398
- Dunham JB, Adams SB, Schroeter RE, Novinger DC (2002) Alien invasions in aquatic ecosystems: toward an understanding of brook trout invasions and potential impacts on inland cutthroat trout in western North America. *Rev Fish Biol Fish*. doi:10.1023/a:1025338203702
- Foote AD, Thomsen PF, Sveegaard S et al (2012) Investigating the potential use of environmental DNA (eDNA) for genetic monitoring of marine mammals. *PLoS One* 7:e41781. doi:10.1371/journal.pone.0041781
- Goldberg CS, Sepulveda A, Ray A et al (2013) Environmental DNA as a new method for early detection of New Zealand mudsnails (*Potamopyrgus antipodarum*). *Freshw Sci* 32:792–800. doi:10.1899/13-046.1
- Guescini M, Sisti D, Rocchi MBL et al (2008) A new real-time PCR method to overcome significant quantitative inaccuracy due to slight amplification inhibition. *BMC Bioinform* 9:326. doi:10.1186/1471-2105-9-326
- Hudy M, Thieling TM, Gillespie N, Smith EP (2008) Distribution, status, and land use characteristics of subwatersheds within the native range of Brook Trout in the eastern United States. *North Am J Fish Manag* 28:1069–1085. doi:10.1577/M07-017.1
- Jane SF, Wilcox TM, Mckelvey KS et al (2015) Distance, flow, and PCR inhibition: eDNA dynamics in two headwater streams. *Mol Ecol Resour* 15:216–227
- Jerde CL, Mahon AR, Chadderton WL, Lodge DM (2011) “Sight-unseen” detection of rare aquatic species using environmental DNA. *Conserv Lett* 4:150–157. doi:10.1111/j.1755-263X.2010.00158.x
- Laramie MB, Pilliod DS, Goldberg CS (2014) Characterizing the distribution of an endangered salmonid using environmental DNA analysis. *Biol Conserv*. doi:10.1016/j.biocon.2014.11.025
- Lodge DM, Turner CR, Jerde CL et al (2012) Conservation in a cup of water: estimating biodiversity and population abundance from environmental DNA. *Mol Ecol* 21:2555–2558. doi:10.1111/j.1365-294X.2012.05600.x
- Mächler E, Deiner K, Steinmann P, Altermatt F (2014) Utility of environmental DNA for monitoring rare and indicator macroinvertebrate species. *Freshw Sci* 33:000. doi:10.1086/678128
- Maggi F (2013) The settling velocity of mineral, biomineral, and biological particles and aggregates in water. *J Geophys Res Ocean* 118:2118–2132. doi:10.1002/jgrc.20086
- Piaggio AJ, Engeman RM, Hopken MW et al (2013) Detecting an elusive invasive species: a diagnostic PCR to detect Burmese python in Florida waters and an assessment of persistence of environmental DNA. *Mol Ecol Resour*. doi:10.1111/1755-0998.12180
- Pilliod DS, Goldberg CS, Arkle RS, Waits LP (2013) Estimating occupancy and abundance of stream amphibians using environmental DNA from filtered water samples. *Can J Fish Aquat Sci* 70:1123–1130
- R Core Team (2013) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>
- Ritz C, Spiess A-N (2008) qpcR: an R package for sigmoidal model selection in quantitative real-time polymerase chain reaction analysis. *Bioinformatics* 24:1549–1551. doi:10.1093/bioinformatics/btn227
- Takahara T, Minamoto T, Doi H (2013) Using environmental DNA to estimate the distribution of an invasive fish species in ponds. *PLoS One* 8:e56584. doi:10.1371/journal.pone.0056584
- Turner CR, Barnes MA, Xu CCY et al (2014) Particle size distribution and optimal capture of aqueous microbial eDNA. *Methods Ecol Evol* 5:676–684. doi:10.1101/001941
- Wilcox TM, McKelvey KS, Young MK et al (2013) Robust detection of rare species using environmental DNA: the importance of primer specificity. *PLoS One* 8:e59520. doi:10.1371/journal.pone.0059520